

Birational classification of varieties

James M^cKernan

UCSB

A little category theory

- The most important part of any category \mathcal{C} are the **morphisms** not the **objects**.

A little category theory

- The most important part of any category \mathcal{C} are the **morphisms** not the **objects**.
- It is the aim of higher dimensional geometry to classify algebraic varieties up to birational equivalence.

A little category theory

- The most important part of any category \mathcal{C} are the **morphisms** not the **objects**.
- It is the aim of higher dimensional geometry to classify algebraic varieties up to birational equivalence.
- Thus the objects are algebraic varieties, but what are the morphisms?

Contraction mappings

- Well, given any morphism $f: X \rightarrow Y$ of normal algebraic varieties, we can always factor f as $g: X \rightarrow W$ and $h: W \rightarrow Y$, where h is finite and g has connected fibres.

Contraction mappings

- Well, given any morphism $f: X \longrightarrow Y$ of normal algebraic varieties, we can always factor f as $g: X \longrightarrow W$ and $h: W \longrightarrow Y$, where h is finite and g has connected fibres.
- Mori theory does **not** say much about finite maps.

Contraction mappings

- Well, given any morphism $f: X \longrightarrow Y$ of normal algebraic varieties, we can always factor f as $g: X \longrightarrow W$ and $h: W \longrightarrow Y$, where h is finite and g has connected fibres.
- Mori theory does **not** say much about finite maps.
- It **does** have a lot to say about morphisms with connected fibres.

Contraction mappings

- Well, given any morphism $f: X \longrightarrow Y$ of normal algebraic varieties, we can always factor f as $g: X \longrightarrow W$ and $h: W \longrightarrow Y$, where h is finite and g has connected fibres.
- Mori theory does **not** say much about finite maps.
- It **does** have a lot to say about morphisms with connected fibres.
- In fact any morphism $f: X \longrightarrow Y$ such that $f_* \mathcal{O}_X = \mathcal{O}_Y$ will be called a **contraction morphism**. If X and Y are normal, this is the same as requiring the fibres of f to be connected.

Curves versus divisors

- So we are interested in the category of algebraic varieties (primarily normal and projective), and contraction morphisms, and we want to classify all contraction morphisms.

Curves versus divisors

- So we are interested in the category of algebraic varieties (primarily normal and projective), and contraction morphisms, and we want to classify all contraction morphisms.
- Traditionally the approved way to study a projective variety is to embed it in projective space, and consider the family of hyperplane sections.

Curves versus divisors

- So we are interested in the category of algebraic varieties (primarily normal and projective), and contraction morphisms, and we want to classify all contraction morphisms.
- Traditionally the approved way to study a projective variety is to embed it in projective space, and consider the family of hyperplane sections.
- In Mori theory, we focus on **curves**, not **divisors**.

Curves versus divisors

- So we are interested in the category of algebraic varieties (primarily normal and projective), and contraction morphisms, and we want to classify all contraction morphisms.
- Traditionally the approved way to study a projective variety is to embed it in projective space, and consider the family of hyperplane sections.
- In Mori theory, we focus on **curves**, not **divisors**.
- In fact a contraction morphism $f: X \longrightarrow Y$ is determined by the curves which it contracts. Indeed Y is clearly determined topologically, and the condition $\mathcal{O}_Y = f_* \mathcal{O}_X$ determines the algebraic structure.

The closed cone of curves

- $\text{NE}(X)$ denotes the **cone of effective curves** of X , the **closure** of the image of the effective curves in $H_2(X, \mathbb{R})$, considered as a cone inside the span.

The closed cone of curves

- $\text{NE}(X)$ denotes the **cone of effective curves** of X , the **closure** of the image of the effective curves in $H_2(X, \mathbb{R})$, considered as a cone inside the span.
- By Kleiman's criteria, any divisor H is ample iff it defines a positive linear functional on

$$\begin{aligned} \text{NE}(X) - \{0\} &\quad \text{by} \\ [C] &\longrightarrow H \cdot C. \end{aligned}$$

The closed cone of curves

- $\text{NE}(X)$ denotes the **cone of effective curves** of X , the **closure** of the image of the effective curves in $H_2(X, \mathbb{R})$, considered as a cone inside the span.
- By Kleiman's criteria, any divisor H is ample iff it defines a positive linear functional on

$$\begin{aligned} \text{NE}(X) - \{0\} &\quad \text{by} \\ [C] &\longrightarrow H \cdot C. \end{aligned}$$

- Given f , set $D = f^*H$, where H is an ample divisor on Y . Then D is **nef**, that is $D \cdot C \geq 0$, for every curve C .

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$.
Moreover the set of such curves is a **face** of $\text{NE}(X)$.

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$. Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$.
Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the
 - faces F of $\text{NE}(X)$ and the

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$.
Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the
 - faces F of $\text{NE}(X)$ and the
 - contraction morphisms f .

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$.
Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the
 - faces F of $\text{NE}(X)$ and the
 - contraction morphisms f .
- So, which faces F correspond to contractions f ?
Similarly which divisors are the pullback of ample divisors?

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$. Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the
 - faces F of $\text{NE}(X)$ and the
 - contraction morphisms f .
- So, which faces F correspond to contractions f ? Similarly which divisors are the pullback of ample divisors?
- We say that a divisor D is **semiample** if $D = f^*H$, for some contraction morphism f and ample divisor H .

Semiample divisors

- Then a curve C is contracted by f iff $D \cdot C = 0$. Moreover the set of such curves is a **face** of $\text{NE}(X)$.
- Thus there is **partial** correspondence between the
 - faces F of $\text{NE}(X)$ and the
 - contraction morphisms f .
- So, which faces F correspond to contractions f ? Similarly which divisors are the pullback of ample divisors?
- We say that a divisor D is **semiample** if $D = f^*H$, for some contraction morphism f and ample divisor H .
- Note that if D is semiample, it is certainly nef.

An easy example

- Suppose that $X = \mathbb{P}^1 \times \mathbb{P}^1$.

An easy example

- Suppose that $X = \mathbb{P}^1 \times \mathbb{P}^1$.
- $\text{NE}(X)$ sits inside a two dimensional vector space. The cone is spanned by $f_1 = [\mathbb{P}^1 \times \{\text{pt}\}]$ and $f_2 = [\{\text{pt}\} \times \mathbb{P}^1]$.

An easy example

- Suppose that $X = \mathbb{P}^1 \times \mathbb{P}^1$.
- $\text{NE}(X)$ sits inside a two dimensional vector space. The cone is spanned by $f_1 = [\mathbb{P}^1 \times \{\text{pt}\}]$ and $f_2 = [\{\text{pt}\} \times \mathbb{P}^1]$.
- This cone has four faces. The whole cone, the zero cone and the two cones spanned by f_1 and f_2 .

An easy example

- Suppose that $X = \mathbb{P}^1 \times \mathbb{P}^1$.
- $\text{NE}(X)$ sits inside a two dimensional vector space. The cone is spanned by $f_1 = [\mathbb{P}^1 \times \{\text{pt}\}]$ and $f_2 = [\{\text{pt}\} \times \mathbb{P}^1]$.
- This cone has four faces. The whole cone, the zero cone and the two cones spanned by f_1 and f_2 .
- The corresponding morphisms are the identity, the constant map to a point, and the two projections.

An easy example

- Suppose that $X = \mathbb{P}^1 \times \mathbb{P}^1$.
- $\text{NE}(X)$ sits inside a two dimensional vector space. The cone is spanned by $f_1 = [\mathbb{P}^1 \times \{\text{pt}\}]$ and $f_2 = [\{\text{pt}\} \times \mathbb{P}^1]$.
- This cone has four faces. The whole cone, the zero cone and the two cones spanned by f_1 and f_2 .
- The corresponding morphisms are the identity, the constant map to a point, and the two projections.
- In this example, the correspondence between faces and contractions is complete and in fact every nef divisor is semiample.

A harder example

- Suppose that $X = E \times E$, where E is a general elliptic curve.

A harder example

- Suppose that $X = E \times E$, where E is a general elliptic curve.
- $\text{NE}(X)$ sits inside a three dimensional vector space. The class δ of the diagonal is independent from the classes f_1 and f_2 of the two fibres.

A harder example

- Suppose that $X = E \times E$, where E is a general elliptic curve.
- $\text{NE}(X)$ sits inside a three dimensional vector space. The class δ of the diagonal is independent from the classes f_1 and f_2 of the two fibres.
- $\text{Aut}(X)$ is large; it contains $\text{SL}(2, \mathbb{Z})$.

A harder example

- Suppose that $X = E \times E$, where E is a general elliptic curve.
- $\text{NE}(X)$ sits inside a three dimensional vector space. The class δ of the diagonal is independent from the classes f_1 and f_2 of the two fibres.
- $\text{Aut}(X)$ is large; it contains $\text{SL}(2, \mathbb{Z})$.
- There are many contractions. Start with either of the two projections and act by $\text{Aut}(X)$.

- On a surface, if $D^2 > 0$, and $D \cdot H > 0$ for some ample divisor, then D is effective by Riemann-Roch.

- On a surface, if $D^2 > 0$, and $D \cdot H > 0$ for some ample divisor, then D is effective by Riemann-Roch.
- As the action of $\text{Aut}(X)$ is transitive, there are no curves of negative self-intersection. Thus $\text{NE}(X)$ is given by $D^2 \geq 0, D \cdot H \geq 0$.

- On a surface, if $D^2 > 0$, and $D \cdot H > 0$ for some ample divisor, then D is effective by Riemann-Roch.
- As the action of $\text{Aut}(X)$ is transitive, there are no curves of negative self-intersection. Thus $\text{NE}(X)$ is given by $D^2 \geq 0, D \cdot H \geq 0$.
- $\text{NE}(X)$ is one half of the classic circular cone $x^2 + y^2 = z^2 \subset \mathbb{R}^3$. Thus many faces don't correspond to contractions.

- On a surface, if $D^2 > 0$, and $D \cdot H > 0$ for some ample divisor, then D is effective by Riemann-Roch.
- As the action of $\text{Aut}(X)$ is transitive, there are no curves of negative self-intersection. Thus $\text{NE}(X)$ is given by $D^2 \geq 0, D \cdot H \geq 0$.
- $\text{NE}(X)$ is one half of the classic circular cone $x^2 + y^2 = z^2 \subset \mathbb{R}^3$. Thus many faces don't correspond to contractions.
- Many nef divisors are not semiample. Indeed, even on an elliptic curve there are numerically trivial divisors which are not torsion.

A much harder example

- Suppose that $X = C_2, C \times C$, modulo the obvious involution, where C is a **general** curve, $g \geq 2$.

A much harder example

- Suppose that $X = C_2, C \times C$, modulo the obvious involution, where C is a **general** curve, $g \geq 2$.
- C_2 corresponds to divisors $p + q$ of degree 2.

A much harder example

- Suppose that $X = C_2, C \times C$, modulo the obvious involution, where C is a **general** curve, $g \geq 2$.
- C_2 corresponds to divisors $p + q$ of degree 2.
- $\text{NE}(X)$ sits inside a two dimensional vector space, spanned by the image δ of the class of the diagonal and the image f of the class of a fibre. In particular the cone is spanned by two rays.

A much harder example

- Suppose that $X = C_2, C \times C$, modulo the obvious involution, where C is a **general** curve, $g \geq 2$.
- C_2 corresponds to divisors $p + q$ of degree 2.
- $\text{NE}(X)$ sits inside a two dimensional vector space, spanned by the image δ of the class of the diagonal and the image f of the class of a fibre. In particular the cone is spanned by two rays.
- $\delta^2 = 1 - g < 0$; it is contracted by the Abel-Jacobi map. So δ spans a ray. But what is the other ray?

A much harder example

- Suppose that $X = C_2, C \times C$, modulo the obvious involution, where C is a **general** curve, $g \geq 2$.
- C_2 corresponds to divisors $p + q$ of degree 2.
- $\text{NE}(X)$ sits inside a two dimensional vector space, spanned by the image δ of the class of the diagonal and the image f of the class of a fibre. In particular the cone is spanned by two rays.
- $\delta^2 = 1 - g < 0$; it is contracted by the Abel-Jacobi map. So δ spans a ray. But what is the other ray?
- Conjecturally it spanned by the class of a curve of self-intersection zero. But why could there not be another curve Σ , such that $\Sigma^2 < 0$?

More Pathologies

- If $S \rightarrow C$ is the projectivisation of a stable rank two vector bundle over a curve of genus $g \geq 2$, then $\text{NE}(S)$ sits inside a two dimensional vector space.

More Pathologies

- If $S \rightarrow C$ is the projectivisation of a stable rank two vector bundle over a curve of genus $g \geq 2$, then $\text{NE}(S)$ sits inside a two dimensional vector space.
- One edge is spanned by the class f of a fibre. The other edge is corresponds to a class α of self-intersection zero.

More Pathologies

- If $S \rightarrow C$ is the projectivisation of a stable rank two vector bundle over a curve of genus $g \geq 2$, then $\text{NE}(S)$ sits inside a two dimensional vector space.
- One edge is spanned by the class f of a fibre. The other edge is corresponds to a class α of self-intersection zero.
- However there is no curve Σ such that the class of C is equal to α .

More Pathologies

- If $S \rightarrow C$ is the projectivisation of a stable rank two vector bundle over a curve of genus $g \geq 2$, then $\text{NE}(S)$ sits inside a two dimensional vector space.
- One edge is spanned by the class f of a fibre. The other edge is corresponds to a class α of self-intersection zero.
- However there is no curve Σ such that the class of C is equal to α .
- Indeed the existence of such a curve would imply that the pullback of S along $\Sigma \rightarrow C$ splits, which contradicts stability.

More Pathologies

- If $S \rightarrow C$ is the projectivisation of a stable rank two vector bundle over a curve of genus $g \geq 2$, then $\text{NE}(S)$ sits inside a two dimensional vector space.
- One edge is spanned by the class f of a fibre. The other edge is corresponds to a class α of self-intersection zero.
- However there is no curve Σ such that the class of C is equal to α .
- Indeed the existence of such a curve would imply that the pullback of S along $\Sigma \rightarrow C$ splits, which contradicts stability.
- We really need to take the closure, to define $\text{NE}(S)$.

Even more Pathologies

- Let $S \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.

Even more Pathologies

- Let $S \longrightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.
- We can perturb one point, so that the nine points are the intersection of two smooth cubics.

Even more Pathologies

- Let $S \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.
- We can perturb one point, so that the nine points are the intersection of two smooth cubics.
- In this case $S \rightarrow \mathbb{P}^1$, with elliptic fibres.

Even more Pathologies

- Let $S \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.
- We can perturb one point, so that the nine points are the intersection of two smooth cubics.
- In this case $S \rightarrow \mathbb{P}^1$, with elliptic fibres.
- The nine exceptional divisors are sections. The difference of any two is not torsion in the generic fibre. Translating by the difference generates infinitely many exceptional divisors.

Even more Pathologies

- Let $S \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.
- We can perturb one point, so that the nine points are the intersection of two smooth cubics.
- In this case $S \rightarrow \mathbb{P}^1$, with elliptic fibres.
- The nine exceptional divisors are sections. The difference of any two is not torsion in the generic fibre. Translating by the difference generates infinitely many exceptional divisors.
- Perturbing, we lose the fibration, but keep the -1 -curves.

Even more Pathologies

- Let $S \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at 9 general points.
- We can perturb one point, so that the nine points are the intersection of two smooth cubics.
- In this case $S \rightarrow \mathbb{P}^1$, with elliptic fibres.
- The nine exceptional divisors are sections. The difference of any two is not torsion in the generic fibre. Translating by the difference generates infinitely many exceptional divisors.
- Perturbing, we lose the fibration, but keep the -1 -curves.
- What went wrong?

The canonical divisor

- The answer in all cases is to consider the behaviour of the canonical divisor K_X .

The canonical divisor

- The answer in all cases is to consider the behaviour of the canonical divisor K_X .
- Recall that the canonical divisor is defined by picking a meromorphic section of $\wedge^n T_X^*$, and looking at its zeroes minus poles.

The canonical divisor

- The answer in all cases is to consider the behaviour of the canonical divisor K_X .
- Recall that the canonical divisor is defined by picking a meromorphic section of $\wedge^n T_X^*$, and looking at its zeroes minus poles.
- The basic moral is that the cone of curves is nice on the negative side, and that if we contract these curves, we get a reasonable model.

The canonical divisor

- The answer in all cases is to consider the behaviour of the canonical divisor K_X .
- Recall that the canonical divisor is defined by picking a meromorphic section of $\wedge^n T_X^*$, and looking at its zeroes minus poles.
- The basic moral is that the cone of curves is nice on the negative side, and that if we contract these curves, we get a reasonable model.
- Consider the case of curves.

Smooth projective curves

- Curves C come in three types:

Smooth projective curves

- Curves C come in three types:
- $C \simeq \mathbb{P}^1$.

Smooth projective curves

- Curves C come in three types:
- $C \simeq \mathbb{P}^1$. K_C is negative.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is negative.
 - C is elliptic, a plane cubic.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is **negative**.
 - C is elliptic, a plane cubic. K_C is **zero**.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is **negative**.
 - C is elliptic, a plane cubic. K_C is **zero**.
 - C has genus at least two.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is **negative**.
 - C is elliptic, a plane cubic. K_C is **zero**.
 - C has genus at least two. K_C is **positive**.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is **negative**.
 - C is elliptic, a plane cubic. K_C is **zero**.
 - C has genus at least two. K_C is **positive**.
- We hope (wishfully?) that the same pattern remains in higher dimensions.

Smooth projective curves

- Curves C come in three types:
 - $C \simeq \mathbb{P}^1$. K_C is **negative**.
 - C is elliptic, a plane cubic. K_C is **zero**.
 - C has genus at least two. K_C is **positive**.
- We hope (wishfully?) that the same pattern remains in higher dimensions.
- So let us now consider surfaces.

Smooth projective surfaces

- Any smooth surface S is birational to:

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 .

Smooth projective surfaces

- Any smooth surface S is birational to:
- \mathbb{P}^2 . $-K_S$ is ample, a Fano variety.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 .

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.
 - $S \rightarrow C$, where K_S is zero on the fibres.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.
 - $S \rightarrow C$, where K_S is zero on the fibres. If C is a curve, the fibres are elliptic curves.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.
 - $S \rightarrow C$, where K_S is zero on the fibres. If C is a curve, the fibres are elliptic curves.
 - K_S is ample.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.
 - $S \rightarrow C$, where K_S is zero on the fibres. If C is a curve, the fibres are elliptic curves.
 - K_S is ample. S is of general type. Note that S is forced to be singular in general.

Smooth projective surfaces

- Any smooth surface S is birational to:
 - \mathbb{P}^2 . $-K_S$ is ample, a **Fano variety**.
 - $S \rightarrow C$, $g(C) \geq 1$, where the fibres are isomorphic to \mathbb{P}^1 . $-K_S$ is relatively ample, a **Fano fibration**.
 - $S \rightarrow C$, where K_S is zero on the fibres. If C is a curve, the fibres are elliptic curves.
 - K_S is ample. S is of general type. Note that S is forced to be singular in general.
- The problem, as we have already seen, is that we can destroy this picture, simply by blowing up. It is the aim of the MMP to reverse the process of blowing up.

The cone theorem

- Let X be a smooth variety, or in general mildly singular. There are two cases:

The cone theorem

- Let X be a smooth variety, or in general mildly singular. There are two cases:
 - K_X is nef.

The cone theorem

- Let X be a smooth variety, or in general mildly singular. There are two cases:
 - K_X is nef.
 - There is a curve C such that $K_X \cdot C < 0$.

The cone theorem

- Let X be a smooth variety, or in general mildly singular. There are two cases:
 - K_X is nef.
 - There is a curve C such that $K_X \cdot C < 0$.
- In the second case there is a K_X -extremal ray R . That is to say R is extremal in the sense of convex geometry, and $K_X \cdot R < 0$.

The cone theorem

- Let X be a smooth variety, or in general mildly singular. There are two cases:
 - K_X is nef.
 - There is a curve C such that $K_X \cdot C < 0$.
- In the second case there is a K_X -extremal ray R . That is to say R is extremal in the sense of convex geometry, and $K_X \cdot R < 0$.
- Moreover, we can contract R , $\phi_R: X \longrightarrow Y$.

The case of surfaces

- Let S be a smooth surface. Suppose that K_S is not nef. Let R be an extremal ray, $\phi: S \longrightarrow Z$. There are three cases:

The case of surfaces

- Let S be a smooth surface. Suppose that K_S is not nef. Let R be an extremal ray, $\phi: S \longrightarrow Z$. There are three cases:
- Z is a point. In this case $S \simeq \mathbb{P}^2$.

The case of surfaces

- Let S be a smooth surface. Suppose that K_S is not nef. Let R be an extremal ray, $\phi: S \longrightarrow Z$. There are three cases:
 - Z is a point. In this case $S \simeq \mathbb{P}^2$.
 - Z is a curve. The fibres are copies of \mathbb{P}^1 .

The case of surfaces

- Let S be a smooth surface. Suppose that K_S is not nef. Let R be an extremal ray, $\phi: S \longrightarrow Z$. There are three cases:
 - Z is a point. In this case $S \simeq \mathbb{P}^2$.
 - Z is a curve. The fibres are copies of \mathbb{P}^1 .
 - Z is a surface. ϕ blows down a -1 -curve.

The MMP for surfaces

- Start with a smooth surface S .

The MMP for surfaces

- Start with a smooth surface S .
- If K_S is nef, then **STOP**.

The MMP for surfaces

- Start with a smooth surface S .
- If K_S is nef, then **STOP**.
- Otherwise there is a K_S -extremal ray R , with associated contraction $\phi: S \longrightarrow Z$.

The MMP for surfaces

- Start with a smooth surface S .
- If K_S is nef, then **STOP**.
- Otherwise there is a K_S -extremal ray R , with associated contraction $\phi: S \longrightarrow Z$.
- If $\dim Z < 2$, then **STOP**.

The MMP for surfaces

- Start with a smooth surface S .
- If K_S is nef, then **STOP**.
- Otherwise there is a K_S -extremal ray R , with associated contraction $\phi: S \rightarrow Z$.
- If $\dim Z < 2$, then **STOP**.
- If $\dim Z = 2$ then replace S with Z , and continue.

The general algorithm

- Start with any birational model X .

The general algorithm

- Start with any birational model X .
- Desingularise X .

The general algorithm

- Start with any birational model X .
- Desingularise X .
- If K_X is nef, then **STOP**.

The general algorithm

- Start with any birational model X .
- Desingularise X .
- If K_X is nef, then **STOP**.
- Otherwise there is a curve C , such that $K_X \cdot C < 0$.
Our aim is to remove this curve or reduce the question to a lower dimensional one.

The general algorithm

- Start with any birational model X .
- Desingularise X .
- If K_X is nef, then **STOP**.
- Otherwise there is a curve C , such that $K_X \cdot C < 0$.
Our aim is to remove this curve or reduce the question to a lower dimensional one.
- By the Cone Theorem, there is an extremal contraction, $\pi: X \longrightarrow Y$, of relative Picard number one such that for a curve C' , $\pi(C')$ is a point iff C' is homologous to C .

Analyzing π

- If the fibres of π have dimension at least one, then we have a Mori fibre space, that is $-K_X$ is π -ample, π has connected fibres and relative Picard number one. We have reduced the question to a lower dimensional one: **STOP**.

Analyzing π


- If the fibres of π have dimension at least one, then we have a Mori fibre space, that is $-K_X$ is π -ample, π has connected fibres and relative Picard number one. We have reduced the question to a lower dimensional one: **STOP**.
- If π is birational and the locus contracted by π is a divisor, then even though Y might be singular, it will at least be \mathbb{Q} -factorial (for every Weil divisor D , some multiple is Cartier).
Replace X by Y and keep going.

π is birational

- If the locus contracted by π is not a divisor, that is, π is small, then Y is not \mathbb{Q} -factorial.

π is birational

- If the locus contracted by π is not a divisor, that is, π is small, then Y is not \mathbb{Q} -factorial.
- Instead of contracting C , we try to replace X by another birational model X^+ , $X \dashrightarrow X^+$, such that $\pi^+: X^+ \rightarrow Y$ is K_{X^+} -ample.



Flips

- This operation is called a **flip**.

Flips

- This operation is called a **flip**.
- Even supposing we can perform a flip, how do we know that this process terminates?

Flips

- This operation is called a **flip**.
- Even supposing we can perform a flip, how do we know that this process terminates?
- It is clear that we cannot keep contracting divisors, but why could there not be an infinite sequence of flips?

Adjunction and Vanishing, I

- In higher dimensional geometry, there are two basic results, adjunction and vanishing.

Adjunction and Vanishing, I

- In higher dimensional geometry, there are two basic results, adjunction and vanishing.
- (Adjunction) In its simplest form it states that given a variety smooth X and a divisor S , the restriction of $K_X + S$ to S is equal to K_S .

Adjunction and Vanishing, I

- In higher dimensional geometry, there are two basic results, adjunction and vanishing.
- (Adjunction) In its simplest form it states that given a variety smooth X and a divisor S , the restriction of $K_X + S$ to S is equal to K_S .
- (Vanishing) The simplest form is Kodaira vanishing which states that if X is smooth and L is an ample line bundle, then $H^i(K_X + L) = 0$, for $i > 0$.

Adjunction and Vanishing, I

- In higher dimensional geometry, there are two basic results, adjunction and vanishing.
- (Adjunction) In its simplest form it states that given a variety smooth X and a divisor S , the restriction of $K_X + S$ to S is equal to K_S .
- (Vanishing) The simplest form is Kodaira vanishing which states that if X is smooth and L is an ample line bundle, then $H^i(K_X + L) = 0$, for $i > 0$.
- Both of these results have far reaching generalisations, whose form dictates the main definitions of the subject.

An illustrative example

- Let S be a smooth projective surface and let $E \subset S$ be a -1 -curve, that is $K_S \cdot E = -1$ and $E^2 = -1$. We want to contract E .

An illustrative example

- Let S be a smooth projective surface and let $E \subset S$ be a -1 -curve, that is $K_S \cdot E = -1$ and $E^2 = -1$. We want to contract E .
- By adjunction, K_E has degree -2 , so that $E \simeq \mathbb{P}^1$. Pick up an ample divisor H and consider $D = K_S + G + E = K_S + aH + bE$.

An illustrative example

- Let S be a smooth projective surface and let $E \subset S$ be a -1 -curve, that is $K_S \cdot E = -1$ and $E^2 = -1$. We want to contract E .
- By adjunction, K_E has degree -2 , so that $E \simeq \mathbb{P}^1$. Pick up an ample divisor H and consider $D = K_S + G + E = K_S + aH + bE$.
- Pick $a > 0$ so that $K_S + aH$ is ample.

An illustrative example

- Let S be a smooth projective surface and let $E \subset S$ be a -1 -curve, that is $K_S \cdot E = -1$ and $E^2 = -1$. We want to contract E .
- By adjunction, K_E has degree -2 , so that $E \simeq \mathbb{P}^1$. Pick up an ample divisor H and consider $D = K_S + G + E = K_S + aH + bE$.
- Pick $a > 0$ so that $K_S + aH$ is ample.
- Then pick b so that $(K_S + aH + bE) \cdot E = 0$. Note that $b > 0$ (in fact typically b is very large).

An illustrative example

- Let S be a smooth projective surface and let $E \subset S$ be a -1 -curve, that is $K_S \cdot E = -1$ and $E^2 = -1$. We want to contract E .
- By adjunction, K_E has degree -2 , so that $E \simeq \mathbb{P}^1$. Pick up an ample divisor H and consider $D = K_S + G + E = K_S + aH + bE$.
- Pick $a > 0$ so that $K_S + aH$ is ample.
- Then pick b so that $(K_S + aH + bE) \cdot E = 0$. Note that $b > 0$ (in fact typically b is very large).
- Now we consider the rational map given by $|mD|$, for $m \gg 0$ and sufficiently divisible.

Basepoint Freeness

- Clearly the base locus of $|mD|$ is contained in E .

Basepoint Freeness

- Clearly the base locus of $|mD|$ is contained in E .
- So consider the restriction exact sequence

$$0 \longrightarrow \mathcal{O}_S(mD - E) \longrightarrow \mathcal{O}_S(mD) \longrightarrow \mathcal{O}_E(mD) \longrightarrow 0.$$

Basepoint Freeness

- Clearly the base locus of $|mD|$ is contained in E .
- So consider the restriction exact sequence

$$0 \longrightarrow \mathcal{O}_S(mD - E) \longrightarrow \mathcal{O}_S(mD) \longrightarrow \mathcal{O}_E(mD) \longrightarrow 0.$$

- Now

$$mD - E = K_S + G + (m - 1)D,$$

and $G + (m - 1)D$ is ample.

Basepoint Freeness

- Clearly the base locus of $|mD|$ is contained in E .
- So consider the restriction exact sequence

$$0 \longrightarrow \mathcal{O}_S(mD-E) \longrightarrow \mathcal{O}_S(mD) \longrightarrow \mathcal{O}_E(mD) \longrightarrow 0.$$

- Now

$$mD - E = K_S + G + (m-1)D,$$

and $G + (m-1)D$ is ample.

- So by Kawamata-Viehweg Vanishing

$$H^1(S, \mathcal{O}_S(mD-E)) = H^1(S, \mathcal{O}_S(K_S+G+(m-1)D)) = 0.$$

Castelnuovo's Criteria

- By assumption $\mathcal{O}_E(mD)$ is the trivial line bundle.
But this is a cheat.

Castelnuovo's Criteria

- By assumption $\mathcal{O}_E(mD)$ is the trivial line bundle.
But this is a cheat.
- In fact by adjunction

$$(K_S + G + E)|_E = K_E + B,$$

where $B = G|_E$.

Castelnuovo's Criteria

- By assumption $\mathcal{O}_E(mD)$ is the trivial line bundle.
But this is a cheat.
- In fact by adjunction

$$(K_S + G + E)|_E = K_E + B,$$

where $B = G|_E$.

- B is ample, so we have the start of an induction.

Castelnuovo's Criteria

- By assumption $\mathcal{O}_E(mD)$ is the trivial line bundle.
But this is a cheat.
- In fact by adjunction

$$(K_S + G + E)|_E = K_E + B,$$

where $B = G|_E$.

- B is ample, so we have the start of an induction.
- By vanishing, the map

$$H^0(S, \mathcal{O}_S(mD)) \longrightarrow H^0(E, \mathcal{O}_E(mD))$$

is surjective. Thus $|mD|$ is base point free and the resulting map $S \longrightarrow T$ contracts E .

The General Case

- We want to try to do the same thing, but in higher dimension. Unfortunately the locus E we want to contract need not be a divisor.

The General Case

- We want to try to do the same thing, but in higher dimension. Unfortunately the locus E we want to contract need not be a divisor.
- Observe that if we set $G' = \pi_* G$, then G' has high multiplicity along p , the image of E (that is b is large).

The General Case

- We want to try to do the same thing, but in higher dimension. Unfortunately the locus E we want to contract need not be a divisor.
- Observe that if we set $G' = \pi_* G$, then G' has high multiplicity along p , the image of E (that is b is large).
- In general, we manufacture a divisor E by picking a point $x \in X$ and then pick H with high multiplicity at x .

The General Case

- We want to try to do the same thing, but in higher dimension. Unfortunately the locus E we want to contract need not be a divisor.
- Observe that if we set $G' = \pi_* G$, then G' has high multiplicity along p , the image of E (that is b is large).
- In general, we manufacture a divisor E by picking a point $x \in X$ and then pick H with high multiplicity at x .
- Next resolve singularities $\tilde{X} \rightarrow X$ and restrict to an exceptional divisor E , whose centre has high multiplicity w.r.t H (strictly speaking a log canonical centre of $K_X + H$).

Singularities in the MMP

- Let X be a normal variety. We say that a divisor $\Delta = \sum_i a_i \Delta_i$ is a **boundary**, if $0 \leq a_i \leq 1$.

Singularities in the MMP

- Let X be a normal variety. We say that a divisor $\Delta = \sum_i a_i \Delta_i$ is a **boundary**, if $0 \leq a_i \leq 1$.
- Let $\pi: Y \longrightarrow X$ be birational map. Suppose that $K_X + \Delta$ is \mathbb{Q} -Cartier. Then we may write

$$K_Y + \Gamma = \pi^*(K_X + \Delta).$$

Singularities in the MMP

- Let X be a normal variety. We say that a divisor $\Delta = \sum_i a_i \Delta_i$ is a **boundary**, if $0 \leq a_i \leq 1$.
- Let $\pi: Y \longrightarrow X$ be birational map. Suppose that $K_X + \Delta$ is \mathbb{Q} -Cartier. Then we may write

$$K_Y + \Gamma = \pi^*(K_X + \Delta).$$

- We say that the pair (X, Δ) is **klt** if the coefficients of Γ are always less than one.

Adjunction II

- To apply adjunction we need a component S of coefficient one.

Adjunction II

- To apply adjunction we need a component S of coefficient one.
- So suppose we can write $\Delta = S + B$, where S has coefficient one. Then

$$(K_X + S + B)|_S = K_S + D.$$

Adjunction II

- To apply adjunction we need a component S of coefficient one.
- So suppose we can write $\Delta = S + B$, where S has coefficient one. Then

$$(K_X + S + B)|_S = K_S + D.$$

- Moreover if $K_X + S + B$ is plt then $K_S + D$ is klt.

Vanishing II

- We want a form of vanishing which involves boundaries.

Vanishing II

- We want a form of vanishing which involves boundaries.
- If we take a cover with appropriate ramification, then we can eliminate any component with coefficient less than one.

Vanishing II

- We want a form of vanishing which involves boundaries.
- If we take a cover with appropriate ramification, then we can eliminate any component with coefficient less than one.
- (Kawamata-Viehweg vanishing) Suppose that $K_X + \Delta$ is **klt** and L is a line bundle such that $L - (K_X + \Delta)$ is big and nef. Then, for $i > 0$,

$$H^i(X, L) = 0.$$

Summary

- We hope that varieties X belong to two types:

Summary

- We hope that varieties X belong to two types:
- X is a **minimal model**: K_X is **nef**. That is $K_X \cdot C \geq 0$, for every curve C in X .

Summary

- We hope that varieties X belong to two types:
- X is a **minimal model**: K_X is **nef**. That is $K_X \cdot C \geq 0$, for every curve C in X .
- X is a **Mori fibre space**, $\pi: X \rightarrow Y$. That is π is **extremal** ($-K_X$ is relatively ample and π has relative Picard one) and π is a **contraction** (the fibres of π are connected) of dimension at least one.

Summary

- We hope that varieties X belong to two types:
 - X is a **minimal model**: K_X is **nef**. That is $K_X \cdot C \geq 0$, for every curve C in X .
 - X is a **Mori fibre space**, $\pi: X \rightarrow Y$. That is π is **extremal** ($-K_X$ is relatively ample and π has relative Picard one) and π is a **contraction** (the fibres of π are connected) of dimension at least one.
- To achieve this birational classification, we propose to use the **MMP**.

Two main Conjectures

To finish the proof of the existence of the MMP, we need to prove the following two conjectures:

Two main Conjectures

To finish the proof of the existence of the MMP, we need to prove the following two conjectures:

Conjecture. (*Existence*) Suppose that $K_X + \Delta$ is kawamata log terminal. Let $\pi: X \rightarrow Y$ be a small extremal contraction.

Then the flip of π exists.

Two main Conjectures

To finish the proof of the existence of the MMP, we need to prove the following two conjectures:

Conjecture. (Existence) Suppose that $K_X + \Delta$ is kawamata log terminal. Let $\pi: X \rightarrow Y$ be a small extremal contraction.

Then the flip of π exists.

Conjecture. (Termination) There is no infinite sequence of kawamata log terminal flips.

Abundance

Now suppose that X is a minimal model, so that K_X is nef.

Abundance

Now suppose that X is a minimal model, so that K_X is nef.

Conjecture. (*Abundance*) *Suppose that $K_X + \Delta$ is kawamata log terminal and nef.
Then $K_X + \Delta$ is semiample.*

Abundance

Now suppose that X is a minimal model, so that K_X is nef.

Conjecture. (*Abundance*) Suppose that $K_X + \Delta$ is kawamata log terminal and nef.
Then $K_X + \Delta$ is semiample.

Considering the resulting morphism $\phi: X \longrightarrow Y$, we recover the Kodaira-Enriques classification of surfaces.