

Is the subdominant part of the energy spectrum due to downscale energy cascade hidden in quasi-geostrophic turbulence?

Eleftherios Gkioulekas* and Ka Kit Tung†

Department of Applied Mathematics, University of Washington, Seattle, WA, United States

In systems governing two-dimensional turbulence, surface quasi-geostrophic turbulence, (more generally α -turbulence), two-layer quasi-geostrophic turbulence, etc., there often exist two conservative quadratic quantities, one “energy”-like and one “enstrophy”-like. In a finite inertial range there are in general two spectral fluxes, one associated with each conserved quantity. We derive here an inequality comparing the relative magnitudes of the “energy” and “enstrophy” fluxes for finite or infinitesimal dissipations, and for hyper or hypo viscosities. When this inequality is satisfied, as is the case of 2D turbulence, where the energy flux contribution to the energy spectrum is small, the subdominant part will be effectively hidden. In SQG turbulence, it is shown that the opposite is true: the downscale energy flux becomes the dominant contribution to the energy spectrum. A combination of these two behaviors appears to be the case in 2-layer QG turbulence, depending on the baroclinicity of the system.

1. INTRODUCTION

The characteristic feature of two-dimensional turbulence is that there are two conserved quantities, kinetic energy and enstrophy. This led Kraichnan (1967), Leith (1968), and Batchelor (1969) to conjecture that there will exist two inertial ranges, one located upscale of the spectral region of injection and another on the downscale side of injection. In the upscale side, it is assumed that there is only an upscale flux of energy, and no flux of enstrophy. On the downscale side, likewise, there is only a downscale flux of enstrophy, and no flux of energy. One then uses a dimensional analysis argument to calculate the energy spectrum $E(k)$ where it is assumed that in each inertial range it depends only on the corresponding single flux and the wavenumber k . The same type of argument was used in the energy cascade of three-dimensional turbulence (Batchelor, 1947; Kolmogorov, 1941a,b). Although three-dimensional turbulence also has two conserved quantities, energy and helicity, one has the option to inject energy without injecting helicity. In two-dimensional turbulence it is not possible to inject energy without injecting enstrophy and vice versa, because the two quantities are related

Initial efforts to simulate the enstrophy cascade yielded confusing reports of various numerical slopes. Consequently, alternative theories have been proposed over the past 30 years to explain them (Moffatt, 1986; Polyakov, 1993; Saffman, 1971). Recently, in carefully set up experiments, it was shown that it is possible to obtain the enstrophy cascade in agreement with the KLB theory (Ishihara and Kaneda, 2001; Lindborg and Alvelius, 2000; Pasquero and Falkovich, 2002). A numerical simulation with very good diagnostics has shown that the inverse energy cascade can be obtained accordingly (Boffetta et al., 2000). There are also however many papers that question the universality of these results (Danilov, 2005; Danilov and Gurarie, 2001a,b; Tran and Bowman, 2003, 2004). A review can be found in

Tableting (2002) and Gkioulekas and Tung (2005c).

Further confusion has resulted from efforts to explain the observed energy spectrum of the atmosphere with the KLB theory. Observations show that there is a robust energy spectrum with slope -3 which transitions at large wavenumbers into slope $-5/3$ (Gage, 1979; Gage and Nastrom, 1986; Nastrom and Gage, 1984; Nastrom et al., 1984). In the KLB theory, on the other hand, one expects that at small wavenumbers the energy spectrum will have slope $-5/3$ from the inverse energy cascade, which will then transition at the forcing wavenumber, into a -3 slope from the direct enstrophy cascade. The apparent contradiction between these two predictions has led to various explanations and debate (Dewan, 1979; Lilly, 1989; Lindborg, 1999; VanZadt, 1982).

It was conjectured by Tung and Orlando (2003a) that the observed atmospheric energy spectrum results from the downscale cascade of enstrophy and energy injected at the large scales by baroclinic instability and dissipated at the smallest length scales. If η_{uv} is the downscale enstrophy flux and ε_{uv} is the downscale energy flux, it was suggested that they would coexist on the downscale side of injection and that their separate contributions to the energy spectrum would give the latter a compound spectral shape, with a -3 slope transitioning to a shallower $-5/3$ slope as the wavenumber increases. The transition from -3 slope to $-5/3$ slope occurs at the transition wavenumber k_t with order of magnitude estimated by $k_t \approx \sqrt{\eta_{uv}/\varepsilon_{uv}}$.

General Circulation models have been shown to be capable of reproducing the Nastrom-Gage spectrum in agreement with observations (Koshyk and Hamilton, 2001; Koshyk et al., 1999; Skamarock, 2004). Although the nature of the nonlinear interactions which give rise to the downscale energy flux changes from quasi-geostrophic to stratified three-dimensional in the mesoscales, as far as the energy spectrum $E(k)$ is concerned it is the existence of a downscale energy flux which gives rise to the $k^{-5/3}$ slope, regardless of the character of the motion. The recent interest, typified by Lindborg (2005), in understanding the $k^{-5/3}$ slope in terms of three-dimensional stratified turbulence is justified, since it is necessary to account for length scales less than 100km in

*Electronic address: lf@amath.washington.edu

†Electronic address: tung@amath.washington.edu

wavelength where the quasi-geostrophic assumption fails. It is the view of the authors that it is equally important to understand why the quasi-geostrophic model is capable of supporting a downscale energy cascade with $k^{-5/3}$ scaling, because one also has to account for the existence of $k^{-5/3}$ scaling at length scales ranging from 700km to 100km where the quasi-geostrophic assumption is presumably valid.

Tung and Orlando (2003a) have demonstrated numerically that a two-layer quasi-geostrophic channel model with thermal forcing, Ekman damping, and hyperdiffusion can also reproduce this compound spectrum. The resolution of these simulations goes down to 100km in wavelength. The diagnostic shown in figure 7 of (Tung and Orlando, 2003a), shows both the constant downscale energy and enstrophy fluxes co-existing in the same inertial range. Recent measurements and data analysis by Cho and Lindborg (2001) have confirmed the existence of a downscale energy flux and estimate $\eta_{uv} \approx 2 \times 10^{-15} \text{s}^{-3}$ and $\varepsilon_{uv} \approx 6 \times 10^{-11} \text{km}^2 \text{s}^{-3}$. From these estimates we find the mean value of the transition scale $k_t = \sqrt{\eta_{uv}/\varepsilon_{uv}} \approx 0.57 \times 10^{-2} \text{km}^{-1}$ and $\lambda_t = 2\pi/k_t \approx 1 \times 10^3 \text{km}$ which has the correct order of magnitude.

This theory of a combined downscale enstrophy cascade and downscale energy cascade is contrary to the widely accepted misconception that the argument by Fjørtoft (1953) forbids a downscale energy flux in two-dimensional turbulence altogether, and through the isomorphism theorem of Charney (1971) also in quasi-geostrophic turbulence. Various aspects of this misconception have been clarified by Merilees and Warn (1975), Tung and Welch (2001) and Gkioulekas and Tung (2005a).

As has been pointed out by previous authors (Borue, 1994; Eyink, 1996), as long as the dissipation terms at large-scale and small scales have finite viscosity coefficients and the inertial ranges exist, the downscale enstrophy flux will be accompanied by a small downscale energy flux, and the upscale energy flux will be accompanied by a small upscale enstrophy flux. Dimensional analysis arguments are premised on the assumption that these additional fluxes can be ignored, consequently the energy spectrum predictions obtained by such arguments are valid only to leading order. While ignoring sub-leading effects can be justified for strictly two-dimensional turbulence, we will argue in this paper that for models of quasi-geostrophic turbulence, such as the two-layer model, the sub-leading contributions can be important in the inertial range and cannot be safely ignored. Predicting the form of these sub-leading corrections requires a subtle mathematical argument, given by Gkioulekas and Tung (2005b), that goes beyond dimensional analysis.

In particular, Gkioulekas and Tung (2005b) have argued that the sub-leading fluxes are associated with a sub-leading downscale energy cascade and a sub-leading inverse enstrophy cascade that contribute *linearly* to the total energy spectrum in addition to the dominant contributions. The two contributions are homogeneous solutions of the underlying statistical theory, which is in fact *linear*. Furthermore, the two homogeneous solutions are independent of each other, so the downscale energy cascade is independent of the downscale enstrophy flux η_{uv} and the downscale enstrophy cascade is inde-

pendent of the downscale energy flux ε_{uv} . As a result, in the downscale inertial range, the total energy spectrum $E(k)$ has the following three contributions:

$$E(k) = E_{uv}^{(\varepsilon)}(k) + E_{uv}^{(\eta)}(k) + E_{uv}^{(p)}(k), \quad \forall k \ell_0 \gg 1, \quad (1)$$

where $E_{uv}^{(\varepsilon)}(k)$, $E_{uv}^{(\eta)}(k)$ are the contributions of the downscale energy and enstrophy cascade, given by

$$\begin{aligned} E_{uv}^{(\varepsilon)}(k) &= a_{uv} \varepsilon_{uv}^{2/3} k^{-5/3} \mathcal{D}_{uv}^{(\varepsilon)}(k \ell_{uv}^{(\varepsilon)}) \\ E_{uv}^{(\eta)}(k) &= b_{uv} \eta_{uv}^{2/3} k^{-3} [\chi + \ln(k \ell_0)]^{-1/3} \mathcal{D}_{uv}^{(\eta)}(k \ell_{uv}^{(\eta)}), \end{aligned} \quad (2)$$

with $\mathcal{D}_{uv}^{(\varepsilon)}$ and $\mathcal{D}_{uv}^{(\eta)}$ describing the dissipative corrections. Here we use the logarithmic correction of Kraichnan (1971), adjusted by the constant χ of Bowman (1996) for the contribution of the enstrophy cascade. We have also assumed without explicit justification that we may ignore the possibility of intermittency corrections to the sub-leading downscale energy cascade. For the downscale enstrophy cascade intermittency corrections have been ruled out by Eyink (2001). For the downscale energy cascade we conjecture that intermittency corrections are small for the same reasons as in three dimensional turbulence. The scales $\ell_{uv}^{(\varepsilon)}$, $\ell_{uv}^{(\eta)}$ are the dissipation length scales for the downscale energy and enstrophy cascade. Finally, $E_{uv}^{(p)}(k)$ is the contribution from the effect of forcing and the sweeping interactions. The latter can become significant via the violation of statistical homogeneity caused by the boundary conditions (see Gkioulekas (2005) for details). Thus, in the inertial range where the effect of forcing and dissipation can be ignored, the energy spectrum will take the simple form in the downscale range:

$$E(k) \approx a_{uv} \varepsilon_{uv}^{2/3} k^{-5/3} + b_{uv} \eta_{uv}^{2/3} k^{-3} [\chi + \ln(k \ell_0)]^{-1/3}. \quad (3)$$

We see that the energy spectrum will take the slope of -3 for small k , and $-5/3$ for large k . The transition from one slope to the other occurs at k_t , given by $\varepsilon_{uv} k_t^2 \sim \eta_{uv}$.

It should be emphasized that the formation of cascades observable in the energy spectrum is by no means guaranteed. There are two prerequisites that need to be satisfied: first, the contribution of the particular solution $E_{uv}^{(p)}(k)$ has to be negligible both downscale and upscale of the injection scale, i.e.

$$\begin{aligned} E_{uv}^{(p)}(k) &\ll E_{uv}^{(\varepsilon)}(k) + E_{uv}^{(\eta)}(k), \quad \forall k \ell_0 \gg 1 \\ E_{ir}^{(p)}(k) &\ll E_{ir}^{(\varepsilon)}(k) + E_{ir}^{(\eta)}(k), \quad \forall k \ell_0 \ll 1. \end{aligned} \quad (4)$$

Second, the dissipative adjustment $\mathcal{D}_{uv}^{(\eta)}(k \ell_{uv}^{(\eta)})$ and $\mathcal{D}_{uv}^{(\varepsilon)}(k \ell_{uv}^{(\varepsilon)})$ of the homogeneous solution has to be such that it does not destroy the power law scaling in the inertial range. Furthermore, the dissipation scales $\ell_{uv}^{(\eta)}$ and $\ell_{uv}^{(\varepsilon)}$ have to be positioned so that the incoming energy and enstrophy can be dissipated.

The idea of two cascades in the same wavenumber region has an interesting precedent in the case of three-dimensional turbulence, where there is interest in understanding the double cascade of helicity and energy (Arad et al., 1999; Chen et al.,

2003; Moiseev and Chkhetiani, 1996). There, the situation is more straightforward because the helicity cascade and the energy cascade reside in separate isotropic sectors of the $SO(3)$ group (Arad et al., 1999; Biferale and Procaccia, 2005). This makes it easier to argue in support of a superposition principle.

In two-dimensional turbulence the situation is more interesting because both cascades reside in the same isotropic sectors. The main argument in support of our conjecture was given in section 2 of Gkioulekas and Tung (2005b). Additional evidence is given in section 3 of the same paper. It should be noted that our main argument exploits the linearity of the exact statistical theory of two-dimensional turbulence (i.e. the complete infinite system of equations governing the relevant fully-unfused structure functions). Nonlinear results, such as the one that was proposed by Lilly (1989), follow from closure models instead of the exact theory. Likewise, phenomenological arguments with the eddy-turnover rate, such as the one in Kraichnan (1971), are essentially coming out of a 1-loop nonlinear closure theory, and would also lead to nonlinear expressions for the energy spectrum. Most closure arguments miss the point that the *exact* statistical theory is in fact *linear*.

It should be noted that the main results of the present paper rely on the statement that the transition wavenumber k_t from -3 scaling to $-5/3$ scaling is given by $k_t \approx \sqrt{\eta_{uv}/\varepsilon_{uv}}$. This statement follows from the superposition principle, but it doesn't require it and can probably also be established with different arguments. For the case of the double cascade of helicity and energy in three-dimensional turbulence, such a different argument was given by Moiseev and Chkhetiani (1996). Furthermore, the dimensional estimate $k_t \approx \sqrt{\eta_{uv}/\varepsilon_{uv}}$ for the transition wavenumber k_t has been confirmed by Tung and Orlando (2003a). Finally, extending the superposition principle to the two-layer QG model (the main focus of this paper), which has three conservation laws instead of just two, remains an open theoretical question.

In two-dimensional turbulence, the fluxes ε_{uv} and η_{uv} are constrained by an inequality that was communicated to us by Sergey Danilov (Gkioulekas and Tung, 2005c). This constraint implies that the contribution of the downscale energy cascade to the energy spectrum is overwhelmed by the contribution of the downscale enstrophy cascade and cannot be seen visually on a plot. This result was conjectured earlier by Smith (2004) who claimed that the downscale energy cascade can never have enough flux to move the transition wavenumber k_t into the inertial range. The two-layer model is a different dynamical system than the two-dimensional Navier-Stokes equations, and the validity of the Danilov inequality in the two-layer model is not obvious (Gkioulekas and Tung, 2005c; Tung, 2004).

In the present paper we will show that in the two-layer model when the Ekman dissipation coefficient ν_E is below a critical value, then the Danilov inequality will be satisfied. We will also argue that the asymmetric presence of Ekman damping on the bottom layer but not the top layer may cause the violation of the Danilov inequality for larger values of the

Ekman dissipation coefficient. In this case, the top layer has more enstrophy than the bottom layer, as is realistic in the atmosphere, and provided that the difference in enstrophy between the two layers is large enough, the downscale energy cascade will be made observable in the energy spectrum. We derive specific conditions on how large this difference needs to be in order for the Danilov inequality to be violated for some wavenumbers k . The simulation of Tung and Orlando (2003a) has shown that it is possible to have an observable downscale energy cascade, which implies a violation of the Danilov inequality. The role of the argument in this paper is to explain how and why this can happen, given that it is a surprising and very unexpected result. We also hope that our paper will raise renewed interest in understanding the phenomenology of the two-layer quasi-geostrophic model.

We will also show that in the surface quasi-geostrophic model, which represents the most extreme case of baroclinicity, the downscale energy cascade becomes completely dominant. An immediate implication of our argument is that the existence of an extensive observable $k^{-5/3}$ in the energy spectrum of the atmosphere has the physical interpretation that the atmosphere is very baroclinic.

The paper is organized as follows. The Danilov inequality is reviewed in section 2 where we make some simple generalizations. Its implications for two-dimensional turbulence are discussed in section 3. The surface quasi-geostrophic model is discussed in section 4 and that the two-layer model in section 5. Conclusions and some further remarks are given in section 6.

2. THE DANILOV INEQUALITY IN ONE-LAYER MODELS

The governing equation for a wide range of one-layer hydrodynamic models takes the form:

$$\frac{\partial \zeta}{\partial t} + J(\psi, \zeta) = \mathcal{D} + \mathcal{F}, \quad (5)$$

where \mathcal{F} is the forcing and \mathcal{D} is the dissipation and $\zeta = -\mathcal{L}\psi$. Here, \mathcal{L} is a linear isotropic operator involving the derivatives with respect to the horizontal coordinates. For a general combination of hyper- and hypo-diffusion:

$$\mathcal{D} = -\nu_0(-\Delta)^p \zeta - \nu_1(-\Delta)^{-h} \zeta, \quad (6)$$

with p, h , positive integers. $p = 1, h = 0$ yields the combination of molecular viscosity and Ekman damping.

For 2D turbulence, \mathcal{L} is given by $\mathcal{L} = -\Delta$, where Δ is the Laplacian operator and the streamfunction ψ is related to the 2D nondivergent velocity as

$$(u, v) = \left(-\frac{\partial \psi}{\partial y}, \frac{\partial \psi}{\partial x} \right). \quad (7)$$

For barotropic QG turbulence, also known as Charney-Hasegawa-Nima (CHM) turbulence (Charney, 1948; Hasegawa et al., 1979; Hasegawa and Mima, 1978), \mathcal{L} is given instead by $\mathcal{L} = -\Delta + \lambda^2$, where λ^2 is a given positive constant. Another interesting family of one-layer models are

the α -turbulence models where $\mathcal{L} = \Lambda^\alpha$ with $\Lambda \equiv (-\Delta)^{1/2}$. The case $\alpha = 1$ corresponds to surface quasi-geostrophic turbulence (SQG) which is an extreme baroclinic model, and not a barotropic model like 2D turbulence or CHM turbulence.

2.1. Conservation laws

Let $\langle\langle f \rangle\rangle$ be the spatial and ensemble average of $f(x, y)$ defined as

$$\langle\langle f \rangle\rangle \equiv \iint \langle f(x, y) \rangle dx dy. \quad (8)$$

There are two inviscid quadratic invariants for (5), which are:

$$A = (1/2)\langle\langle -\psi\zeta \rangle\rangle, \quad (9)$$

$$B = (1/2)\langle\langle \zeta^2 \rangle\rangle. \quad (10)$$

Note that B is conserved for all linear operators \mathcal{L} , whereas the conservation law of A requires that \mathcal{L} be self-adjoint, i.e.

$$\langle\langle f(\mathcal{L}g) \rangle\rangle = \langle\langle (\mathcal{L}f)g \rangle\rangle. \quad (11)$$

For example in 2D turbulence it is well-known that,

$$E \equiv (1/2)\langle\langle (u^2 + v^2) \rangle\rangle = (1/2)\langle\langle |\nabla\psi|^2 \rangle\rangle \quad (12)$$

$$= (1/2)\langle\langle -\psi\zeta \rangle\rangle \equiv A, \quad (13)$$

is the kinetic energy of the 2D fluid, and

$$G \equiv (1/2)\langle\langle \zeta^2 \rangle\rangle \equiv B, \quad (14)$$

is the enstrophy. In section 4 we see that A and B have a different physical interpretation in SQG turbulence.

The spectra $A(k)$ and $B(k)$ of the conserved quadratics A and B are defined as

$$A(k) = \frac{1}{2} \frac{d}{dk} \langle\langle -\psi^{<k} \zeta^{<k} \rangle\rangle, \quad (15)$$

$$B(k) = \frac{1}{2} \frac{d}{dk} \langle\langle (\zeta^{<k})^2 \rangle\rangle, \quad (16)$$

with $\psi^{<k}$ and $\zeta^{<k}$ the streamfunction and vorticity fields with all the Fourier wavenumbers greater than k in magnitude filtered out. The relationship, $\zeta = -\mathcal{L}\psi$, translates into the spectral relationships in the Fourier space

$$\hat{\zeta}(\mathbf{k}) = L(|\mathbf{k}|)\hat{\psi}(\mathbf{k}), \quad B(k) = L(k)A(k), \quad (17)$$

where

$$\hat{\psi}(\mathbf{k}) = \iint \psi(\mathbf{x}) e^{-i\mathbf{k}\cdot\mathbf{x}} d\mathbf{x}. \quad (18)$$

We will assume that $L(k) > 0$, so that both $A(k)$ and $B(k)$ are positive. Furthermore, we will assume that $L(k)$ is a monotonically increasing function of k . In 2D turbulence, $L(k) = k^2$; in CHM turbulence, $L(k) = k^2 + \lambda^2$; in α -turbulence, $L(k) = k^\alpha$, and in SQG, $L(k) = k$.

Furthermore, $A(k)$ and $B(k)$ satisfy the following spectral equations:

$$\frac{\partial A(k)}{\partial t} + \frac{\partial \Pi_A(k)}{\partial k} = -D_A(k) + F_A(k) \quad (19)$$

$$\frac{\partial B(k)}{\partial t} + \frac{\partial \Pi_B(k)}{\partial k} = -D_B(k) + F_B(k). \quad (20)$$

Here, $D_A(k)$ and $D_B(k)$ are the spectral dissipation rate of $A(k)$ and $B(k)$, respectively, with

$$D_B(k) = L(k)D_A(k), \quad (21)$$

$$D_A(k) = [\nu_0 k^{2p} + \nu_1 k^{-2h}]A(k) > 0, \quad (22)$$

for a combination of hyper- and hypo-viscosities. Furthermore, $F_A(k)$ and $F_B(k)$ are the spectra of forcing also related by $F_B(k) = L(k)F_A(k)$, and, $\Pi_A(k)$ and $\Pi_B(k)$ are the spectral fluxes of A and B . The Leith (1968) constraint on the fluxes generalizes to

$$\frac{\partial \Pi_B(k)}{\partial k} = L(k) \frac{\partial \Pi_A(k)}{\partial k}. \quad (23)$$

The conservation laws for A and B arise from the following boundary conditions on $\Pi_A(k)$ and $\Pi_B(k)$:

$$\lim_{k \rightarrow 0^+} \Pi_A(k) = \lim_{k \rightarrow +\infty} \Pi_A(k) = 0 \quad (24)$$

$$\lim_{k \rightarrow 0^+} \Pi_B(k) = \lim_{k \rightarrow +\infty} \Pi_B(k) = 0. \quad (25)$$

2.2. The Danilov inequality

Assuming that the injection (forcing) of A and B occurs in $[k_1, k_2]$, then at steady state, we have, from (19) and (20):

$$\Pi_A(k) = \int_k^{+\infty} D_A(q) dq, \quad \text{for } k > k_2 \quad (26)$$

$$\Pi_B(k) = \int_k^{+\infty} D_B(q) dq, \quad \text{for } k > k_2 \quad (27)$$

$$\Pi_A(k) = - \int_0^k D_A(q) dq, \quad \text{for } 0 < k < k_1 \quad (28)$$

$$\Pi_B(k) = - \int_0^k D_B(q) dq, \quad \text{for } 0 < k < k_1, \quad (29)$$

since $F_A(k) = 0$ and $F_B(k) = 0$ for $0 < k < k_1$ and $k > k_2$. For wavenumbers $k > k_2$, we have therefore

$$L(k)\Pi_A(k) - \Pi_B(k) = \int_k^{+\infty} [L(k) - L(q)]D_A(q) dq < 0. \quad (30)$$

Similarly, for wavenumbers $0 < k < k_1$, we have:

$$L(k)\Pi_A(k) - \Pi_B(k) = - \int_0^k [L(k) - L(q)]D_A(q) dq < 0. \quad (31)$$

Consequently, for all wavenumbers $k \in (0, k_1) \cup (k_2, +\infty)$ not in the forcing range, we have:

$$L(k)\Pi_A(k) - \Pi_B(k) < 0. \quad (32)$$

This inequality was brought to our attention by Danilov (2004, personal communication) for the case of 2D turbulence. We shall see later in this paper that although this inequality has a trivial mathematical derivation in one-layer models, it becomes nontrivial for the two-layer QG model. In all cases, what is interesting is the physical understanding that follows from studying this inequality, and not the mathematical derivations per se.

Previously, Fjroft (1953) and Eyink (1996) derived a similar, but looser, bound, for the downscale energy flux Π_E in two-dimensional turbulence: $\Pi_E(k) < \eta_0/k^2$, involving the total rate of enstrophy injection η_0 . This looser inequality is often used to show (Salmon, 1998) that in two-dimensional turbulence with an infinite downscale range, the energy flux $\Pi_E(k)$, vanishes. For the case of small but finite viscosity where the downscale spectral range is finite, the energy flux does not vanish.

The significance of the inequality (32) is that it decides whether the transition wavenumber k_t is within the inertial range, thus making a transition from the leading cascade to the subleading cascade observable in the energy spectrum $E(k)$. Whether this happens depends on the baroclinicity of the system, as we will show below by considering three different cases: two-dimensional turbulence which is completely barotropic, SQG turbulence which is completely baroclinic, and the two-layer QG model which lies in between with respect to baroclinicity.

3. THE CASE OF 2D TURBULENCE

We begin with the case of 2D turbulence in finite domain with finite viscosity for the infrared and ultraviolet dissipations. This is a generalization of the theory by Kraichnan (1967) of infinite domain with infinitesimal dissipation. In the inertial range on the downscale side of injection, $\Pi_A(k) \approx \epsilon_{uv}$, and $\Pi_B(k) \approx \eta_{uv}$ provided that we ignore a small dissipative contribution. The inequality (32) implies that $\epsilon_{uv}k^2 < \eta_{uv}$ for all k in this inertial range. The energy spectrum in (1), valid in the inertial range, can be rewritten to leading order, omitting the logarithmic correction:

$$E(k) \approx C_1 \epsilon_{uv}^{2/3} k^{-5/3} + C_2 \eta_{uv}^{2/3} k^{-3} \quad (33)$$

$$\approx C_2 \eta_{uv}^{2/3} k^{-3} \left(1 + \frac{C_1}{C_2} \left(\frac{\epsilon_{uv} k^2}{\eta_{uv}} \right)^{2/3} \right) \quad (34)$$

$$\approx C_2 \eta_{uv}^{2/3} k^{-3}, \quad (35)$$

where we use $\epsilon_{uv}k^2 < \eta_{uv}$ and the assumption $C_1 \leq C_2$. This sequence of steps is valid *asymptotically* in the limit of large separation between the forcing scale and the dissipation scale, for wavenumbers k in the inertial range. A similar argument can be made for the inertial range upscale of injection.

It follows that on the downscale side of injection the dominant cascade is the enstrophy cascade with $E(k) \sim k^{-3}$, and on the upscale side of injection the dominant cascade is the inverse energy cascade with $E(k) \sim k^{-5/3}$. By ‘‘dominant’’ we mean that even for finite Reynolds numbers the contributions of the subleading downscale energy cascade and the subleading inverse enstrophy cascade are hidden for *all* the wavenumbers k in the inertial range. A violation of the condition $C_1 \leq C_2$ can allow, in principle, a transition to $k^{-5/3}$ scaling very near the neighborhood of the dissipation range. The Danilov inequality by itself cannot completely rule out such an effect, but so far as we know, it has never been clearly observed in numerical simulations of the enstrophy cascade.

This argument supports the conjecture by Smith (2004) that in 2D turbulence, on the downscale side of injection, we have no transition to shallower scaling $E(k) \sim k^{-5/3}$. His other conjecture, that the same result also holds for the two-layer QG model, is not true in general and will be discussed later in this paper.

It should be noted that in the foregoing arguments it is *assumed* that an inertial range exists either upscale or downscale of injection. Unlike the case of 3D turbulence, where the downscale energy cascade is very robust, it is well known that in 2D turbulence there are circumstances where the leading inverse energy cascade (Danilov, 2005; Danilov and Gurarie, 2001a,b; Gkioulekas and Tung, 2005c) or the leading downscale enstrophy cascade (Tran and Bowman, 2003, 2004; Tran and Shepherd, 2002) may fail to appear as expected. Some of these issues are also relevant to the case of α -turbulence (Tran, 2004). In general, the failure of cascades is to be attributed to the absence of a sufficiently strong large-scale dissipation sink. Since the observational evidence suggests that cascades exist in atmospheric turbulence, we will simply assume that without further discussion.

4. THE CASE OF SQG TURBULENCE

There has been considerable confusion over the physical interpretation of the surface quasi-geostrophic model. Although its mathematical formulation is in the form of a one-layer model, it represents a three-dimensional system that corresponds to the baroclinic limit of the three-dimensional quasi-geostrophic model. Once that is taken into account, the physical interpretation of the spectra $A(k)$ and $B(k)$ and the physical implications of the Danilov inequality have to be revised. The viewpoint that we would like to put forth in this section is that whereas two-dimensional turbulence is an extreme case where the enstrophy cascade is completely dominant, SQG turbulence is the other extreme case where it is the downscale energy cascade that is completely dominant.

4.1. 3D interpretation of SQG turbulence

As derived by Charney (1971), 3D QG flow conserves the 3D potential vorticity ξ , which is advected horizontally by the streamfunction ψ . Here, both ψ and ξ are 3D fields. For con-

stant Coriolis parameter f , the governing conservation law for ξ takes the form:

$$\frac{\partial \xi}{\partial t} + J(\psi, \xi) = 0, \quad (36)$$

with ξ given by

$$\xi = \Delta \psi + \frac{f^2}{\rho_0} \frac{\partial}{\partial z} \left(\frac{\rho_0}{N^2} \frac{\partial \psi}{\partial z} \right) \equiv \mathcal{P}\psi, \quad (37)$$

where $\rho_0(z)$ is the ambient air density, and $N^2(z)$ the Brunt-Väisälä frequency. Here we have omitted the forcing and dissipation terms. The streamfunction ψ is also linked with the potential temperature Θ via the hydrostatic relation

$$\Theta = \frac{g}{N} \frac{T}{T_0} = \frac{f}{N} \frac{\partial \psi}{\partial z}. \quad (38)$$

The potential temperature is also governed by a 2D-like equation

$$\frac{\partial \Theta}{\partial t} + J(\psi, \Theta) + \frac{T_0 N^2}{g} w = 0, \quad (39)$$

where w is the vertical velocity field.

In SQG the potential vorticity ξ is assumed, a priori, to be identically zero for $z > 0$. The streamfunction ψ is solved from $\xi = \mathcal{P}\psi = 0$. With ρ_0 and N^2 taken to be constants, the horizontal Fourier transform of $\psi(x, y, z, t)$ is obtained as

$$\hat{\psi}(\mathbf{k}, z, t) = \hat{\psi}_0(\mathbf{k}, t) e^{-|\mathbf{k}|(N/f)z}, \quad (40)$$

using the boundedness boundary condition as $z \rightarrow \infty$.

Most of the dynamics in this model are occurring at the surface $z = 0$, where the boundary condition of vanishing vertical velocity w applied to the potential temperature (Θ) equation leads to:

$$\frac{\partial \Theta}{\partial t} + J(\psi, \Theta) = \mathcal{D} + \mathcal{F}, \quad (41)$$

where Θ now plays the role of the conserved quantity ζ in (5). Here we have also introduced thermal forcing and dissipation: $\mathcal{D} = \nu \Delta \Theta$ is the thermal diffusion, and

$$\mathcal{F} = Q = \alpha_E (\Theta_0 - \Theta), \quad (42)$$

is the thermal heating in the commonly used form of Newtonian cooling (see Tung and Orlando (2003b)) which includes a ‘‘forcing’’ term $\alpha_E \Theta_0$ and the ‘‘Ekman damping’’ term $-\alpha_E \Theta$. This equation is to be solved on a 2D surface $z = 0$. It has the same form as the vorticity equation for 2D turbulence (e.g. (5)), except that the spectral relationship between the advected quantity Θ and the advecting field ψ is given instead by

$$\hat{\Theta}(\mathbf{k}, z, t) = \frac{f}{N} \frac{\partial}{\partial z} \left(\hat{\psi}_0(\mathbf{k}, t) e^{-|\mathbf{k}|(N/f)z} \right) \quad (43)$$

$$= -|\mathbf{k}| [\hat{\psi}_0(\mathbf{k}, t) e^{-|\mathbf{k}|(N/f)z}] \quad (44)$$

$$= -|\mathbf{k}| \hat{\psi}_0(\mathbf{k}, z, t), \quad (45)$$

which reduces to $\hat{\Theta}(\mathbf{k}, t) = -|\mathbf{k}| \hat{\psi}(\mathbf{k}, t)$ at $z = 0$. Thus SQG corresponds to $L(k) = k$.

It has been shown by Charney (1971), and more generally by Tung and Orlando (2003b), that the 3D QG energy density

$$\mathcal{E} \equiv \frac{1}{2} \rho_0 \left[|\nabla \psi|^2 + \frac{f^2}{N^2} \left(\frac{\partial \psi}{\partial z} \right)^2 \right], \quad (46)$$

is an invariant (i.e. independent of time), *when integrated over the 3D domain*. \mathcal{E} is the sum of the kinetic energy density \mathcal{E}_K and the potential energy density \mathcal{E}_P which are given by

$$\mathcal{E}_K = (1/2) \rho_0 (u^2 + v^2) = (1/2) \rho_0 |\nabla \psi|^2 \quad (47)$$

$$\mathcal{E}_P = \frac{1}{2} \rho_0 \left(\frac{f}{N} \right)^2 \left(\frac{\partial \psi}{\partial z} \right)^2 = \frac{1}{2} \rho_0 \Theta^2. \quad (48)$$

For SQG, using Parseval’s identity, the energies integrated over the horizontal surface are given by

$$E_P = \langle \mathcal{E}_P \rangle = \frac{1}{2} \rho_0 \langle \Theta^2 \rangle, \quad (49)$$

$$E_K = \langle \mathcal{E}_K \rangle = \frac{1}{2} \rho_0 \langle |\nabla \psi|^2 \rangle \quad (50)$$

$$= \frac{1}{2} \rho_0 \int \langle (i\mathbf{k} \hat{\psi}(\mathbf{k}, t)) \cdot (-i\mathbf{k} \hat{\psi}^*(\mathbf{k}, t)) \rangle d\mathbf{k} \quad (51)$$

$$= \frac{1}{2} \rho_0 \int \langle |-k \hat{\psi}(\mathbf{k}, t)|^2 \rangle d\mathbf{k} = \frac{1}{2} \rho_0 \langle \Theta^2 \rangle = E_P. \quad (52)$$

It is thus seen that the kinetic energy density and the available potential energy density, when integrated horizontally, are equipartioned, and that

$$2B \equiv \langle \Theta^2 \rangle = (E_P + E_K) / \rho_0 = E / \rho_0, \quad (53)$$

is the total energy at the lower surface. The 3D energy is, instead

$$\begin{aligned} E_{3D} &\equiv \int_0^\infty \langle \mathcal{E} \rangle dz = \int_0^\infty \rho_0 dz \langle \Theta^2 \rangle \\ &= \int_0^\infty \rho_0 dz \iint dk_x dk_y \langle |\hat{\Theta}|_{z=0}^2 \rangle e^{-2|\mathbf{k}|(N/f)z} \\ &= \frac{1}{2} \rho_0 \iint dk_x dk_y \frac{f}{N|\mathbf{k}|} \langle \hat{\Theta} \hat{\Theta}^* |_{z=0} \rangle \\ &= \frac{1}{2} \rho_0 \iint \frac{f}{N} \langle -\hat{\psi} \hat{\Theta}^* |_{z=0} \rangle dk_x dk_y \\ &= \frac{1}{2} \rho_0 \frac{f}{N} \langle (-\psi \Theta |_{z=0}) \rangle = \rho_0 (f/N) A, \end{aligned}$$

with A defined earlier as $A \equiv (1/2) \langle (-\psi \Theta) \rangle$. Previous authors have made use of the similarity between the form of vorticity equation (5) in 2D turbulence and the temperature equation (41) in SQG turbulence to identify, by analogy, A as the ‘‘energy’’ and B as the ‘‘enstrophy’’ (Held et al., 1995; Pierrehumbert et al., 1994). As pointed out in Tung and Orlando (2003b) and also here, $2B$ is the total energy integrated over the lower surface, and includes kinetic

plus available potential energy. The physical interpretation for A was not given, but can now be seen to be the total energy integrated over the 3D domain. There is no potential enstrophy ($\xi^2/2$) per se in SQG turbulence, because potential vorticity ξ has been taken to be zero identically. Consequently the flux of potential enstrophy in SQG is exactly equal to zero, thus $k^2\Pi_{E_{2D}}(k) - \Pi_G(k) > 0$ and $k^2\Pi_{E_{3D}}(k) - \Pi_G(k) > 0$.

4.2. The energy spectrum in SQG turbulence

The argument of the previous section can be extended to the case of SQG turbulence and to the even wider case of α -turbulence, which includes both the case of SQG for $\alpha = 1$, and the case of 2D for $\alpha = 2$. Here $L(k) = k^\alpha$ and we assume $\alpha > 0$. Using the linear superposition principle discussed in Gkioulekas and Tung (2005b) and assuming the existence of inertial ranges, the spectrum of $A(k)$ and $B(k)$ are, in the downscale inertial range:

$$\begin{aligned} A(k) &\approx C_1(\Pi_A)^{2/3}k^{-\frac{7}{3}+\frac{1}{3}\alpha} + C_2(\Pi_B)^{2/3}k^{-\frac{7}{3}-\frac{1}{3}\alpha} \quad (54) \\ B(k) &= |k|^\alpha A(k). \quad (55) \end{aligned}$$

Here, Π_A and Π_B are the constant fluxes on the downscale side of the forcing range. The inequality (32) becomes $k^\alpha\Pi_A < \Pi_B$ for all k in the inertial ranges. Consequently, for wavenumbers k in the inertial range, the spectrum $A(k)$ is given by

$$\begin{aligned} A(k) &\approx C_2(\Pi_B)^{2/3}k^{-\frac{7}{3}-\frac{1}{3}\alpha} \left(1 + \frac{C_1}{C_2} \left(\frac{\Pi_A k^\alpha}{\Pi_B} \right)^{2/3} \right) \quad (56) \\ &\approx C_2(\Pi_B)^{2/3}k^{-\frac{7}{3}-\frac{1}{3}\alpha}, \quad (57) \end{aligned}$$

again in the limit of large separation between the forcing scale and the dissipation scale and provided that $C_1 \leq C_2$. Thus, in the downscale range, there is no observable transition and therefore:

$$A(k) \approx C_2(\Pi_B)^{2/3}k^{-\frac{7}{3}-\frac{1}{3}\alpha}, \quad (58)$$

$$B(k) \approx C_2(\Pi_B)^{2/3}k^{-\frac{7}{3}+\frac{2}{3}\alpha}. \quad (59)$$

For the SQG model, in the downscale inertial range, the energy spectrum is the same as that for $2B(k)$ in α -turbulence with $\alpha = 1$:

$$E(k) = 2B(k) \approx C_1\varepsilon_{3D}^{2/3}k^{-1} + C_2\varepsilon_{2D}^{2/3}k^{-5/3}. \quad (60)$$

Here ε_{3D} is the downscale energy flux of 3D energy A and ε_{2D} is the downscale energy flux of 2D energy B . From the Danilov inequality we learn that the visible energy spectrum in the inertial range downscale from the injection scale is given by

$$E(k) \approx C_2\varepsilon_{2D}^{2/3}k^{-5/3}. \quad (61)$$

This $k^{-5/3}$ energy spectrum is now predicted by our theory. The flux ε_{2D} is not the ‘‘enstrophy’’ flux but is the 2D flux of

2D energy $2B = E$ at the $z = 0$ layer. Numerical simulations have reproduced this energy cascade and appear to indicate a small deviation from the $k^{-5/3}$ slope due to conjectured intermittency corrections (Celani et al., 2004).

5. THE CASE OF TWO-LAYER QG TURBULENCE

The results in the previous sections demonstrate that QG turbulence can exhibit a variety of behaviors. Barotropic models usually possess an energy spectrum with -3 spectral slope, as in 2D turbulence, while SQG turbulence, which is baroclinic (with its exponential decay with height), is expected to have a spectrum with a $-5/3$ slope, in the downscale inertial range. Two-layer QG models have both a barotropic and a baroclinic component (see Salmon (1978, 1980, 1998)), and we therefore expect a mixture of -3 and $-5/3$ slopes depending on the degree of baroclinicity. In terms of the Danilov inequality, we expect that it will hold for small wavenumbers and fail for large wavenumbers.

A relatively realistic two-layer model applicable to studying atmospheric turbulence in the troposphere was adopted in Tung and Orlando (2003a). In this model forcing is due to thermal heating, which injects energy directly into the baroclinic part of the total energy. The two-layer fluid sits atop of an Ekman boundary layer near the ground, which introduces Ekman pumping in the lower layer (Holton, 1979) but *not* in the upper layer. If one artificially adds an identical Ekman damping in the upper layer it can be easily shown that Danilov’s inequality applies, and we leave the proof to the interested reader. Here we will discuss the case of asymmetric Ekman damping.

For the multiple-layer QG model, the governing equations can be rewritten in the form of the conservation law with $\zeta = -\mathcal{L}\psi$, if we make \mathcal{L} into a matrix and ψ into a column vector. We will discuss the more general theory of multi-layers in a future paper. Here we only wish to explain why and how the Danilov inequality can fail in the two-layer QG model.

5.1. The formal setup

Two-layer QG models conserve potential vorticity in each layer in the absence of forcing and damping. In the forced-dissipative case, the governing equations read:

$$\text{Top layer: } \frac{\partial \zeta_1}{\partial t} + J(\psi_1, \zeta_1) = \mathcal{D}_1 + \mathcal{F}_1 \quad (62)$$

$$\text{Bottom layer: } \frac{\partial \zeta_2}{\partial t} + J(\psi_2, \zeta_2) = \mathcal{D}_2 + \mathcal{F}_2, \quad (63)$$

where

$$\zeta_1 = \Delta\psi_1 - \frac{k_R^2}{2}(\psi_1 - \psi_2), \quad \zeta_2 = \Delta\psi_2 + \frac{k_R^2}{2}(\psi_1 - \psi_2),$$

are the potential vorticity in each layer. $k_R \equiv (2\sqrt{2}f)/(hN)$ is the Rossby radius of deformation wavenumber and is taken as a given constant (h is the height). The dissipation terms, \mathcal{D}_i , include momentum dissipation of relative vorticity, $\Delta\psi_i$,

in each layer, and Ekman damping from the lower boundary layer:

$$\mathcal{D}_1 = \nu(-\Delta)^{p+1}\psi_1, \quad \mathcal{D}_2 = \nu(-\Delta)^{p+1}\psi_2 - \nu_E\Delta\psi_2. \quad (64)$$

The forcing terms can be shown to satisfy

$$\mathcal{F}_1 = +\frac{k_R^2}{2f}Q, \quad \mathcal{F}_2 = -\frac{k_R^2}{2f}Q, \quad (65)$$

where Q is the radiative heating term in the temperature equation.

The two inviscid quadratic invariants are the energy E and the total layer potential enstrophies G_1 and G_2 given by

$$E \equiv \langle\langle \psi_1\zeta_1 + \psi_2\zeta_2 \rangle\rangle \quad (66)$$

$$G_1 \equiv \langle\langle \zeta_1^2 \rangle\rangle, \quad G_2 \equiv \langle\langle \zeta_2^2 \rangle\rangle. \quad (67)$$

The energy and enstrophy spectra are defined as

$$E(k) \equiv \frac{d}{dk} \langle\langle \psi_1^{<k} \zeta_1^{<k} \rangle\rangle + \frac{d}{dk} \langle\langle \psi_2^{<k} \zeta_2^{<k} \rangle\rangle, \quad (68)$$

$$G_1(k) \equiv \frac{d}{dk} \langle\langle \zeta_1^{<k} \zeta_1^{<k} \rangle\rangle, \quad (69)$$

$$G_2(k) \equiv \frac{d}{dk} \langle\langle \zeta_2^{<k} \zeta_2^{<k} \rangle\rangle, \quad (70)$$

and the total enstrophy spectrum $G(k)$ is $G(k) = G_1(k) + G_2(k)$. We define the streamfunction spectra

$$U_1(k) \equiv \frac{d}{dk} \langle\langle \psi_1^{<k} \psi_1^{<k} \rangle\rangle, \quad (71)$$

$$U_2(k) \equiv \frac{d}{dk} \langle\langle \psi_2^{<k} \psi_2^{<k} \rangle\rangle, \quad (72)$$

$$C(k) \equiv \frac{d}{dk} \langle\langle \psi_1^{<k} \psi_2^{<k} \rangle\rangle, \quad (73)$$

and also $U(k) \equiv U_1(k) + U_2(k)$. It is easy to show the triangle inequality $2C(k) \leq U(k)$. The energy and enstrophy spectra are related with the streamfunction spectra via

$$E(k) = (k^2 + k_R^2/2)U(k) - k_R^2C(k) \quad (74)$$

$$G(k) = (k^4 + k^2k_R^2 + k_R^4/2)U(k) - k_R^2(2k^2 - k_R^2)C(k). \quad (75)$$

From the standpoint of examining the Danilov inequality, it is most convenient to work with the streamfunction spectra. However, following Salmon (1978, 1980, 1998), for physical understanding it is useful to work with the energy and enstrophy spectra. Furthermore, it is helpful to distinguish between barotropic energy and baroclinic energy as follows: Let $\psi \equiv (\psi_1 + \psi_2)/2$ and $\tau \equiv (\psi_1 - \psi_2)/2$. So, $\psi_1 = \psi + \tau$ and $\psi_2 = \psi - \tau$. Now we define three spectra $E_K(k)$, $E_P(k)$, and $E_C(k)$ in terms of ψ and τ :

$$E_K(k) \equiv 2k^2 \frac{d}{dk} \langle\langle \psi^{<k} \psi^{<k} \rangle\rangle, \quad (76)$$

$$E_P(k) \equiv 2(k^2 + k_R^2) \frac{d}{dk} \langle\langle \tau^{<k} \tau^{<k} \rangle\rangle, \quad (77)$$

$$E_C(k) \equiv 2k^2 \frac{d}{dk} \langle\langle \psi^{<k} \tau^{<k} \rangle\rangle. \quad (78)$$

Here $E_K(k)$ is the barotropic energy spectrum and $E_P(k)$ the baroclinic energy spectrum. It is easy to show that the definitions are self-consistent, i.e. $E(k) = E_K(k) + E_P(k)$. The relation between the energy spectra and the enstrophy spectra can be now written in terms of two constraint equations:

$$G_1(k) = (1/2)[k^2E_K(k) + (k^2 + k_R^2)E_P(k)] + (k^2 + k_R^2)E_C(k) \quad (79)$$

$$G_2(k) = (1/2)[k^2E_K(k) + (k^2 + k_R^2)E_P(k)] - (k^2 + k_R^2)E_C(k). \quad (80)$$

Thus we see that the physical interpretation of $E_C(k)$ is that it represents the difference in potential enstrophy distribution between the two layers, and it is given by

$$E_C(k) = \frac{G_1(k) - G_2(k)}{2(k^2 + k_R^2)}. \quad (81)$$

5.2. Controlled necessary condition

The right-hand-side of the Danilov inequality reads

$$k^2\Pi_E(k) - \Pi_G(k) = \int_k^{+\infty} dq (k^2D_E(q) - D_G(q)) \equiv \int_k^{+\infty} dq \Delta(k, q), \quad (82)$$

so the key question is whether $\Delta(k, q)$ is positive or negative for wavenumbers $k < q < k_{max}$. Here k_{max} is either the truncation wavenumber in the numerical model, or, in the theoretical case of infinite resolutions, is the hyperviscosity dissipation wavenumber, beyond which the spectral enstrophy dissipation rate becomes negligible.

With some straightforward but tedious calculations, it can be shown that the dissipation rate $D_E(k)$ for the total energy, and the dissipation rate $D_G(k)$ for the total potential enstrophy are given by

$$D_E(k) = 2\nu k^{2p+2}U(k) + 2\nu_E k^2 U_2(k) \quad (83)$$

$$D_G(k) = 2\nu k^{2p+2}E(k) + \nu_E k^2 [(2k^2 + k_R^2)U_2(k) - k_R^2 C(k)], \quad (84)$$

thus it follows that

$$\Delta(k, q) = k^2 D_E(q) - D_G(q) = 2\nu q^{2p+2} [k^2 U(q) - E(q)] + \nu_E k_R^2 q^2 C(q) + \nu_E q^2 (2k^2 - 2q^2 - k_R^2) U_2(q) \quad (85)$$

$$= 2\nu q^{2p+2} [(k^2 - q^2)U(q) + (k_R^2/2)(2C(q) - U(q))] + \nu_E k_R^2 q^2 (C(q) - U_2(q)) + 2\nu_E q^2 (k^2 - q^2) U_2(q). \quad (86)$$

The first and third terms in this expression are always negative. Consequently, a *necessary* condition for *violating* the Danilov inequality is that the second term has to be positive, i.e. $C(q) - U_2(q) \geq 0$. Otherwise, if $C(q) - U_2(q) \leq 0$, then the Danilov inequality will be satisfied. A physical interpretation of this condition will be given in section 5.4.

A controlled sufficient condition to satisfy the Danilov inequality can be derived in terms of the physical parameters of the problem by noting that $C(q) - U_2(q) \leq (1/2)(U_1(q) - U_2(q))$. It follows that

$$\Delta(k, q) = 2\nu q^{2p+2} [(k^2 - q^2)U(q) + (k_R^2/2)(2C(q) - U(q))] + \nu_E k_R^2 q^2 (C(q) - U_2(q)) + 2\nu_E q^2 (k^2 - q^2) U_2(q) \quad (87)$$

$$\leq 2\nu q^{2p+2} [(k^2 - q^2)U(q) + (k_R^2/2)(2C(q) - U(q))] + \nu_E k_R^2 q^2 (1/2)(U_1(q) - U_2(q)) + 2\nu_E q^2 (k^2 - q^2) U_2(q) \quad (88)$$

$$= [2\nu q^{2p+2} (k^2 - q^2) + (1/2)\nu_E k_R^2 q^2] U_1(q) + [2(\nu q^{2p+2} + \nu_E q^2)(k^2 - q^2) - (1/2)\nu_E k_R^2 q^2] U_2(q) \quad (89)$$

$$\leq [2\nu q^{2p+2} (k^2 - q^2) + (1/2)\nu_E k_R^2 q^2] U_1(q). \quad (90)$$

Here, we have used the inequality $2C(k) \leq U(k)$ to eliminate the $(k_R^2/2)(2C(q) - U(q))$ term. We have also eliminated the $U_2(q)$ term because it is unconditionally negative. This leads to the following controlled *sufficient* condition to *satisfy* Danilov's inequality:

$$\nu_E < 4\nu k_{\max}^{2p} \left(\frac{k_{\max}}{k_R} \right)^2. \quad (91)$$

Equivalently, a *necessary* condition to *violate* Danilov's inequality is

$$\nu_E > 4\nu k_{\max}^{2p} \left(\frac{k_{\max}}{k_R} \right)^2. \quad (92)$$

It is interesting to note that in the numerical simulation of the two-layer model the algorithm adopted by Tung and Orlando (2003a) for determining the magnitude of the hyperviscosity coefficient is $\nu_E \gg \nu k_{\max}^{2p}$, for all but the last twenty wavenumbers k in the dissipation range. Tung and Orlando (2003a) obtained an energy spectrum with the compound slope configuration and the transition wavenumber k_t occurred in the inertial range downscale from injection in agreement with the condition $k_t \approx \sqrt{\eta_{uv}/\epsilon_{uv}}$, thus implying a violation of Danilov's inequality.

5.3. An uncontrolled necessary and sufficient condition

The question now arises: is it possible to derive a sufficient condition to violate the Danilov inequality of the form $\nu_E k_R^2 \geq \Lambda \nu k_{\max}^{2p+2}$ for some universal constant Λ ? So far as we know, this is not possible. However, it is possible to derive an *uncontrolled* necessary and sufficient condition for violating the Danilov inequality.

We begin with defining

$$2C(q) = \lambda(q)U(q) \quad (93)$$

$$U_2(q) = u(q)U(q) \quad (94)$$

Here $0 \leq u(q) \leq 1$ and $-1 \leq \lambda(q) \leq 1$. We may thus rewrite everything in terms of $U(q)$ by employing

$$C(q) - U_2(q) = (\lambda(q)/2 - u(q))U(q) \quad (95)$$

$$2C(q) - U(q) = (\lambda(q) - 1)U(q) \quad (96)$$

Then we can rewrite $\Delta(k, q)$, as follows:

$$\Delta(k, q) = 2\nu q^{2p+2} [(k^2 - q^2)U(q) + (k_R^2/2)(2C(q) - U(q))] + \nu_E k_R^2 q^2 (C(q) - U_2(q)) + 2\nu_E q^2 (k^2 - q^2) U_2(q) \quad (97)$$

$$= q^2 U(q) [-\nu q^{2p} (2(q^2 - k^2) + k_R^2(1 - \lambda(q))) + \nu_E k_R^2 (\lambda(q)/2 - u(q)) + 2(k/k_R)^2 u(q) - 2(q/k_R)^2 u(q)]. \quad (98)$$

It easy to see that a sufficient condition to get $\Delta(k, q) \geq 0$ is

$$\nu_E k_R^2 (\lambda(q)/2 - u(q) + 2(k/k_R)^2 u(q) - 2(q/k_R)^2 u(q)) \geq \nu q^{2p} (2(q^2 - k^2) + k_R^2 (1 - \lambda(q))). \quad (99)$$

The necessary condition to violate the Danilov inequality $C(q) - U_2(q) \geq 0$, which was derived previously, implies that $\lambda(q)/2 - u(q) \geq 0$. Provided that we assume the stronger condition

$$\lambda(q)/2 - u(q) + 2(k/k_R)^2 u(q) - 2(q/k_R)^2 u(q) \geq 0, \quad (100)$$

we may rewrite our sufficient condition as:

$$\frac{\nu_E k_R^2}{\nu q^{2p}} \geq \frac{2(q^2 - k^2) + k_R^2 (1 - \lambda(q))}{\lambda(q)/2 - u(q) + 2(k/k_R)^2 u(q) - 2(q/k_R)^2 u(q)}. \quad (101)$$

To violate the Danilov inequality at wavenumber k , this sufficient condition must hold for all q such that $k < q < k_{max}$. Since the numerator is always positive, the condition (100) is in fact a stronger necessary condition for violating the Danilov inequality.

5.4. Physical interpretation of necessary conditions

We would like now to discuss the plausibility of the necessary condition $\lambda(q)/2 - u(q) > 0$ and the stronger necessary condition (100). To this end, we rewrite these conditions equivalently in terms of the physical energy and enstrophy spectra.

It is easy to write $U_1(k)$, $U_2(k)$, and $C(k)$ in terms of $E_K(k)$, $E_P(k)$, and $E_C(k)$:

$$U_1(k) = \frac{d}{dk} \langle \langle (\psi + \tau)^{<k} (\psi + \tau)^{<k} \rangle \rangle \quad (102)$$

$$= \frac{d}{dk} \langle \langle \psi^{<k} \psi^{<k} \rangle \rangle + 2 \frac{d}{dk} \langle \langle \psi^{<k} \tau^{<k} \rangle \rangle + \frac{d}{dk} \langle \langle \tau^{<k} \tau^{<k} \rangle \rangle \quad (103)$$

$$= \frac{E_K(k)}{2k^2} + \frac{E_P(k)}{2(k^2 + k_R^2)} + \frac{E_C(k)}{k^2}, \quad (104)$$

and with a similar argument we find

$$U_2(k) = \frac{E_K(k)}{2k^2} + \frac{E_P(k)}{2(k^2 + k_R^2)} - \frac{E_C(k)}{k^2} \quad (105)$$

$$C(k) = \frac{E_K(k)}{2k^2} - \frac{E_P(k)}{2(k^2 + k_R^2)}. \quad (106)$$

We may thus write $\lambda(k)$ and $u(k)$ in terms of $E_K(k)$, $E_P(k)$, and $E_C(k)$:

$$\lambda(k) = \frac{2C(k)}{U(k)} = 2 \frac{\frac{E_K(k)}{2k^2} - \frac{E_P(k)}{2(k^2 + k_R^2)}}{\frac{E_K(k)}{2k^2} + \frac{E_P(k)}{2(k^2 + k_R^2)}} \quad (107)$$

$$= \frac{(k^2 + k_R^2)E_K(k) - k^2 E_P(k)}{(k^2 + k_R^2)E_K(k) + k^2 E_P(k)}, \quad (108)$$

$$(109)$$

and

$$u(k) = \frac{U_2(k)}{U(k)} = \frac{\frac{E_K(k)}{2k^2} + \frac{E_P(k)}{2(k^2 + k_R^2)} - \frac{E_C(k)}{k^2}}{\frac{E_K(k)}{2k^2} + \frac{E_P(k)}{2(k^2 + k_R^2)}} \quad (110)$$

$$= \frac{1}{2} \frac{(k^2 + k_R^2)E_K(k) + k^2 E_P(k) - 2(k^2 + k_R^2)E_C(k)}{(k^2 + k_R^2)E_K(k) + k^2 E_P(k)}. \quad (111)$$

and the necessary condition $\lambda(q)/2 - u(q) \geq 0$ can now be rewritten as

$$\lambda(q)/2 - u(q) = \frac{(q^2 + k_R^2)E_C(q) - q^2 E_P(q)}{(q^2 + k_R^2)E_K(q) + q^2 E_P(q)} > 0. \quad (112)$$

The denominator is obviously positive, consequently the condition is equivalent to

$$G_1(q) - G_2(q) = 2(q^2 + k_R^2)E_C(q) > 2q^2 E_P(q), \quad (113)$$

for all q such as $k < q < k_{max}$.

This is a very interesting result. The requirement, in part, is that there should be more enstrophy on the top layer than the bottom layer, i.e. $G_1(q) > G_2(q)$. It is reasonable to expect this if there is more dissipation on the bottom layer than the top layer. This is the case for the model we are considering where there is Ekman damping at the bottom layer but not at the top layer. However, the actual condition also requires that the difference should be larger than the potential energy $E_P(q)$ multiplied with $2q^2$. In our model, as has been pointed out by Salmon (1978, 1980, 1998), the energy is injected into the system at large scales as baroclinic energy, and most of this energy is converted into barotropic energy near the Rossby wavenumber k_R . It is therefore reasonable to expect that this necessary condition will be satisfied for wavenumbers $q > k_R$.

It should be noted that this is only a necessary condition. A sufficient condition would require furthermore that equation (101) be valid. The necessary precondition for that is the stronger requirement

$$\frac{\lambda(q)/2 - u(q)}{u(q)} > 2 \left(\frac{q}{k_R} \right)^2 - 2 \left(\frac{k}{k_R} \right)^2. \quad (114)$$

This condition can be rewritten equivalently as

$$\frac{G_1(q) - G_2(q) - 2q^2 E_P(q)}{2G_2(q) + k_R^2 (E_K(q) - E_P(q))} > 2 \left(\frac{q}{k_R} \right)^2 - 2 \left(\frac{k}{k_R} \right)^2. \quad (115)$$

and it can be simplified further to give

$$G_1(q) - \left(1 + 4 \frac{q^2 - k^2}{k_R^2} \right) G_2(q) > 2q^2 E_K(q) - 2k^2 (E_K(q) - E_P(q)). \quad (116)$$

The constraint $u(q) \geq 0$ implies that

$$2G_2(q) + k_R^2(E_K(q) - E_P(q)) \geq 0, \quad (117)$$

and subtracting this inequality from the necessary condition above gives the following simplification:

$$G_1(q) - (1 + 4(q/k_R)^2)G_2(q) > 2q^2E_K(q), \quad (118)$$

for all q such that $k < q < k_{max}$.

This condition, which is also a necessary condition for violating the Danilov inequality, places an even stronger constraint on the difference between the enstrophy between the two layers. The fly in the ointment is the $4(q/k_R)^2G_2(q)$ term. If we want to go from the transition scale at about 700km in wavelength down to length scales of a few kilometers, then $4(q/k_R)^2$ increases at least by four orders of magnitude. In the numerical simulations of Tung and Orlando (2003a), the variability of $4(q/k_R)^2$ is relatively small because they only go down to 100km in wavelength. We may therefore conjecture that a violation of the Danilov inequality requires that the ratio $(k_d/k_R)^2$, where k_d is the dissipation wavenumber, must be small. In other words, the subleading downscale energy cascade is expected to be observable in a numerical simulation with small separation of scales between k_R and k_d , but not in a numerical simulation where the separation between these scales is large. A trend of diminishing downscale energy flux with increasing numerical resolution, and thus with increasing separation of scales, has already been seen in the simulations of Tung and Orlando (2003a).

It should be noted, of course, that the quasi-geostrophic model is not valid at length scales much smaller than 100km in wavelength. When three-dimensional effects become relevant the conservation of enstrophy is violated by the nonlinearity itself which reflects itself in additional nonlinear dissipative contributions to $D_G(k)$. These terms make it all the more easier to violate the Danilov inequality for wavenumbers k that correspond to scales less than 100km in wavelength, thereby preserving the very extensive downscale energy cascade observed in the Nastrom-Gage spectrum.

6. CONCLUSIONS AND DISCUSSION

The classical KLB theory of 2D turbulence relies for its mathematical simplicity and elegance on two unrealistic assumptions: that the domain is infinite, and that the Reynolds number approaches infinity. When these two assumptions are relaxed, the situation becomes more complicated. The downscale enstrophy cascade is accompanied with a hidden downscale energy cascade, and similarly the inverse energy cascade is accompanied with a hidden inverse enstrophy cascade. This is true as long as the leading cascades themselves exist, which requires the presence of a sufficiently strong dissipation sink at small wavenumbers. The fluxes associated with the subleading cascades are constrained by the Danilov inequality, and as a result the subleading cascades cannot contribute large enough terms to the energy spectrum to create an observable effect. This picture represents a generalization

of the KLB theory to finite inertial ranges and finite dissipations. This situation changes, however, in baroclinic models of quasi-geostrophic turbulence.

The surface quasi-geostrophic model represents an extreme baroclinic case where the entire behavior in the three-dimensional domain is constrained by the behavior of the system at the $z = 0$ layer. In this model there is no enstrophy, and the dominant feature is the downscale energy cascade.

We have shown that in the two-layer quasi-geostrophic model, the violation of the Danilov inequality is possible only as a result of asymmetric Ekman damping operating on only one of the two-layers. This creates an imbalance between the amount of enstrophy accumulated in one layer versus the amount accumulated in the other layer, and the downscale energy cascade will become observable on the condition that this imbalance is sufficiently large. We have derived in the present paper a sufficient condition for *not* violating the Danilov inequality which explains why the $k^{-5/3}$ spectrum has not been observed in some of the previous simulations of the two-layer model. We have also derived a necessary and sufficient condition for violating the Danilov inequality, but it is an uncontrolled condition. The numerical simulation by Tung and Orlando (2003a) has confirmed that a double cascade with the transition wavenumber located in the inertial range can be realized. This can only occur when the Danilov inequality is violated for some wavenumbers k in the inertial range. The parameterization of the Ekman damping in that simulation does in fact satisfy the necessary condition derived in this paper.

As long as we operate within the framework of multiple-layer models with a finite number of layers, one cannot rule out the alternative theory that the atmospheric energy spectrum might reflect a double downscale cascade of helicity and energy instead of enstrophy and energy (see discussion in section 6.5 of Branover et al. (1999), and figure 3 of Bershadskii et al. (1993)). However, most of the current debate has been focused on the somewhat mysterious nature of the very extensive and robust $k^{-5/3}$ spectrum.

Our work in the present paper explains why it can be reproduced in numerical simulations that use baroclinic models, while the same effect cannot be realized in simulations of two-dimensional turbulence. On the other hand our work here does not rule out the possibility that the shallower part of the spectrum observed by Nastrom and Gage (1984) over the mesoscales can be due to dynamics other than QG, whether it is barotropic or baroclinic, especially for scales of 100 km or less (see e.g. Lindborg (2005) with Boussinesq dynamics). Our present work serves to point out that over the larger scales ($\gtrsim 600$ km), where the transition to a shallower spectrum occurs, baroclinic QG theory by itself is a viable mechanism for explaining the transition from -3 to $-5/3$ slopes.

Furthermore, as proposed first by Tung and Orlando (2003a), the downscale energy flux, which is important in explaining the $k^{-5/3}$ energy spectrum over the mesoscales in most theories, originates at larger scales (the synoptic scales). Its contribution to the energy spectrum is hidden for smaller wavenumbers under the k^{-3} part of the spectrum, and then emerges for larger k past the transition scale. It remains an

open question, one that is beyond the scope of this paper, to explain how this downscale energy flux can be continued into length scales too small for QG theory to describe, and how it is eventually dissipated.

Acknowledgments

It is a pleasure to thank Dr. Sergey Danilov for his e-mail correspondence which led us to the development of the ideas in this paper. Constructive comments from an anonymous reviewer are also gratefully acknowledged. The research is supported by the National Science Foundation, under grants DMS-03-27658 and ATM-01-32727.

References

- Arad, I., V. L'vov, and I. Procaccia, 1999: Correlation functions in isotropic and anisotropic turbulence: The role of the symmetry group. *Phys. Rev. E*, **59**, 6753–6765.
- Batchelor, G., 1947: Kolmogorov's theory of locally isotropic turbulence. *Proc. Camb. Phil. Soc.*, **43**, 533–559.
- 1969: Computation of the energy spectrum in homogeneous, two dimensional turbulence. *Phys. Fluids Suppl. II*, **12**, 233–239.
- Bershanskii, A., E. Kit, and A. Tsinober, 1993: Spontaneous breaking of reflexional symmetry in real quasi-two-dimensional turbulence: stochastic travelling waves and helical solitons in atmosphere and laboratory. *Proc. R. Soc. Lond. A*, **441**, 147–155.
- Biferale, L. and I. Procaccia, 2005: Anisotropy in turbulent flows and in turbulent transport. *Phys. Rep.*, **414**, 43–164.
- Boffetta, G., A. Celani, and M. Vergassola, 2000: Inverse energy cascade in two dimensional turbulence: deviations from gaussian behavior. *Phys. Rev. E*, **61**, 29–32.
- Borue, V., 1994: Inverse energy cascade in stationary two dimensional homogeneous turbulence. *Phys. Rev. Lett.*, **72**, 1475–1478.
- Bowman, J., 1996: On inertial-range scaling laws. *J. Fluid. Mech.*, **306**, 167–181.
- Branover, H., A. Eidelman, E. Golbraikh, and S. Moiseev, 1999: *Turbulence and structures: chaos, fluctuations, and helical self organization in nature and the laboratory*. Academic Press, San Diego.
- Celani, A., M. Cencini, A. Mazzino, and M. Vergassola, 2004: Active and passive fields face to face. *New J. Phys.*, **6**, 72.
- Charney, J., 1948: On the scale of atmospheric motions. *Geophys. Publ.*, **17**, 1–17.
- 1971: Geostrophic turbulence. *J. Atmos. Sci.*, **28**, 1087–1095.
- Chen, Q., S. Chen, and G. Eyink, 2003: The joint cascade of energy and helicity in three-dimensional turbulence. *Phys. Fluids*, **15**, 361–374.
- Cho, J. and E. Lindborg, 2001: Horizontal velocity structure functions in the upper troposphere and lower stratosphere. 1. Observations. *J. Geophys. Res.*, **106 D10**, 10,223–10,232.
- Danilov, S., 2005: Non-universal features of forced 2d turbulence in the energy and enstrophy ranges. *Discrete and Continuous Dynamical Systems B*, **5**, 67–78.
- Danilov, S. and D. Gurarie, 2001a: Forced two-dimensional turbulence in spectral and physical space. *Phys. Rev. E*, **63**, 061208.
- 2001b: Non-universal features of forced two dimensional turbulence in the energy range. *Phys. Rev. E*, **63**, 020203.
- Dewan, E., 1979: Stratospheric spectra resembling turbulence. *Science*, **204**, 832–835.
- Eyink, G., 1996: Exact results on stationary turbulence in two dimensions: Consequences of vorticity conservation. *Physica D*, **91**, 97–142.
- 2001: Dissipation in turbulent solutions of 2D Euler equations. *Nonlinearity*, **14**, 787–802.
- Fjørtoft, R., 1953: On the changes in the spectral distribution of kinetic energy for two dimensional non-divergent flow. *Tellus*, **5**, 225–230.
- Gage, K., 1979: Evidence for a $k^{-5/3}$ law inertial range in mesoscale two-dimensional turbulence. *J. Atmos. Sci.*, **36**, 1950–1954.
- Gage, K. and G. Nastrom, 1986: Theoretical interpretation of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft during GASP. *J. Atmos. Sci.*, **43**, 729–740.
- Gkioulekas, E., 2005: Remarks on the Frisch framework of hydrodynamic turbulence and the quasi-lagrangian formulation, submitted to *Physica D*, [nlin.CD/0506064].
- Gkioulekas, E. and K. Tung, 2005a: A new proof on net upscale energy cascade in 2D and QG turbulence, submitted to *J. Fluid Mech.*, [nlin.CD/0512005].
- 2005b: On the double cascades of energy and enstrophy in two dimensional turbulence. Part 1. Theoretical formulation. *Discrete and Continuous Dynamical Systems B*, **5**, 79–102.
- 2005c: On the double cascades of energy and enstrophy in two dimensional turbulence. Part 2. Approach to the KLB limit and interpretation of experimental evidence. *Discrete and Continuous Dynamical Systems B*, **5**, 103–124.
- Hasegawa, A., C. MacLennan, and Y. Kodama, 1979: Nonlinear behaviour and turbulence spectra of drift waves and Rossby waves. *Phys. Fluids*, **22**, 2122–2129.
- Hasegawa, A. and K. Mima, 1978: Pseudo-three-dimensional turbulence in magnetized nonuniform plasma. *Phys. Fluids*, **21**, 87–92.
- Held, I. M., R. T. Pierrehumbert, S. T. Garner, and K. Swanson, 1995: Surface quasi-geostrophic dynamics. *J. Fluid. Mech.*, **282**, 1–20.
- Holton, J. R., 1979: *An introduction to dynamic meteorology*. Academic Press, San Diego.
- Ishihara, T. and Y. Kaneda, 2001: Energy spectrum in the enstrophy transfer range of two-dimensional forced turbulence. *Phys. Fluids*, **13**, 544–547.
- Kolmogorov, A., 1941a: Dissipation of energy in the locally isotropic turbulence. *Dokl. Akad. Nauk. SSSR*, **32**, 16–18, english translation published in volume 434 of *Proc. R. Soc. Lond. A*.
- 1941b: The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *Dokl. Akad. Nauk. SSSR*, **30**, 301–305, english translation published in volume 434 of *Proc. R. Soc. Lond. A*.
- Koshyk, J. and K. Hamilton, 2001: The horizontal kinetic energy spectrum and spectral budget simulated by a high-resolution troposphere-stratosphere mesosphere GCM. *J. Atmos. Sci.*, **58**, 329–348.
- Koshyk, J., K. Hamilton, and J. Mahlman, 1999: Simulation of the $k^{-5/3}$ mesoscale spectral regime in the GFDL SKYHI general circulation model. *Geophys. Res. Lett.*, **26**, 843–846.
- Kraichnan, R., 1967: Inertial ranges in two dimensional turbulence. *Phys. Fluids*, **10**, 1417–1423.
- 1971: Inertial range transfer in two and three dimensional turbulence. *J. Fluid. Mech.*, **47**, 525–535.
- Leith, C., 1968: Diffusion approximation for two dimensional turbulence. *Phys. Fluids*, **11**, 671–673.
- Lilly, D., 1989: Two dimensional turbulence generated by energy sources at two scales. *J. Atmos. Sci.*, **46**, 2026–2030.
- Lindborg, E., 1999: Can the atmospheric kinetic energy spectrum be explained by two dimensional turbulence? *J. Fluid. Mech.*, **388**, 259–288.
- 2005: The effect of rotation on the mesoscale energy cascade in

- the free atmosphere. *Geophys. Res. Lett.*, **32**, L01809.
- Lindborg, E. and K. Alvelius, 2000: The kinetic energy spectrum of two dimensional enstrophy turbulence cascade. *Phys. Fluids*, **12**, 945–947.
- Merilees, P. and T. Warn, 1975: On energy and enstrophy exchanges in two-dimensional non-divergent flow. *J. Fluid. Mech.*, **69**, 625–630.
- Moffatt, H., 1986: *Advances in Turbulence*, G.Comte-Bellot and J. Mathieu, eds., Springer-Verlag, Berlin, 284.
- Moiseev, S. and O. Chkhetiani, 1996: Helical scaling in turbulence. *JETP*, **83**, 192–198.
- Nastrom, G. and K. Gage, 1984: A climatology of atmospheric wave number spectra of wind and temperature observed by commercial aircraft. *J. Atmos. Sci.*, **42**, 950–960.
- Nastrom, G., K. Gage, and W. Jasperson, 1984: The atmospheric kinetic energy spectrum, $10^0 - 10^4$ km. *Nature*, **310**, 36–38.
- Pasquero, C. and G. Falkovich, 2002: Stationary spectrum of vorticity cascade in two dimensional turbulence. *Phys. Rev. E*, **65**, 056305.
- Pierrehumbert, R. T., I. M. Held, and K. L. Swanson, 1994: Spectra of local and non-local two-dimensional turbulence. *Chaos, Solitons and Fractals*, **4**, 1111–1116.
- Polyakov, A., 1993: The theory of turbulence in two dimensions. *Nuclear Physics B*, **396**, 367–385.
- Saffman, P., 1971: On the spectrum and decay of random two dimensional vorticity distributions at large Reynolds number. *Stud. Appl. Math.*, **50**, 377.
- Salmon, R., 1978: Two-layer quasi-geostrophic turbulence in a simple special case. *Geophys. Astrophys. Fluid Dyn.*, **10**, 25–52.
- 1980: Baroclinic instability and geostrophic turbulence. *Geophys. Astrophys. Fluid Dyn.*, **15**, 167–211.
- 1998: *Lectures on Geophysical Fluid Dynamics*. Oxford University Press, New York.
- Skamarock, W., 2004: Evaluating mesoscale NWP models using kinetic energy spectra. *Month. Weath. Rev.*, **132**, 3019–3032.
- Smith, K., 2004: Comment on: The k^{-3} and $k^{-5/3}$ energy spectrum of the atmospheric turbulence: quasi-geostrophic two level model simulation. *J. Atmos. Sci.*, **61**, 937–942.
- Tabeling, P., 2002: Two dimensional turbulence: a physicist approach. *Phys. Rep.*, **362**, 1–62.
- Tran, C., 2004: Nonlinear transfer and spectral distribution of energy in α turbulence. *Physica D*, **191**, 137–155.
- Tran, C. and J. Bowman, 2003: On the dual cascade in two dimensional turbulence. *Physica D*, **176**, 242–255.
- 2004: Robustness of the inverse cascade in two-dimensional turbulence. *Phys. Rev. E*, **69**, 036303.
- Tran, C. and T. Shepherd, 2002: Constraints on the spectral distribution of energy and enstrophy dissipation in forced two dimensional turbulence. *Physica D*, **165**, 199–212.
- Tung, K., 2004: Reply to comments by K. Shafer Smith. *J. Atmos. Sci.*, **61**, 943–948.
- Tung, K. and W. Orlando, 2003a: The k^{-3} and $k^{-5/3}$ energy spectrum of the atmospheric turbulence: quasi-geostrophic two level model simulation. *J. Atmos. Sci.*, **60**, 824–835.
- 2003b: On the differences between 2D and QG turbulence. *Discrete and Continuous Dynamical Systems B*, **3**, 145–162.
- Tung, K. and W. Welch, 2001: Remarks on note on geostrophic turbulence. *J. Atmos. Sci.*, **58**, 2009–2012.
- VanZadt, T., 1982: A universal spectrum of buoyancy waves in the atmosphere. *Geophys. Res. Lett.*, **9**, 575–578.