

Smaller Inner Ear Sensory Epithelia in *Neurog1* Null Mice Are Related to Earlier Hair Cell Cycle Exit

V. Matei,¹ S. Pauley,¹ S. Kaing,² D. Rowitch,² K.W. Beisel,¹ K. Morris,¹ F. Feng,¹ K. Jones,³ J. Lee,³ and B. Fritsch^{1*}

We investigated whether co-expression of *Neurog1* and *Atoh1* in common neurosensory precursors could explain the loss of hair cells in *Neurog1* null mice. Analysis of terminal mitosis, using BrdU, supports previous findings regarding timing of exit from cell cycle. Specifically, we show that cell cycle exit occurs in spiral sensory neurons in a base-to-apex progression followed by cell cycle exit of hair cells in the organ of Corti in an apex-to-base progression, with some overlap of cell cycle exit in the apex for both hair cells and spiral sensory neurons. Hair cells in *Neurog1* null mice show cell cycle exit in an apex-to-base progression about 1–2 days earlier. *Atoh1* is expressed in an apex-to-base progression rather than a base-to-apex progression as in wildtype littermates. We tested the possible expression of *Atoh1* in neurosensory precursors using two *Atoh1-Cre* lines. We show *Atoh1-Cre* mediated β -galactosidase expression in delaminating sensory neuron precursors as well as undifferentiated epithelial cells at E11 and E12.5. PCR analysis shows expression of *Atoh1* in the otocyst as early as E10.5, prior to any histology-based detection techniques. Combined, these data suggest that low levels of *Atoh1* exist much earlier in precursors of hair cells and sensory neurons, possibly including neurosensory precursors. Analysis of *Atoh1-Cre* expression in E18.5 embryos and P31 mice reveal β -galactosidase stain in all hair cells but also in vestibular and cochlear sensory neurons and some supporting cells. A similar expression of *Atoh1-LacZ* exists in postnatal and adult vestibular and cochlear sensory neurons, and *Atoh1* expression in vestibular sensory neurons is confirmed with RT-PCR. We propose that the absence of NEUROG1 protein leads to loss of sensory neuron formation through a phenotypic switch of cycling neurosensory precursors from sensory neuron to hair cell fate. *Neurog1* null mice show a truncation of clonal expansion of hair cell precursors through temporally altered terminal mitosis, thereby resulting in smaller sensory epithelia. *Developmental Dynamics* 234:633–650, 2005. © 2005 Wiley-Liss, Inc.

Key words: hair cells; sensory neurons; inner ear; neurotrophins; cell fate; cell cycle exit

Received 28 February 2005; Revised 24 June 2005; Accepted 8 July 2005

INTRODUCTION

Atoh1 and *Neurog1* belong to an ancient family of basic Helix-Loop-Helix (bHLH) genes that are involved in cell fate determination across phyla and

systems, typically being expressed in proliferating precursors. For example, in coelenterates, an ATONAL-like protein determines fate between muscle and neurons (Seipel et al., 2004). In mammals, various bHLH genes de-

termine the relative frequency of distinct retinal cell types (Akagi et al., 2004; Kim et al., 2005), regulate the size and progression of precursor fate in the olfactory epithelium (Wu et al., 2003), are involved in cell fate deter-

¹Department of Biomedical Science, Creighton University, Omaha, Nebraska

²Department of Molecular Biology, Harvard University, Boston, Massachusetts

³Department of Molecular, Cellular and Developmental Biology, University of Colorado, Boulder, Colorado

Grant sponsor: NIH; Grant numbers: RO1 DC005590, R01 DC04279, 1 C06 RR17417-01.

*Correspondence to: Bernd Fritsch, Ph.D., Dept. Biomed. Sciences, Creighton University, Omaha, NE 68178.

E-mail: Fritsch@Creighton.edu

DOI 10.1002/dvdy.20551

Published online 6 September 2005 in Wiley InterScience (www.interscience.wiley.com).

mination of neurons and switch between neuron and glial cell fates in the brain (Anderson et al., 1997; Bertrand et al., 2002), and determine the fate of progenitors of the gut epithelium (Yang et al., 2001). Except for continuously proliferating systems, these bHLH genes seem to be down-regulated in neonatal mammals within days after birth (Akazawa et al., 1995). As in these other neuronal and non-neuronal systems, at least three bHLH transcription factors are involved in the development of hair cells and sensory neurons of the ear: *Neurog1* is essential for sensory neuron development (Ma et al., 1998), *Atoh1* is essential for hair cell differentiation (Bermingham et al., 1999), and *Neurod1* is required for sensory neuron survival (Liu et al., 2000; Kim et al., 2001). In apparent agreement with other developing systems in which cell fate alterations have been reported after one or more of these genes have been mutated (Akagi et al., 2004), *Neurog1* null mutants show not only loss of all sensory neuron formation but in addition a severe reduction of hair cells in several sensory epithelia (Ma et al., 2000). As reported for seemingly comparable bHLH gene interactions in the eye, spinal cord, and the olfactory system (Gowan et al., 2001), it was hypothesized that this hair cell reduction indicates a molecular interaction of NEUROG1 with undisclosed proteins (Ma et al., 2000) in some common precursors of both cell types (Fritzsch et al., 2000; Fritzsch and Beisel, 2003).

In apparent contradiction to this idea are recent data from an elegant study, which suggest that *Atoh1* is a differentiation factor of postmitotic hair cells that plays no role in hair cell precursor selection in the cochlea (Chen et al., 2002). These conclusions were obtained using an *Atoh1* promoter fragment to drive eGFP (Gowan et al., 2001) as well as immunocytochemistry for ATOH1; both sets of data did not show *Atoh1* signal prior to cessation of proliferation of hair cell precursors. However, the transgenic construct used in this study requires the presence of *Atoh1* protein to be upregulated and may not show the earliest expression of *Atoh1* transcript. Low levels of protein may not be detected using immunocytochemis-

try. Indeed, previous and more recent work has shown somewhat earlier *Atoh1* expression using other techniques (Bermingham et al., 1999; Lanford et al., 2000; Zine et al., 2001; Woods et al., 2004). Other data showed an even earlier upregulation of *Atoh1*-eGFP in some vestibular hair cells at a time when both sensory neurons and hair cells become postmitotic in the vestibular system (Radde-Gallwitz et al., 2004). However, the onset of expression of *Atoh1* in the ear has not been investigated using more sensitive techniques such as RT-PCR or Q-PCR. It seems, therefore, fair to say that at the moment we do not know when *Atoh1* is first upregulated and what cell type is initially expressing *Atoh1*.

While *Atoh1* is clearly involved in hair cell differentiation (Izumikawa et al., 2005), it appears that sensory epithelia composed of undifferentiated cells that express *Atoh1-LacZ* and *Bdnf-LacZ* form even in *Atoh1* null mice and receive afferent and efferent innervation (Fritzsch et al., 2005a). In addition, recent data (Radde-Gallwitz et al., 2004) support previous results on expression of the transcription factor *Islet1* in differentiating vestibular sensory neurons and hair cells (Adam et al., 1998). These *Islet1* data agree with information gathered with the neurotrophin BDNF; *Bdnf* mRNA is also expressed in differentiating hair cells and sensory neuron precursors that appear to delaminate from the future sensory epithelia (Farinas et al., 2001; Fritzsch et al., 2002). Together these data are compatible with the idea that some sensory neurons and hair cells may be derived from common neurosensory precursors (Fritzsch and Beisel, 2004). Indeed, recent lineage tracing work in chicken has shown some lineage relationships of sensory neurons and hair cells (Sato and Fekete, 2005).

These data leave several questions unanswered: It remains unclear whether the detection of the earliest appearance of *Atoh1* expression is compromised by the sensitivity of the employed technique, leading to false-negative results. For example, it is possible that the delay in generating eGFP in the *Atoh1* transgenic mice may not show the earliest expression of ATOH1 protein (needed to drive the

promoter) and certainly will not show the earliest upregulation of *Atoh1* message. Likewise, limited expression of *Atoh1* mRNA may be below the detection threshold of in situ based techniques and thus may lead to false-negative results suggesting, for example, the absence of *Atoh1* expression in neurosensory precursors.

For this study we had six aims: First, we wanted to compare *Bdnf* expression in *Atoh1* and *Neurog1* simple and double null mice. Second, we wanted to determine terminal mitosis as revealed by BrdU in wildtype and *Neurog1* null mice. Third, we wanted to identify the earliest expression of *Atoh1* in control and *Neurog1* null mice using various techniques. Fourth, we wanted to study the expression of *Atoh1* as revealed by *Atoh-Cre* expression in Rosa 26 mice to provide evidence supporting the idea that some hair cells and sensory neurons are progeny of common neurosensory cells. Fifth, we wanted to show the expression of *Atoh1* in vestibular and spiral sensory neurons as revealed by *Atoh1-eGFP*, *Atoh1-LacZ* staining, and Q-PCR, suggesting that this labeling is not an integration artifact (Lumpkin et al., 2003). Sixth, we wanted to determine the significance of this expression in sensory neurons using a biological readout system and show whether absence of *Atoh1* may affect path-finding properties of sensory neurons. Lastly, we wanted to develop a model that can explain how *Neurog1* absence mediates hair cell reduction. For this model, we assume a possible NEUROG1-ATOH1 protein interaction in common precursors of hair cells and sensory neurons (neurosensory precursors) and that *Neurog1* absence skews cell fate decisions of common neurosensory precursors toward hair cells.

MATERIALS AND METHODS

Mouse Lines Used to Generate Single and Compound Mutant Mice

Two *Atoh1* null alleles were used in this study. One allele carried a Hprt cassette (*Atoh1-Hprt*) and the other one carried the *LacZ* reporter gene (*Atoh1-LacZ*) instead of the coding sequence (Ben-Arie et al., 1997, 2000).

To analyze the ATOH1/NEUROG1 interaction further, we generated doubly heterozygotic mice that carried the *Neurog1* null allele (Ma et al., 1998) in combination with the *Atoh1-Hprt* allele and crossed this line with a line carrying the *Bdnf* null allele combined with *LacZ* replacement (*Bdnf-LacZ*) previously described (Jones et al., 1994; Fritzscht et al., 2005a).

Mice with a *LacZ* or *Hprt* marker were bred from heterozygotes as previously described (Birmingham et al., 1999; Fritzscht et al., 2005a). Embryos were collected at embryonic day 11.5 (E11.5), E12.5, E16.5, and E18.5 and several heterozygotic animals were fixed at 2, 5, and 9 months of age. Older embryos and adults were perfusion or immersion fixed, after appropriate anesthesia, with 4% PFA for 30 min. Ears were rapidly dissected in PBS, and subsequently reacted for Lac-Z histochemistry overnight at room temperature (Farinas et al., 2001). Some of these ears of heterozygotic and homozygotic *Atoh1* mutants were also immunostained for nerve fiber stain using tubulin antibodies as previously described (Fritzscht et al., 1997) and anti β -galactosidase. All animal breeding and treatment was approved by an IACUC protocol no. 0630.

In older embryos, the neurotrophin *Bdnf* is expressed only in differentiated hair cells (Farinas et al., 2001) but may play a role in bHLH gene upregulation in earlier stages (Ito et al., 2003). We studied the expression of *Bdnf-LacZ* distribution in control, *Atoh1*, and *Neurog1* null littermates. To achieve this, we crossed *Atoh1-Hprt/Neurog1* doubly heterozygotes with *Bdnf* heterozygote animals. In the offspring, we selected an *Atoh1-Hprt/Neurog1/Bdnf-LacZ* compound heterozygote male and bred this animal with *Atoh1-hprt/Neurog1* heterozygote females. The resulting offspring produced *Atoh1-Hprt* null mutants with *Bdnf-LacZ* heterozygosity at the expected frequency of 1 in 8 embryos. Ears of E11.5, E12.5, and E18.5 perfusion fixed animals were reacted for β -galactosidase histochemistry as described above. Embryonic mice were from timed pregnancies and were, in addition, staged (Theiler, 1989).

In these animals, we also investigated the distribution of cells in which the terminal mitosis was visualized with BrdU injections (100 μ g/g). Injec-

tions were administered for each litter only once, at noon, to a given pregnant mouse of days E9.5–17.5 (with noon after vaginal plug considered as embryonic day 0.5). Animals were perfusion fixed at E18.5, genotyped and the ears processed for BrdU imaging. Briefly, ears were treated for 1 h with 1N HCl, washed, blocked with bovine serum, and incubated for 1 h with anti-BrdU antibody conjugated to Alexa 647 (Invitrogen). Ear epithelia were mounted flat on a slide in glycerol, coverslipped, and viewed with a confocal microscope (Biorad Radiance 2000 or Zeiss LSM). Assessment of labeled cells was achieved by first focusing on the surface of the organ of Corti using the 488-nm laser band. Subsequently, stacks of images were collected through the hair cells and supporting cells to include the nucleus of hair cells, which appeared as a black hole against the autofluorescence of the cytoplasm. In a second channel, the emission of the Alexa 647 was recorded, showing the distribution of BrdU in the nuclei. Nuclei were scored as labeled when 50% or more of the surface was stained in the Z-axis collapsed stacks. Identification of nuclei was supported by the fact that in most cases at least some minor specks of BrdU labeling appeared. Following this high-power analysis using a 40 \times 1.3 NA lens, we took overview images to illustrate the distribution of BrdU-labeled cells for illustration purposes only. In selected cases, we verified that the cells identified as hair cells were indeed hair cells by showing colocalization of BrdU in nuclei of cells labeled with *Bdnf*- β -galactosidase reaction product. To enhance imaging, the β -galactosidase reaction product was photoactivated to yield a 488-nm excitable fluorescing product (Matei et al., 2005).

Some wildtype and *Atoh1* null mutant embryos were also used for neuronal tract tracing as previously described (Fritzscht et al., 2005a) to study the possible effect of *Atoh1* absence on the pattern of innervation. In addition, some ears of *Atoh1* null and wildtype animals were reacted with acetylated tubulin to reveal the pattern of innervation. Ears were whole mounted in glycerol and viewed with the Zeiss LSM confocal system.

The bHLH gene *Neurod1* (formerly *NeuroD*) is expressed in all sensory neurons and in some differentiated hair cells in embryos older than E14.5 (Kim et al., 2001). We studied the *Neurod1-LacZ* expression in control and *Neurog1* null littermates. For this, we crossed *Neurog1* heterozygotes with *Neurod1-LacZ* heterozygotic mice. In the offspring, we selected *Neurog1/Neurod1-LacZ* compound heterozygote males and bred these animals with *Neurog1* heterozygote females. We bred two litters of E12.5–E18.5 embryos in which we had two offspring that were *Neurog1* null mutants and that were also heterozygous for *Neurod1-LacZ*. Ears of these E18.5 embryos were perfusion fixed and reacted for β -galactosidase histochemistry as described above.

After β -galactosidase reaction, ears were dissected and mounted flat for analysis of the sensory epithelia area. Some ears were embedded in soft epoxy resin, sectioned at 20 μ m thickness, and analyzed in a compound light-microscope (Nikon Eclipse 800). Images were captured using a CCD camera and processed using ImagePro or using the Zeiss LSM multiphoton confocal system and 2 photon activation (Matei et al., 2005). Images were compiled into plates using CorelDraw.

***Atoh1*-Cre Lines**

To generate this line, we used the JQ2-*Atoh1* promoter fragment originally described by (Gowan et al., 2001) and used to analyze *Atoh1* upregulation in developing hair cells (Chen et al., 2002) provided by Dr. J. Johnson. This fragment was ligated to the bacteriophage P1 Cre recombinase cDNA and injected into mouse embryo pronuclei to generate two transgenic mouse lines that carried the *Atoh1-Cre*. Integrity of the construct was checked by Southern blot. Transgenic animals were mated with Rosa 26 lines to demonstrate expression of Cre using the conditional LacZ reporter system of this line. We generated two transgenic strains of mice with this construct. Mice of one strain were fixed in 4% paraformaldehyde at E12.5 and E18.5, mice of the second strain were fixed at E11 and at 1 month of age, the latter after lethal anesthesia by perfusion. These stages were chosen to capture the earliest

expression detectable with these techniques as well as the long-term expression. Ears were dissected and either reacted as whole mounts for β -galactosidase or sectioned and reacted for β -galactosidase. Ears and sections were viewed in an Olympus EZ dissection scope or a Nikon Eclipse 800 and images were taken using a Coolsnap camera and Metamorph software. We also used a newly developed technique to photoactivate the BCI reaction product of X-Gal to enhance visibility (Matei et al., 2005).

RT-PCR was done on dissected inner ear tissues at E10.5 and E11.5 and Q-PCR was done on vestibular ganglia dissected from P0 and P10 ears using gene-specific primers according to a published procedure (Beisel et al., 2000) to determine the absence or presence of *Atoh1* transcripts. Care was taken that this tissue was not contaminated with adjacent brain tissue known to express *Atoh1*. The samples were treated with DNase to eliminate any trace genomic DNA. Afterwards, the quality of the RNA was assessed using the Agilent 2100 Bioanalyzer. Primer/probe sets were designed over introns to eliminate any possible amplification from contaminating genomic DNA. In the reactions themselves, no Template Control samples were used and none of these amplified a PCR product. First-strand synthesis was done using a T7-dT(24) primer (Beisel et al., 2000). A primary amplification was done using *mAtoh1-896 FOR* (5' CATCACCTTCGCACCGCCTC CTC 3') and a T7-specific oligomer at standard PCR conditions with 35 cycles. A secondary amplification step was done using the nested primer sets of *mAtoh1-1094 FOR* (5' CTAACA GCGATGATGGCACAGAA-G 3') and *mAtoh1-1777 REV* (5' AGTGATGAAGTGCGTGTATT CTGG 3') with 30 cycles. A primary amplification was done using *mAtoh1-896 FOR* (5' CATCACCTTCGCACCGCCTCCTC 3') and a T7-specific oligomer. A secondary amplification step was done using the nested primer sets of *mAtoh1-1094 FOR* (5' CTAACAGCGATGATGGCACAGAAG 3') and *mAtoh1-1777 REV* (5' AGTGATGAAGTGCGTGTATTCTGG 3'). We also dissected vestibular ganglia of newborn (P0; N = 3) and 7-day-old mice (P7; N = 2) and processed these ganglia for Q-PCR. We




Fig. 1.




Fig. 2.

used the primers outlined above and compared the *Atoh1* transcripts with *Neurod1* (a bHLH gene present in sensory neurons), *Neurog1* (a bHLH gene absent in postnatal sensory neurons), and ribosomal RNA.

RESULTS

Comparison of *Bdnf* Expression in *Atoh1* and *Neurog1* Simple and Double Null Mice

We previously showed that *Neurog1* null mice lose not only all sensory neurons but also show loss and reorganization of hair cells in all sensory epithelia (Ma et al., 2000). We also showed that undifferentiated cells that express *Atoh1-LacZ* develop in *Atoh1* null mice and attract afferent fibers likely because many of these cells co-express *Bdnf* (Fritzsche et al., 2005a), a known major attractor for growing ear sensory neuron processes (Tessarollo et al., 2004). Here we wanted to analyze how the temporal expression of *Bdnf* is altered in the absence of *Atoh1*, *Neurog1*, or both bHLH genes using *Bdnf-LacZ* as a marker for hair cells and sensory neu-

ron precursors, possibly including common neurosensory precursors.

The following differences and similarities in *Bdnf-LacZ* expression were apparent in the E12.5 and E18.5 ears of these mutants. First, the canal cristae and the apex of the cochlea showed roughly identical expression of *Bdnf*, regardless of age or genetic background and whether differentiated hair cells existed or not (Fig. 1A–K). Second, delaminating sensory neurons showed expression of *Bdnf* as they exited from the utricle, saccule, and ductus reuniens/basal turn of the cochlea of *Atoh1* null and wildtype embryos (Fig. 1E,F). As expected, no *Bdnf*-positive sensory neurons were visible outside the otocyst in *Neurog1* null mice at any time during development (Fig. 1C,G). Interestingly, delaminating cells were clearly positive for *Bdnf*- β -galactosidase in *Atoh1-Hprt* null mice despite the fact that no *Bdnf* was expressed in the areas they delaminated from (Fig. 1F). This *Bdnf* positivity suggests that sensory neuron precursors are capable of *Bdnf* up-regulation as they delaminate, but does not exclude the possibility that *Bdnf* is already present in wildtype

neurosensory precursors, giving rise to *Bdnf*-positive hair cells and sensory neurons. Lastly, there was a conspicuous lack of expression of *Bdnf* in the basal turn, utricle, and saccule of *Atoh1* null mutants (Fig. 1B,F). This absence of *Bdnf* expression in *Atoh1* null mutants contrasted sharply with the profound expression of *Bdnf* in distinctly labeled cells of the utricle in *Neurog1* null mice (Fig. 1F,G). We show *Bdnf-LacZ*-positive cells in the utricle of E12.5 *Neurog1* null mice (Fig. 1G) and *Bdnf-LacZ*-positive cells in the ductus reuniens of E18.5 *Neurog1* null mice (Fig. 1C). The frequency of these cells and their distribution was variable and we are currently investigating whether such cells are hair cells by crossing *Neurog1* with *Atoh1-LacZ*.

Since sensory neurons, but not hair cells of utricle and saccule, express *Bdnf* in *Atoh1* null mutants (Fig. 1B,F), we generated mice that were homozygotic null for both *Atoh1* and *Neurog1* combined with *Bdnf-LacZ* heterozygosity. These mice showed the combined phenotype of *Neurog1* null mice (shortened cochlea, reduction of saccular recess, absence of sen-

Fig. 1. Expression patterns of *Bdnf-LacZ* in E18.5 (A–D, H–K), E12.5 (E–G) ears of wildtype (A, E, K), *Atoh1* null mice (B, F, J), *Neurog1* null mice (C, G, I), and *Neurog1/Atoh1* double null mice (D, H) is shown. Note that the canal cristae (HC, AC) and the cochlear apex are strongly labeled in all cases. However, the basal turn shows no *Bdnf-LacZ* staining in *Atoh1* null mice (B, F) and *Neurog1/Atoh1* double null (D) mice. *Neurog1* null and *Neurog1/Atoh1* double null mice show decreased size of all six sensory epithelia (C, D) with the cochlea showing only one, tightly coiled turn. The saccule (S) is forming a small appendix on the smaller utricle in *Neurog1* null mice (C) and *Bdnf-LacZ* positive cells appear in the ductus reuniens (DR, vertical arrow) and the GER (horizontal arrow). In E12.5 embryos, *Bdnf-LacZ* positive sensory neurons delaminate in wildtype (E) and *Atoh1* null mice (F) from the utricle (U; arrow), the ductus reuniens (DR) leading from the saccule (S) to the cochlea (arrow), and the upper middle turn of the cochlea (arrow). No *Bdnf-LacZ*-positive sensory neurons and no ganglion exist in *Neurog1* null mutants (G; no Ggl). Note that delaminating cells may upregulate *Bdnf-LacZ* in *Atoh1* null mice de novo (F) since no *Bdnf-LacZ* is expressed in the utricle and saccule in *Atoh1* null mutants. In contrast to the absence of *Bdnf-LacZ*-positive cells in *Atoh1* null mice (F) is the more profound expression of *Bdnf-LacZ* in the utricle of *Neurog1* null mice (G), which have a saccule (S) formed of few cells (out of focus in G). These data suggest a differential effect of *Atoh1* and *Neurog1* on *Bdnf-LacZ* expression in hair cells, in particular in the utricle, saccule, and basal turn in older embryos. Direct comparison of the anterior crista in various genetic backgrounds shows the additive effect of *Neurog1* and *Atoh1* null mutation on the size and degree of differentiation of this sensory epithelium (H–K). AC, anterior crista; DR, ductus reuniens; Ggl, ganglion; HC, horizontal crista; PC, posterior crista; S, saccule; U, utricle. Scale bar = 100 μ m.

Fig. 2. The terminal mitosis of sensory neurons (Spgl) and hair cells (HC) is shown in wild type mice as revealed with BrdU injection followed by delayed recovery at E18.5 (B–E). Differentiated hair cells can be identified at this stage using either sections of *Bdnf-LacZ* heterozygotic animals (A) or by imaging flat mounted organ of Corti (OC) using a confocal microscope to generate an image stack (stack in A) that extends from the surface to the nuclei of hair cells (A,B). Such a stack of images, if collapsed into a single image (B), will show the nuclei as black holes (B). Fluorescent antibody staining of BrdU will reveal labeling (shown in red in B–E) that allows us to assess at higher magnifications the degree of labeling as over 50% (white arrows in B) or less than 50% (black arrow in B). As the organ of Corti is regularly organized into one row of inner hair cells (IHC in A,B) and three rows of outer hair cells (OHC in A,B) with nuclei much larger than those in cells of the adjacent greater and lesser epithelial ridge (GER, LER), hair cell identification is reliable with this approach. Low-power images (C,D) can be used to assess the overall distribution of labeled cells and thus allow imaging the spatial changes over time following single injections of BrdU at different times in different litters. Injections of BrdU at E11.5 (C) show prominently labeled sensory neuron nuclei in the upper middle turn that were apparently undergoing DNA synthesis for a terminal mitosis at this time as indicated by the almost complete filling of the nucleus of these cell nuclei with anti-BrdU antibody. In contrast, hair cells (arrows) show only puncta of BrdU staining in the nuclei except for the apex (C), suggesting that most hair cell precursors went through two or more rounds of division before their terminal mitosis. An E12.5 injection of BrdU labels hair cell nuclei (arrow) in the basal turn and scattered spiral ganglion cells in the apical half (B,D). Combining BrdU labeling with *Bdnf-LacZ* staining shows that indeed some hair cells, positive for the green fluorescent product (E) have red, BrdU-labeled nuclei (arrow in E). Many supporting cell nuclei throughout the cochlea are also positive for BrdU. GER, greater epithelial ridge; IHC, inner hair cell; LER, lesser epithelial ridge; OC, organ of Corti; OHC, outer hair cell; Spgl, spiral ganglion. Scale bar = 10 μ m (A,B), 100 μ m (C–E).

TABLE 1. Mutation of *Atoh1*, *Neurog1*, and double *Atoh1/Neurog1* Affects the Size and Length of Cochlea and Anterior Crista as Judged by Measuring the Area of *Bdnf* Expression and/or Total Length

	WT	<i>Atoh1</i> Null	<i>Atoh1</i> /WT (% of length)	<i>Neurog1</i> null	<i>Neurog1</i> /WT (% of length)	<i>Atoh1- Neurog1</i> double null	<i>Atoh1-Neurog1</i> T° ler (% of length)
Cochlea							
<i>Bdnf</i> (μm)	5,495 ± 68	3,722 ± 37	68	2,356 ± 29	42	1,329 ± 18	24%
Total (μm)	5,776 ± 86	5,407 ± 78	93	3,083 ± 48	53	3,003 ± 76	52%
Anterior crista							
Width (μm)	237 ± 21	125 ± 19	53	193 ± 44	81	79 ± 32	33%
Length (μm)	608 ± 35	451 ± 38	74	413 ± 35	68	225 ± 86	37%
Area (μm ²)	107,350 ± 193	36,487 ± 283	34	58,003 ± 175	54	12,755 ± 88	12%
	N = 6	N = 6		N = 4		N = 4	

sory neurons; Fig. 1D; Table 1) and *Atoh1* (absence of *Bdnf* expression in utricle, saccule, and basal turn of cochlea; Fig. 1D).

The data on more prominent expression of *Bdnf* in cells of the utricle of *Neurog1* null mice at E12.5 are compatible with at least three scenarios: (1) hair cell precursors exit the cell cycle in younger animals and upregulate *Bdnf*; (2) neurosensory precursors convert to a hair cell fate and remain positive for *Bdnf*; (3) sensory neurons fail to delaminate, remain as *Bdnf*-positive cells, that eventually die and are replaced by hair cells forming later. In each case, more cells should be positive for *Bdnf* in the utricle of *Bdnf* null mice at an earlier stage, precisely what we demonstrated (Fig. 1G). The assumption of earlier exit from the cell cycle of hair cells can be verified by demonstrating alterations in the time of terminal mitosis of hair cells in the *Neurog1* and *Atoh1* null mice, our next experiment. Such data can also show whether cells that exit the cycle at this early stage remain or die. If such cells would die, BrdU-positive hair cells should not be found at later stages after an early embryonic injection of BrdU. Such hair cells would have replaced the dead cells by cells without BrdU labeling, thus allowing us to distinguish between the three possible scenarios.

Terminal Mitosis as Revealed by *BrdU* in Wildtype and *Neurog1* Null Mice

We injected BrdU once in pregnant mice ranging between E9.5 and E17.5

and analyzed the distribution of cells that showed BrdU-Alexa 647 antibody labeling of at least 50% of the area of a given nucleus at E18.5. We used the simple fact that differentiated hair cells form at this stage one row of inner and three rows of outer hair cells to identify hair cells. As is obvious from sections of the cochlea using our hair cell marker *Bdnf-LacZ* (Fig. 2A), hair cell nuclei are distinctly distributed from the supporting cell nuclei. Thus careful focusing of a flat mounted cochlea allows clear identification of hair cells and their nuclei using autofluorescence of the cytoplasm in combination with nuclear size and distribution to distinguish between the large hair cell nuclei and the smaller nuclei of cells of the greater epithelial ridge (GER; Fig. 2A,B) and lesser epithelial ridge (LER; Fig. 2A,B). Such a criterion provided information on cells that had their terminal mitosis shortly after the time of injection as clearance of BrdU takes around 2 h. Analysis in other systems has shown that the cell cycle length changes over the time of our analysis from about 8 h around E9.5 to about 12 h around E16.5 (Calegari and Huttner, 2003). Any continued cell cycling, therefore, would have reduced the BrdU content of the nuclei to less than 50% (Fig. 2B) within 8–12 h after the injection. Such reduction of BrdU results in specks of BrdU-Alexa 647 labeling that amounted to less than 50% of a given nucleus (Fig. 2B). This technique provides a clear picture of the distribution of cells that underwent terminal mitosis shortly after BrdU injection and remained

mitotically quiescent afterwards. This technique will also reveal decreasingly labeled cells derived from precursors labeled at the time of BrdU application but that continue to cycle. Such decrease in labeling will allow detailing further the topological progression of cells that undergo terminal mitosis at earlier or later stages. Of course, cell death would eliminate such cells and their BrdU labeling. The whole mount approach used here will allow identification of cells using histological criteria established over 100 years ago (Ruben, 1967). In our hands, some of the commonly used antibodies to highlight hair cells showed artifacts or no staining above background, probably due to the acid treatment necessary to uncover the BrdU epitope for the antibody recognition of BrdU. In addition, we also used in selected cases *Bdnf-LacZ* heterozygotic animals, photoactivated the β-galactosidase reaction product that is only in hair cells in the basal turn, and showed directly that the cells we identified as hair cells based on topology and cytology criteria are indeed hair cells (Fig. 2A,E).

Using these criteria in whole-mounted ears, we detected labeled spiral sensory neurons in a base-to-apical progression and hair cells in an apex-to-base progression as previously described (Ruben, 1967). Specifically, injections of BrdU at E9.5 and E10.5 and imaging at E18.5 showed label in only a few neurons in the basal turn with only punctate labeling in the apex (data not shown). Injections at E11.5 and imaging the distribution of nuclear staining at E18.5 showed that neurons were labeled in

the basal and middle spiral turn (Fig. 2C). In contrast, neurons in the apex had less than 50% filled nuclei, suggesting that they had at least one or more additional mitosis after the BrdU application. Using the same criterion, we found only a few labeled hair cells in the apical-most tip of the organ of Corti. However, a trail of less labeled hair cell nuclei extended throughout the middle turn almost to the basal hook (Fig. 2C).

Injections of BrdU at E12.5 labeled only neurons in the apex of the cochlea and many hair cells in the apical half turn of the organ of Corti (Fig. 2B,D). As in earlier stages, there was a sharp boundary toward the apex of labeled hair cells whereas the boundary toward the basal turn was fuzzy with many cells showing spotty BrdU labeling. Injections at E13.5 labeled very few spiral sensory neurons and showed labeled hair cells extending throughout the middle and basal turn. These injections also showed profound but spotty labeling of other cell types throughout the cochlea, making it difficult to identify the labeled hair cells at low magnification. However, using the above-described procedure of generating a single focal stack through the hair cells alone, we were able to identify a few hair cells near the basal tip that showed profound nuclear labeling. Injections at E14.5 labeled no sensory neurons and only a few hair cells in the most basal tip of the organ of Corti where the rows of outer hair cells become reduced (data not shown).

These data suggest that in the apex, the first hair cells become postmitotic around the time the last spiral sensory neurons become postmitotic, with an apparent spatial and temporal overlap. In contrast, in the base, hair cells become postmitotic up to three days after spiral sensory neurons. These data suggest that, overall, sensory neurons and hair cells are among the first cells to exit cell cycling in opposing gradients as previously demonstrated (Ruben, 1967).

***Neurog1* Null Mice**

We next investigated the effects of *Neurog1* on time of hair cell terminal mitosis in *Neurog1* null littermates of our wildtype analyses presented above. In


animals injected at E9.5 to E11.5 and analyzed at E18.5, we found no labeling in sensory neurons, consistent with previous data that suggest complete abrogation of any sensory neuron formation in *Neurog1* null mice (Ma et al., 1998, 2000). However, in the middle to apical turn of the shortened and widened organ of Corti of these mutants, we found numerous hair cells in which the nuclei were at least 50% filled with the BrdU antibody fluorescence (Fig. 3B,C). Injections after E13.5 did not reveal any hair cells with labeled nuclei, suggesting that the entire phase of hair cell proliferation was shifted in *Neurog1* null mutants to an approximately 1–2 days earlier start of hair cell terminal mitosis compared to wildtype littermates (Fig. 3A). Data in the vestibular endorgans were less obvious, presumably because of the prolonged time over which hair cells are generated in these epithelia (Ruben, 1967) and the lack of a simple linear progression of terminal mitosis along one axis, as in the cochlea. In summary, the data on the cochlea suggest that abrogation of *Neurog1* leads to a terminal mitosis of hair cells in the cochlea that temporally overlaps with the time of terminal mitosis of sensory neurons in wildtype littermates.

We next wanted to confirm and extend this finding with other markers that show a delayed expression in hair cells compared to sensory neurons. *Neurod1* is upregulated in sensory neurons immediately after *Neurog1* (Ma et al., 1998) but appears only late in development in some hair cells, progressing from base to apex (Kim et al., 2001). We, therefore, bred mice that were null for *Neurog1* and heterozygous for *Neurod1*. Wild type E18.5 littermates showed no *Neurod1*-positive hair cells in the apex, and had only few inner hair cells displaying *Neurod1*-mediated β -galactosidase staining (Fig. 3D,F). In contrast, *Neurog1* null mice showed numerous *Neurod1*- β -galactosidase-positive hair cells in the apex that were scattered across the multiple rows of hair cells of these mutants (Fig. 3E,G). As previously shown (Fig. 1A,C), the absence of *Neurod1*-positive sensory neurons (Fig. 3D) results in the *Neurog1* null mice in a much tighter coiling of the cochlea. At the same magnification, we can therefore show the entire apex of the *Neurog1* null mice but not of the

wildtype littermate (Fig. 3D,E). These data show upregulation of *Neurod1* in many more hair cells throughout the organ of Corti of E18.5 *Neurog1* null mice (Fig. 3F,G). Such data are consistent with the idea that at least some of the hair cells that undergo 1–2-day earlier terminal mitosis are derived from neurosensory precursors that cannot differentiate into neurons in the absence of *Neurog1* but become hair cells instead. Other genes with similar expression profiles are needed to verify whether this is peculiar to *Neurod1* or is a general effect of NEUROG1 protein abrogation.

In conclusion, these data confirm previous findings that absence of *Neurog1* affects not only sensory neuron formation but also hair cell formation (Ma et al., 2000). We now show that absence of *Neurog1* leads to a terminal mitosis of hair cells up to two days earlier and results in upregulation of *Neurod1* in many more hair cells of E18.5 embryos throughout the cochlea. These data are consistent with the notion of a *Neurog1/Atoh1* interaction in the common neurosensory (hair cell/sensory neuron) precursors to regulate the terminal mitosis and cell fate of cell types derived from these proliferating precursors. A simple way to explain these effects of *Neurog1* null mutation would be that ATOH1 protein-mediated onset of differentiation interferes with the cell cycle progression thus resulting in premature terminal mitosis of hair cells and truncation of the proliferative capacity of those sensory neuron/hair cell precursors. More complex interactions would also be compatible with these data such as paracrine interactions between neuronal and sensory precursors. We, therefore, next wanted to see whether *Atoh1* is expressed early enough to be in mitotically active neurosensory precursors and whether this expression changes in the *Neurog1* null mice, thus supporting the idea that some sensory neurons of the mammalian ear show coexpression of *Neurog1* and *Atoh1*, which interact either directly or through yet-to-be-determined intermediaries to allow full clonal expansion of their common and/or separate precursors.

Fig. 3. We show a comparison of temporal patterns of terminal mitosis in the cochlea of the wild type and *Neurog1* null littermates imaged at E18.5 (A–C), and of *Neurod1-LacZ* expression at E18.5 in a *Neurod1-LacZ* heterozygote wild type and *Neurog1* null littermate (D–G). In the absence of *Neurog1*, the hair cells exhibit alteration of the temporal pattern of terminal mitosis. BrdU injection at E9.5 and examination of the cochlea at E18.5 reveals some fully labeled nuclei (i.e., terminal mitotic cells) in the apex of the *Neurog1* mutant cochlea (arrows in C; green is autofluorescence of hair cells). BrdU injected at E11.5 labels cells in the middle turn of the organ of Corti (OC) with only specks of BrdU labeling in wildtype (A) and shows prominent nuclear labeling of spiral ganglion neurons (Spgl in A). In *Neurog1* null littermates, there are no spiral ganglion cells (B). Instead, a prominent labeling of hair cell nuclei of the organ of Corti indicates a premature terminal mitosis of hair cells in the middle turn in a pattern similar to the proliferation of spiral ganglion neurons in wildtype (compare A,B). Similar effects on altered expression patterns are found in *Neurod1* expression in *Neurog1* null mutants and wildtype littermates (D–G). At E18.5, spiral neurons are strongly labeled for *Neurod1* in the wild-type ear (D). In contrast to the very faint labeling of the wild-type organ of Corti, the *Neurod1* null ear (E) shows profound *Neurod1-LacZ* staining in many hair cells. Note the absence of spiral ganglion neurons in *Neurog1* null mice (compare D and E) and the multiple rows of outer hair cells (OHC) in the apex of the mutant cochlea (compare F and G). IHC, inner hair cell; OC, organ of Corti; Spgl, spiral ganglion. Scale bar = 100 μ m (A,B, D,E), 10 μ m (C, F, G).



Earliest Expression of *Atoh1* in Control and *Neurog1* Null Mice as Revealed by Various Techniques

We investigated whether the temporally shifted terminal mitosis of hair cells of *Neurog1* null mice also lead to an upregulation of *Atoh1*. This would indicate not only that the onset of terminal mitosis of hair cells is accelerated but also that ATOH1-mediated hair cell differentiation in *Neurog1* null mice is accelerated compared to wildtype mice. We studied the expression of *Atoh1* in the ear of E11- to E13.5-old mice. In the wildtype littermates, the E12.5 in situ data showed a very faint staining near the forming vestibular epithelia. No differences other than a reduction in size of vestibular epithelia were observed in the *Neurog1* null littermate and wildtype control animals. However, a difference was observed in the cochlea of E13 *Neurog1* null mice compared to age-matched littermates. In the *Neurog1* null mice, we obtained an *Atoh1* signal




Fig. 4.

in the cochlear duct that was strongest near the upper middle turn, fell off steeply near the apex, and tapered off in the lower basal turn (Fig. 4C). In contrast, the wildtype littermates showed a fainter in situ signal exclusively in the basal turn (Fig. 4B). Our β -galactosidase stain in *Atoh1-LacZ* heterozygotic animals showed a comparable temporal and spatial onset of *Atoh1* expression to *Atoh1* detected by in situ hybridization (Fig. 4E). We also obtained some *Atoh1*-eGFP expressing embryos (courtesy of Dr. P. Chen and N. Segil) and could not detect any eGFP signal in the cochlea of mice that were at least one day older than the mice of our lines (Fig. 4A), thus confirming their reported observations (Chen et al., 2002).

We next checked whether absence of eGFP signal in these lines is due to a different timetable of *Atoh1* upregulation or due to delayed eGFP upregulation owing to the construct. Our *Atoh1-LacZ*-positive ears showed a weak signal in the basal turn of the cochlea (Fig. 4E) in an animal in which we could not detect any eGFP, suggesting that this technique fails to detect the *Atoh1* upregulation visible with the other two techniques employed here. Most importantly, these data suggest that in our mouse lines there is a potential spatio-temporal overlap of detectable *Atoh1* expression in proliferating precursors only in the base of the cochlea, but not in the mid-

dle turn and apex, supporting previous suggestions (Chen et al., 2002).

We also tested whether we could detect *Atoh1* signal using PCR technique. All three samples of E11.5 embryonic ears exhibited *Atoh1* expression (data not shown), while 1 out of two samples from E10.5 embryos were positive for *Atoh1* (Fig. 5A). The RNase treated controls did not have any amplification products. These data suggest that the appearance of *Atoh1* transcripts in the developing inner ear is at or around E10.5. This is the earliest time *Atoh1* is expressed in detectable amounts in the developing ear.

In summary, these data imply that *Neurog1* expression affects spatial upregulation of *Atoh1* in apical hair cells that exit the cell cycle about two days earlier. In the absence of *Neurog1*, the hair cells that show this earlier terminal mitosis also show spatial upregulation of *Atoh1*. However, the gradient does not follow the previously reported pattern of upregulation of *Atoh1* in wildtype mice and progresses instead from near the apex toward the base. Interestingly, *Atoh1* expression shows little temporal alteration (only about $\frac{1}{2}$ day in the apex) compared to terminal mitosis (two days), suggesting that those two events are not tightly coupled. We have recently identified several other bHLH genes in the ear using PCR and their expression profiles could show a more pro-

found upregulation in *Neurog1* null mice, something we are actively investigating. Compared to PCR data, our histological techniques do reveal expression of *Atoh1* at least 24 h later (E10.5 vs. E11.5), suggesting that such techniques are less than appropriate to exclude expression of low levels of *Atoh1* in common neurosensory precursors. We, therefore, attempted to show such expression using yet another, more sensitive technique, *Atoh1-Cre* expression in Rosa 26 mice.

Expression of *Atoh1* as Revealed by *Atoh-Cre* Expression in Rosa 26 Mice: Evidence That Some Hair Cells and Sensory Neurons May Be the Progeny of Common Neurosensory Cells

Cre-mediated expression of the *LacZ* reporter results in the Rosa 26 line in β -galactosidase staining wherever the CRE-recombinase was and is active (Ohyama and Groves, 2004). Recently, an *Atoh1* promoter fragment used to drive eGFP (Gowan et al., 2001) was engineered by us to drive *Cre*. If *Atoh1* is expressed in neurosensory precursors, the *Atoh1-Cre* should show β -galactosidase staining in sensory neurons as well as hair cells. We investigated the ears of E 11, E12.5, and E18.5 and P7 of two *Atoh1-Cre/Rosa 26* mice. In E12.5 embryos of one *Atoh1-Cre* line, we found expression of β -galactosidase only in single cells in what appeared to be several vestibular sensory epithelia patches of the utricle, saccule, anterior, horizontal, and posterior crista. In the cochlea, staining was in the medial wall of the basal and middle turn (Fig. 4D). In addition to these intraepithelial cells, β -galactosidase-positive cells were found outside the inner ear epithelia in the vestibular ganglion and the forming spiral ganglion (Fig. 4D). In the hindbrain and midbrain, *Atoh1-Cre*-mediated β -galactosidase stain displayed a pattern consistent with the expression of *Atoh1* as revealed in the same aged *Atoh1-LacZ* brains (data not shown). This expression was verified in a total of four ears and two brains of E12.5 mice.

The second *Atoh1-Cre* line showed a more profound β -galactosidase stain-

Fig. 4. The earliest expression of *Atoh1* as revealed by different techniques (A,B,D,E) in wildtype and *Neurog1* null mice (C) is compared. At E13.5, *Atoh1*-eGFP shows a profound signal in the vestibular sensory epithelia and in some delaminating cells (arrows in A). However, there is no detectable *Atoh1*-eGFP signal in the cochlea. In contrast, in situ hybridization (B) and *Atoh1-LacZ* (E) show *Atoh1* expression in the basal turn of E13 and E13.5 animals (B, F). Comparison of *Atoh1-LacZ* (F) and *Atoh1-Cre/Rosa26* (D) shows that more cells are *Atoh1*-positive in the vestibular sensory epithelia of *Atoh1-LacZ* animals at an earlier stage (F). In contrast, *Atoh1-Cre* shows additional expression in delaminating spiral ganglion neurons (Spgl) near the base and vestibular ganglion neurons (Vgl) near the utricle and saccule (D). Of all the techniques employed here, only *Atoh1*-eGFP shows some of these delaminating cells as well (compare A, D). *Atoh1* in situ hybridization in *Neurog1* null mutants shows a more apical expression in the E13 cochlea (C). This expression suggests a differential spatial upregulation in the upper middle turn of the cochlea instead of the basal turn as in wildtype littermates (B, C). Note the reduced size of vestibular sensory epithelia (compare B and C) in *Neurog1* mutants. The earliest expression of all tissue-based technique was the few β -galactosidase-positive cells in the otocyst wall (indicated by the white dotted line) and in the vestibular ganglion of E11-day-old *Atoh1-Cre* embryos (G-I), using photoactivation of the β -galactosidase reaction product (green labeled cells). Those cells were immediately adjacent to the ingrowing fibers targeted to the utricle, anterior and horizontal crista (H) labeled with tubulin antibody (red). Note that a few cells positive for *Atoh1-Cre* were found outside the otocyst wall in the forming vestibular ganglion (I). Combining these data suggests that the sensitivity of the different techniques varies substantially and several need to be compared to evaluate early onset of *Atoh1* expression. AC, anterior crista; HC, horizontal crista; PC, posterior crista; S, saccule, Spgl, spiral ganglion; U, utricle; Vgl, vestibular ganglion. Scale bar = 100 μ m.




Fig. 5. *Atoh1* expression as revealed in the embryonic ear with PCR (35 cycles; **A**) and in the vestibular ganglion of postnatal animals with Q-PCR (**B**). *Atoh1* presence was evaluated using PCR in two ears of E10.5 mice. Note that only one ear is positive for the PCR product. All three ears of E11.5-old-mice were *Atoh1*-positive (data not shown), suggesting that *Atoh1* expression in the ear starts around E10.5. Q-PCR of vestibular ganglia shows expression of *Atoh1* after 37 cycles at P0 and P7 using both 3' and 5' primers. No *Neurog1* was detectable in these cells, but *Neurod1*, known to be expressed in these cells, was detectable at about the same level as *Atoh1*. These data show the presence of *Atoh1* mRNA in the vestibular ganglion in postnatal animals.




Fig. 6.




Fig. 7.

ing and we chose this line to evaluate the earliest *Atoh1*-expression, enhancing the β -galactosidase reaction product with a recently developed photoactivation process (Matei et al., 2005)

and imaging with the confocal microscope. These data showed that as early as E11, cells are *Atoh1-Cre*-positive and appear in distinct areas of the ear that appear to correspond to

future sensory epithelia (Fig. 4G–I). In addition to cells in the otocyst wall, we also found a few cells distributed among the differentiating vestibular sensory neurons (Fig. 4I). Labeling of neurons with acetylated tubulin revealed that developing nerve fibers were near the *Atoh1-Cre*-positive cells. These data suggest that this technique allows detection of *Atoh1* in tissue as early as approximately 12 h after we can first detect *Atoh1* using RT-PCR.

Some positive cells were also found outside the sensory epithelia in *Atoh1-eGFP* ears (Fig. 4A) suggesting that the expression of *Atoh1* is possibly transient and thus cannot be detected with in situ hybridization or *Atoh1-LacZ*. Whether the *Atoh1* fragment-mediated signal is due to an expression in common neurosensory precursors or reflects an independent and transient upregulation in delaminating sensory neurons in addition to hair cell precursors remains unclear as we could not detect it in proliferating cells.

At E18.5, β -galactosidase was found in almost all hair cells of all sensory epithelia, with the exception of the most apical tip of the cochlea (Fig. 6A,B). In addition, *Atoh1-Cre*-mediated *LacZ* expression was found in spiral ganglion neurons, vestibular ganglion neurons, pillar cells, and other supporting cells (Fig. 6B). We also investigated this expression of *Atoh1-LacZ* in hair cells of E18.5 *Neurog1* null mice combined with *Atoh1-LacZ* heterozygosity and obtained comparable results. Combined, these data suggest that the most apical tip

of the cochlea, the area in which the first hair cells become postmitotic, has a profound delay of *Atoh1* upregulation of several days, in particular in wildtype animals, but also in *Neurog1* null mice. Cells other than hair cells were also *Atoh1-Cre*- β -galactosidase positive. These cells were particularly frequent in the forming spiral prominence (Fig. 6A,B). In addition, few supporting cells of the cochlea and the vestibular epithelia were also β -galactosidase-positive. Interestingly, some cells of the non-sensory epithelium of the ear were also β -galactosidase-positive. These cells were particularly prominent in the extension of the sacculus into the endolymphatic duct (Fig. 6A,B) but also as an extension of the cruciate eminence of the canal cristae into the canals, in particular the anterior and posterior canal.

The ears of the 1-month-old second *Atoh1-Cre-Rosa 26* line (P31) also showed expression other than the hair cells of the sensory epithelia (Fig. 7B,C,G). This expression showed three prominent distribution patterns, sensory neurons, supporting cells in the sensory epithelia (in particular pillar cells; Fig. 7G) and non-sensory cells of the ear outside the sensory epithelia such as a few cells in the stria vascularis (Fig. 7B). Almost all sensory neurons were positive in both the spiral ganglion and the vestibular ganglion in this line with little staining elsewhere (Fig. 7B,C) with labeling being so prominent that the nerve fibers showed a faint β -galactosidase reaction product (Fig. 7C).

These data obtained from two

Atoh1-Cre transgenic lines suggest that *Atoh1-Cre*-expressing precursors may not only give rise to hair cells but also to sensory neurons. Since positive cells are found within 12 h of the first possible detection of *Atoh1* transcript inside and outside the otocyst, we suggest that these cells are progeny of *Atoh1*-positive precursors. Other cells found positive in both lines are some supporting cells and non-sensory epithelia of the ear. We will next focus on the possibility that *Atoh1* is indeed expressed in sensory neurons where our two *Atoh1-Cre* lines showed strong expression (Figs. 6A,B, 7B,C).

Expression of *Atoh1* in Vestibular and Spiral Sensory Neurons as Revealed by *Atoh1-eGFP*, *Atoh1-LacZ* Staining and Q-PCR Suggested That This Labeling Is Not an Integration Artifact

Atoh1-Cre mediated β -galactosidase could provide a false-positive signal that is related to the genomic integration of the promoter fragment. To test for this possibility, we investigated the ears of E12.5 and P7 *Atoh1-eGFP* mice (courtesy of Drs. Chen and Segil). Initially, we could not detect any staining outside the sensory epithelia. However, using the Zeiss LSM confocal system color unmixing algorithm, we were able to extract a weak eGFP signal from the background in scattered spiral sensory neurons of P7 (Fig. 7A) mice. We also found few

Fig. 6. Expression of *Atoh-Cre* as revealed with a *Rosa 26* line is shown for E18.5 embryo ears (A–D). Hair cells are most profoundly positive for the β -galactosidase reaction product (A–C) and can be seen in whole ears (A), flat mounted cochlea (B), or sections imaged with confocal microscope (C, level of section is indicated by dotted line in B). However, additional cells in the cochlea are also positive (A,B). Note that *Atoh1-Cre* staining can be shown in some spiral ganglion neurons (Spgl in A,B; SpglN in D) at any level of magnification (A,B,D). Note that the X-Gal reaction product deposited at places of β -galactosidase activity tends to form small deposits outside of cells (small blue specks in C) but is most prominently deposited in the cytoplasm of *Atoh1-Cre*-positive cells like inner and outer hair cells and spiral ganglion neurons (C,D). Dotted line in B indicates the section plane with C,D indicated as squares. IHC, inner hair cell; OC, organ of Corti; OHC, outer hair cell; S, saccule; Spgl, spiral ganglion; SpglN, spiral ganglion neuron; U, utricle. Scale bar = 100 μ m (A,B), 20 μ m (C,D).

Fig. 7. Expression of *Atoh1* (A,I) in hair cells and sensory neurons of juveniles and adults is shown as revealed with different techniques. With all these techniques, hair cells are most profoundly positive (C, D, F, G, H). However, additional cells in the cochlea are also positive for *Atoh1* using either *Atoh1-eGFP* (A, D), *Atoh1-Cre* (B, C, G), or *Atoh1-LacZ* (H) and can be identified as pillar cells. Additional label is found in cells of the stria vascularis (near B). Note that *Atoh1-LacZ* staining in spiral or vestibular sensory neurons can be readily demonstrated in postnatal animals of all ages with any of these techniques (A, B, C, E, I). High-power images show that the vestibular ganglion neurons (I) are approximately 50 μ m in diameter and thus cannot be confused with any other cell type in the vestibular ganglion. The similarities in detail of the sensory neuron expression obtained with all three techniques support the idea that at least some vestibular and spiral neurons express *Atoh1*. However, the expression of β -galactosidase is particularly profound in spiral and vestibular neurons of *Atoh1-Cre* mice, suggesting that there may be a build up of β -galactosidase over time, leading to a more obvious reaction in these large cells including a faint labeling in the nerve leading to the sensory epithelia. IVG, inferior vestibular ganglion; OC, organ of Corti; S, saccule; Spgl, spiral ganglion; SVG, superior vestibular ganglion; U, utricle. Scale bar = 50 μ m (D,G–I) and 100 μ m (A–C,E,F).

eGFP-positive cells delaminating in E12.5 *Atoh1*-eGFP mice (Fig. 4A). Detailed comparison of *Atoh1-Cre* and *Atoh1*-eGFP thus confirms staining in hair cells and some sensory neurons, but also shows some differences in these three transgenic lines using the same *Atoh1* promoter fragment in additional staining of cells in the cochlea. Most importantly, the signal in delaminating cells of E12.5 *Atoh1*-eGFP mice was stronger than in sensory neurons at P7 (Figs. 4A, 7A), suggesting that *Atoh1* might be transiently upregulated. Moreover, only one of our two *Atoh1-Cre* transgenic lines showed labeling in a large fraction of sensory neurons, suggesting that the possibly transient expression of *Atoh1* in these cells might be more widespread. While it is virtually inconceivable that this expression of *Atoh1* in sensory neurons presents an integration artifact that would need to be similar in all three transgenic lines, we nevertheless wanted to obtain independent confirmation of the validity of the expression in sensory neurons indicated by the *Atoh1* promoter fragment to demonstrate that this expression in sensory neurons is truly not an aberration of *Atoh1* promoter integration.

We reasoned that if *Atoh1* staining is indeed in sensory neurons and not an artifact of the *Atoh1* promoter fragment, then we should be able to detect *Atoh1-LacZ* expression in the spiral and vestibular sensory neurons of juvenile and adult mice and should also detect it with Q-PCR. We investigated this possibility by staining the ears of 9-day-, 2-month-, 5-month-, and 9-month-old *Atoh1-LacZ* heterozygotic mice for β -galactosidase. Our data demonstrated that indeed there is an expression of β -galactosidase not only in hair cells but also in sensory neurons. Moreover, the overall distribution was not unlike that found in E18.5 *Atoh1-Cre* mice and showed only a small fraction of cells to be positive for *Atoh1-LacZ* (Fig. 7E,I). The distribution of these cells in the cochlea showed a more profound cellular staining in the apex, the area in which *Neurog1* null mutants show the most profound effect on the spatial expression of *Atoh1* and on temporal changes of hair cell terminal mitoses. Vestibular ganglion neurons are among the largest cells in the ear, 40–50 μ m in

size, and cannot be mistaken for any other cell type in the vestibular ganglia (Fig. 7I). Interestingly, all transgenic lines as well as adult *Atoh1-LacZ* lines showed prominent expression in virtually all hair cells and most pillar cells (Figs. 6B, 7D,G,H) implying that these cells should also be considered as *Atoh1*-positive cells.

We next tested whether spiral sensory neurons express *Atoh1* and if so how this expression changes quantitatively over time. Previous work had shown that *Atoh1* is sharply downregulated in the brain (Akazawa et al., 1995) and it could, therefore, be possible that the *Atoh1-LacZ* expression is unrelated to actual *Atoh1* protein and gene expression. We, therefore, investigated the expression of *Atoh1* using in situ hybridization and Q-PCR at P0 and P10. With both techniques, we confirmed expression of *Atoh1* in neonatal and older mice in the sensory neurons of the cochlea at virtually identical levels compared to newborns (Fig. 5B). Together these data firmly establish the presence of *Atoh1* in spiral and vestibular ganglion neurons.

Absence of *Atoh1* May Affect Pathfinding Properties of Sensory Neurons

We next wanted to understand whether *Atoh1* expression in sensory neurons is biologically meaningful. We reasoned that pathfinding properties of sensory neurons are most likely to be disturbed and investigated this issue using tract tracing and immunocytochemistry for tubulin. The fibers projecting to the posterior crista have previously been demonstrated to provide a good readout for molecular perturbation of pathfinding properties such as neurotrophin misexpression (Tessarollo et al., 2004) or semaphorin signaling (Gu et al., 2003). Specifically, *Bdnf* null mutant mice lose all canal crista innervation and do not receive many fibers even when sensory neurons cannot degenerate in the absence of *Bax* (Hellard et al., 2004), suggesting that hair cells lacking BDNF are not able to attract fibers. Loss of hair cells late in development as in *Pou4f3* null mutants (Xiang et al., 2003) or absence of differentiated hair cells as in E18.5 *Atoh1* null mutants is compatible with retention of

canal epithelia innervation (Fritzsch et al., 2005a). Moreover, misexpression of *Bdnf* under *Ntf3* promoter control redirects vestibular fibers away from vestibular hair cells and even to areas devoid of any hair cells (Tessarollo et al., 2004). Thus, lack of hair cell differentiation in *Atoh1* null mice combined with the presence of BDNF protein in canal epithelia should have no effect on the initial growth of canal epithelia afferent fibers.

Consistent with this expectation, we could not detect any obvious alterations in pathfinding at E11.5 (Fig. 8A,B), but found aberrations at E12.5 (Fig. 8C,D) and E13.5 (Fig. 8F,G). Specifically, fibers tended to branch profusely off the main branchlet to the posterior crista with only few fibers entering the epithelium (Fig. 8D,E) in *Atoh1* null mice. The aberrations observed resembled those previously reported (Gu et al., 2003) in targeted disruption of semaphorin signaling through its *neuropilin 1* receptor (Fig. 8E). In contrast, wildtype mice showed bifurcation of the posterior crista branchlet to enter the two hemicristae of the posterior cristae epithelium. Similar data were obtained for the anterior and horizontal crista and suggest that around the onset of *Bdnf*-mediated rerouting of sensory processes (Tessarollo et al., 2004), *Atoh1* null affects targeting of afferents near the sensory epithelia in a manner reminiscent of that in *Sema3a* signaling-disrupted mutants (Gu et al., 2003).

In summary, we show here that *Neurog1* null mice have a reduction in the number of hair cells and a reduced longitudinal growth of the cochlea. This is in part mediated by a terminal mitosis of hair cells of up to two days earlier and, to a lesser extent, by topographically altered upregulation of *Atoh1* in the apical turn of the cochlea and a more prominent expression of *Neurod1* in *Neurog1* null mice. We demonstrate the presence of *Atoh1* in sensory neurons using three *Atoh1* transgenic lines, *Atoh1-LacZ* heterozygotes, and Q-PCR of vestibular ganglion neurons and show a possible effect of *Atoh1* absence on neuronal pathfinding properties. Combined, these data support the notion that *Atoh1* is expressed in some common hair cell and sensory neuron precursors, in particular in the co-

Fig. 8. The effect of *Atoh1* null mutation (B, D, G) on the pattern of innervation of the posterior crista is compared with a wildtype (A, C, F) and a *npn1^{-sema3a}* mutant (E) as revealed with lipophilic tracers (A–F) and acetylated tubulin immunocytochemistry (G). The initial growth toward the posterior crista is unaffected in E11.5 embryos (A,B). However, afferent fibers continue to branch outside the sensory epithelium in both *Atoh1* null and *npn1^{-sema3a}* mutant embryos (D,E) whereas fibers have entered and branch inside the sensory epithelium in wildtype littermates (C). By E13.5, afferents have entered the epithelium and branched to reach the *Bdnf*-positive undifferentiated hair cells of the *Atoh1* null mutant. This branching is more profuse in wildtype animals and the fibers have split into two areas, each supplying the *Bdnf*-positive hair cells of the hemichristae (F). These data suggest that the absence of *Atoh1* interferes with the homing of afferent fibers near the target sensory epithelium. Scale bar = 100 μ m.




Fig. 8.

Fig. 9. This summary diagram shows the spatiotemporal progression of sensory neuron and hair cell terminal mitosis, and *Atoh1* expression in wildtype (top) and *Neurog1* null mice (bottom). Our data and those of Ruben (1967) suggest that in wildtype, the spiral ganglion terminal mitosis (Spgl) progresses from the base of the cochlea to the apex whereas the proliferation of hair cells of the organ of Corti progresses from apex to base with a broad overlap of terminal mitosis of both cell types in the apex around E11.5–12.5. The *Neurog1* null ear has no spiral neurons, and shows a premature onset and cessation of terminal mitosis in hair cells in an apex-to-base gradient, along a time line comparable to sensory neurons in wildtype mice. *Neurog1* null mice have fewer hair cells in a shortened and widened cochlea. *Neurog1* null mice also show some alteration in the topology of *Atoh1* expression. The right half of the scheme presents a hypothetical relationship between neurosensory precursors that give rise to both hair cells and sensory neurons. The model assumes that some neurosensory precursors will undergo a progressive up-regulation of *Atoh1* in *Neurog1*-positive precursors until eventually they will turn these cells into hair cell-producing precursors. It also assumes a mutual inhibition of *Neurog1* and *Atoh1* in the same precursor. According to this model, absence of NEUROG1 protein in *Neurog1* null mutants causes premature commitment to hair cell fate and thus a terminal mitosis of hair cells at earlier developmental stages as well as truncation of the progenitor pool expansion, resulting in a shortened cochlea. [Modified after Ma et al., 1999; Gowan et al., 2001; Bertrand et al., 2002.]




Fig. 9.

chlea and saccule. What remains to be shown is that common neurosensory clones exist in mice as in chicken (Sato and Fekete, 2005) and that *Neurog1* is also co-expressed with *Atoh1* in neurosensory precursors giving rise to hair cells and neurons, a possibility that was recently suggested (Raft, unpublished data).

DISCUSSION

Past research in the ear has demonstrated that three bHLH genes are essential for the formation and/or differentiation of the two types of neurosensory cells of the ear, the hair cells, and the sensory neurons (Ma et al., 1998; Bermingham et al., 1999; Liu et al., 2000; Kim et al., 2001). Additional information on *Neurog1* and *Atoh1* null mutations have shown that loss of sensory neuron formation in *Neurog1* null mice affects hair cell formation (Ma et al., 2000). In *Atoh1* null mutants, undifferentiated and/or dying cells still form in the topology normally occupied by hair cells (Chen et al., 2002; Fritzscht et al., 2005a). It still is unclear how the interaction between *Neurog1* and *Atoh1*, resulting in overlapping effects upon the ear neurosensory epithelium, could be reconciled, especially in light of the complex interactions of bHLH genes known in other developing sensory systems (Bertrand et al., 2002; Wu et al., 2003; Akagi et al., 2004). The *Neurog1* and *Atoh1* interaction also suggests a possible clonal relationship between some sensory neurons and hair cells in mammals (Fritzscht and Beisel, 2004), which was recently demonstrated in chickens (Sato and Fekete, 2005). As previously pointed out, our data and those of others (Bermingham et al., 1999; Chen et al., 2002; Fritzscht et al., 2005a) do not support the recent idea that *Atoh1* is not only essential for hair cell differentiation but also for sensory epithelia formation (Woods et al., 2004). In fact, our data show that even in combined *Atoh1/Neurog1* double null mutants, there is a formation of *Bdnf-LacZ*-positive cells in a topology that closely corresponds to some of the sensory epithelia found in wildtype littermates, suggesting that neither of these two bHLH genes is essential for the formation of undifferentiated sensory epithelia precursors and their maintenance up to E18.5 (Fig. 1D). The ob-

served reduction in total length of the cochlea in *Neurog1* null mice exceeds the length of *Bdnf-LacZ*-expressing cells in *Atoh1* null mice. The data on length and area changes in *Neurog1* null, *Atoh1* null and double null mice suggest a simple additive effect, implying that *Neurog1* affects both hair cell and non-sensory epithelium formation (Fig. 1; Table 1). These data are more in line with the idea that other genes are responsible for setting up the neurosensory precursor domains of the ear (Zou et al., 2004; Daudet and Lewis, 2005; Kiernan et al., 2005).

Atoh1 in Sensory Neurons

With almost any technique employed to reveal *Atoh1* in the ear, we are able to demonstrate *Atoh1* expression not only in the hair cells but also in some spiral and vestibular sensory neurons (Figs. 4–7). In the spiral ganglion, we find labeled cells in particular in the apex, an area where past research suggests a more prominent presence of type II spiral neurons that project to the outer hair cells (Ryugo, 1992; Rubel and Fritzscht, 2002). The reported lack of projection deeper into the undifferentiated organ of Corti of the remaining sensory neuron afferent fibers of *Atoh1* null mutants at E18.5 (Fritzscht et al., 2005a) could indicate a lack of projection of type II neurons. Alternatively, it could relate to the limited attraction provided by the undifferentiated cells that express *Atoh1-LacZ* and *Bdnf-LacZ* (Fritzscht et al., 2005a) as these cells are apparently negative for any other hair cell or supporting cell protein marker tested thus far (Chen et al., 2002; Woods et al., 2004).

It remains somewhat unclear why none of the previous publications has indicated such an expression of *Atoh1* in sensory neurons. In large part, this may be related to the more prominent expression of *Atoh1* in hair cells combined with the known effects of *Atoh1* on hair cell maturation (Bermingham et al., 1999; Izumikawa et al., 2005) that diverted a complete inner ear expression analyses. In part this appears to be inherent to some of the techniques employed, like the *Atoh1-eGFP* reporter, which requires additional techniques to extract the rather faint signal from the background. A

compounding factor is likely the early formation of bone around the sensory neurons, which preclude proper penetration of detection solutions, thus leading to false negatives as we can detect *Atoh1* in spiral and vestibular sensory neurons in older animals in which bone encloses these neurons. Lastly, without reason to suspect expression of *Atoh1*, one would not dissect the vestibular ganglia and process them for PCR to detect the clearly present *Atoh1* signal (Fig. 5B).

In summary, *Atoh1* expression is not only in hair cells but also in some sensory neurons and may be functionally meaningful for the development of specific projection patterns (Fig. 8).

Neurog1 and Hair Cell Terminal Mitosis and Differentiation

Our past work has shown that many hair cells do not form in *Atoh1* null mice (Ma et al., 2000), in particular in the saccule and the cochlea. We now show that combined mutation of *Atoh1* and *Neurog1* leads to a simple additive effect of the *Atoh1* phenotype on the *Neurog1* phenotype (Fig. 1; Table 1). Our data confirm previous reports on a base-to-apex progression of sensory neuron and apex-to-base progression of hair cell terminal mitosis (Ruben, 1967). Minor differences in the timing of these events relate to differences in staging the mice and differences in gestational length of the mouse strains used. Most importantly, this past work has indicated that apical hair cells may become postmitotic before apical sensory neurons. We now show that there is a widespread overlap of timing of terminal mitosis in apical sensory neurons and hair cells around E11.5–E12.5 (Fig. 2). On closer examination of terminal mitosis, *Neurod1* and *Atoh1* expression in *Neurog1* null embryos shows both terminal mitosis and *Neurod1* and *Atoh1* expressions are temporally and/or spatially altered in *Neurog1* null mice. In fact, terminal mitosis of hair cells occurs up to two days earlier in *Neurog1* null mice. This temporal acceleration of hair cell terminal mitosis, combined with our inability to demonstrate earlier expression of *Atoh1*, implies that other genes are responsible for the cell fate

switch from sensory neurons to hair cells in the *Neurog1* null mice: instead of being able to differentiate as sensory neurons, postmitotic cells now become hair cells. The altered expression of *Neurod1* and *Atoh1* in the cochlea with a spatial shift toward the apex suggests that *Neurog1* is not only affecting cell fate selection but also expression of other bHLH genes.

The simplest way of explaining these data as well as the expression of *Atoh1* in sensory neurons would be the assumption that neurosensory precursors co-express *Atoh1* and *Neurog1* and switch their cell fate to hair cell differentiation in *Neurog1* null mice. (Fig. 9). However, we cannot exclude other models, namely that *Neurog1* absence disrupts coordinated cell cycling of nearby cells and affects their gene expression profiles, thus causing *Neurod1* upregulation in hair cells. This alternative scenario was previously raised in developing dorsal root ganglion (Ma et al., 1999). Clearly, the truncation of hair cell formation may relate to either direct effects in neurosensory precursors through interactions of these bHLH genes as previously reported (Gowan et al., 2001) or may interfere with the coordinated inhibitory processes or collaboration with other factors that set up sensory epithelia (Adam et al., 1998; Daudet and Lewis, 2005).

Similar effects on reduced clonal expansion and changes in cell fate assignment have been reported for null mice of the winged helix gene *Bf1* (now *Foxg1*). In these mutants, the clonal expansion of cortical precursors is truncated through premature terminal mitosis and cell fate is limited to only one cortical neuron type (Hanashima et al., 2002, 2004). It appears that the *Neurog1* null mutation has comparable effects on neurosensory precursors of the ear, limiting the fate choice to one (hair cells) and initiating an earlier terminal mitosis of these cells resulting in clonal compression. Clearly, not only are fewer hair cells formed in the smaller epithelia of *Neurog1* null mice, but these mutants also miss all sensory neurons, suggesting that NEUROG1 protein is essential for the complete clonal expansion of the entire neurosensory population of the ear, possibly including some hair cell precursors.

In summary, much like in the spinal cord (Gowan et al., 2001), these data could be most easily explained by assuming an inhibitory interaction of *Neurog1* and *Atoh1* in a set of neurosensory precursor cells of the ear (Fig. 9). Absence of this interaction in *Neurog1* null mice leads to temporal shifting of hair cell terminal mitosis through a phenotype switch of postmitotic cells. Recent work in chicken has demonstrated such a clonal relationship exists at a limited scale (Satoh and Fekete, 2005) and our data are compatible with such a relationship for mice but do not prove it. Direct evidence is still needed for mammals.

***Atoh1* and Hair Cell Terminal Mitosis**

Obviously, our *Atoh1* in situ hybridization data as well as our *Atoh1-LacZ* data strongly suggest that *Atoh1* expression is at best in a small set of cycling hair cell precursors in the basal turn, the last hair cells to become postmitotic in the cochlea. These data are in agreement with almost all other data on *Atoh1* expression in the cochlea published thus far (Birmingham et al., 1999; Zine et al., 2001; Chen et al., 2002). Only some recent *Atoh1-LacZ* data suggest a somewhat more profound expression of *Atoh1* throughout the cochlea at E13.5 (Woods et al., 2004), which partially agrees with our *Atoh1-LacZ* data (Fig. 4E). Combined, our data confirm the previous conclusion based on *Atoh1-eGFP* expression that *Atoh1* is not expressed in detectable levels in cycling precursors (Chen et al., 2002), except for a few cells at the basal tip of the organ of Corti.

A low level of *Atoh1* may be present but this possibility cannot be ruled out with in situ hybridization, *Atoh1-LacZ*, or *Atoh1-eGFP*. Indeed, our *Atoh1-Cre* data show somewhat earlier upregulation of *Atoh1* expression in the future vestibular epithelia at E11 (Fig. 4G) and the cochlea basal turn at E12.5 (Fig. 4D) as well as β -galactosidase labeling in sensory neurons at these developmental stages (Fig. 4). These data imply that some *Atoh1* expression in amounts not detectable by current histological techniques exist in the growing organ of Corti several hours or days prior to

E12.5. Recent data on another bHLH gene, *MyoD*, showed that more sensitive in situ techniques can demonstrate cellular distribution prior to detection with more established in situ techniques, providing cellular distribution information for already existing RT-PCR data (Gerhart et al., 2004). Indeed, it appears that there is a delay of approximately 12 h between the first detection of *Atoh1* with PCR and with *Atoh1-Cre* (Figs. 4,5). This *Atoh1* expression revealed with the second *Cre* line is particularly remarkable as it is so early despite the fact that this reporter must be delayed through the multiple steps required for expression and eventual detection. This is apparent by the earlier *Atoh1-Cre* activation and detection in sensory neurons in *Atoh1-Cre/Rosa 26* lines. In the light of these data, combined with the clear effect of *Neurog1* on changes in timing of terminal mitosis of hair cells, we suggest that the spatio-temporal overlap of *Atoh1* expression with proliferating hair cell precursors might be more profound than our data and those of others suggest (Chen et al., 2002) as they may be biased toward detection only of larger amounts of *Atoh1* transcript or ATOH1 protein, rather than the earliest onset of expression. Using the newly developed *Atoh1-Cre* line combined with photoactivation of the β -galactosidase reaction product (Figs. 2, 4) should allow us to capture such expression in neurosensory precursors, if it exists.

As outlined above, it is possible that none of the employed histological techniques may detect the earliest onset of *Atoh1* expression. Such early upregulation of *Atoh1-Cre/Rosa 26* signal indicates that *Atoh1* transcription may exist in cochlea precursor cells as early as E10.5, a time in which the majority of hair cells are not yet postmitotic. We assume that this early expression is in common neurosensory precursors of sensory neurons and hair cells, leading to the formation of enough ATOH1 protein to activate the *Atoh1* promoter fragment that will drive the *Cre* expression that will eventually activate the LacZ reporter system of the Rosa 26 line, a notion supported by the early expression revealed in the second *Atoh1-Cre* line at E11 (Fig. 4G). However, while

not very likely, we cannot exclude that most hair cells and sensory neurons arise from discrete progenitors and that the early *Atoh1-Cre* expression reflects an early upregulation of *Atoh1* in some sensory neurons.

In many other systems, quantitative relationships within clones are determined by one or more bHLH genes through regulation of both cell fate and numbers of cells of a given fate (Bertrand et al., 2002; Akagi et al., 2004). Elimination of one bHLH gene may lead to replacement of a given cell type through clonal expansion and regulation of cell fate using another bHLH gene (Ma et al., 1999). In the ear, there appears to be a dichotomy of the bHLH gene channeling towards either a hair cell or sensory neuron fate. Thus, abrogation of *Neurog1* may cause premature upregulation of *Neurod1* and possibly other genes that can, however, in the context of the ear only cooperate in hair cell differentiation (Fig. 9). The temporal acceleration of terminal mitosis of hair cells will move these cells out of synchrony with other developing gene expression patterns, thus leading to the formation of multiple rows of hair cells and disorganization of apical specializations (Ma et al., 2000) presumably by moving hair cell differentiation out of synchrony with polarity-determining gene expressions such as *Vangl* (Amonlirdviman et al., 2005). Closer examination of *Vangl* upregulation (Montcouquiol et al., 2003) in *Neurog1* null mice and control littermates is needed to support this notion. Alternatively, the more profound upregulation of *Neurod1*-driven *LacZ* reporter in these hair cells of *Neurog1* null mice may affect cell polarity organization more directly.

Interestingly, our data on *Atoh1* expression, regardless of any genetic background, show that apical hair cells express *Atoh1* only in early neonates, but become postmitotic between E11.5 and E12.5. In contrast to *Atoh1* null mice, where *Atoh1* absence apparently leads to rapid degeneration of many of the undifferentiated *Atoh1-lacZ*-positive precursors found in these mutants (Chen et al., 2002; Fritzsche et al., 2005a), *Atoh1*-negative postmitotic hair cells of the apex in wildtype mice seem to do just fine for several days, even in *Atoh1* null mice

(Fritzsche et al., 2005a). Several explanations for this conflicting data are apparent. It is possible that cells require ATOH1 protein once the upstream regulators have initiated *Atoh1* expression and failure of ATOH1 protein formation leads to cell death within several hours. Available data on base-to-apex progression of apoptosis would be compatible with such an idea (Chen et al., 2002). Since the lack of ATOH1 protein will invariably lead to apoptosis, the reported base-to-apex progression of apoptosis in *Atoh1* null mutants might also relate to inherited specific maximal times postmitotic hair cells can survive without a specified amount of ATOH1 protein.

***Atoh1* Expression in Sensory Neurons May Be Biologically Meaningful**

The expression of *Atoh1* in the vestibular ganglion neurons, sensory neurons known to project to the saccule and the posterior crista (Maklad and Fritzsche, 1999), appears to relate to a possible effect of *Atoh1* null mutation on targeted afferent fiber growth (Fig. 8). Examination of initial vestibular fiber growth has shown unusual deviations of fibers near the target epithelia that could not be explained by lack of neurotrophins in the target epithelia (Fritzsche et al., 2005a). Clearly, *Bdnf*, the most important neurotrophin to guide and maintain afferent fibers to the crista organs (Fritzsche et al., 2004; Tessarollo et al., 2004), is expressed in these epithelia even in mutants lacking both *Neurog1* and *Atoh1* (Fig. 1H–J). Moreover, *Atoh1* null mice show a prominent innervation of the posterior crista as late as E18.5 (Fritzsche et al., 2005a) thus ruling out that *Bdnf* expression might be the problem as it would have resulted in loss of all crista innervation by E14.5 (Fritzsche et al., 2004; Tessarollo et al., 2004). We now have extended this initial finding and show aberrations that resemble those previously reported for *Npn1^{sema3a}*-mutant mice (Gu et al., 2003). We suggest that the expression of *Atoh1* in vestibular sensory neurons mediates certain path-finding properties of sensory neurons either by regulating neuropilin expression (Gu et al., 2003) or by affecting the BDNF/TRPC pathway for

guiding growth cones of vestibular ganglion neurons (Li et al., 2005; Wang and Poo, 2005). Such path-finding properties are essential for the connection of insect mechanosensory neurons with the CNS and may even be mediated by *Atoh1* knocked into *atonal* in flies (Ben-Arie et al., 2000; Wang et al., 2002). Governing path-finding may also be a property of *Atoh1* in CNS neurons in which it is expressed (Ben-Arie et al., 1997; Birmingham et al., 1999, 2001). Such involvement of *Atoh1* in regulating the molecular basis of path-finding may directly relate to the ancestry of the atonal gene family and their association with the sensory ciliated neuron (Fritzsche and Beisel, 2003) and the sensory system development and evolution (Fritzsche and Beisel, 2003; Fritzsche et al., 2005b). Testing this suggestion requires conditional null mutation of *Atoh1* only in sensory neurons using a *Neurog1*-mediated *Cre* expression. Alternatively, analyzing the expression of *Neuropilin1*, *TrkB*, and *TrpC* in wildtype and *Atoh1* null ears could reveal differential regulation of these genes known to be involved in path-finding.

In summary, our data support the notion that some sensory neurons express low levels of *Atoh1*, particularly those of the apex of the cochlea and the inferior vestibular ganglion known to innervate the saccule and posterior crista. *Neurog1* abrogation leads to earlier terminal mitosis of hair cells, somewhat displaced upregulation of *Atoh1* and *Neurod1*, and down-regulation of clonal expansion of hair cell precursors (Fig. 9). Consistent with recent data in chicken (Sato and Fekete, 2005), our data are compatible with a clonal relationship of some sensory neurons and hair cells in mice, being derived from common neurosensory precursors. However, this does by no means imply that all hair cells and all sensory neurons share a common neurosensory ancestor and more data are needed to demonstrate the extent of the postulated common neurosensory precursor. For translational research, it would be extremely helpful if the molecular switches from sensory neuron precursor to hair cell precursors are better understood and can be molecularly manipulated. Ultimately, this may lead to implantation of a single neurosensory

precursor population that can give rise to both sensory neurons and hair cells. Indeed, sensory neuron precursors that have the capacity to proliferate *in vitro* have recently been isolated from human cochlea (Rask-Andersen et al., 2005) and thus offer a possible therapeutic approach in combination with the above outlined insights.

ACKNOWLEDGMENTS

This work was supported by grants from NIH (RO1 DC005590 to B.F.; RO1 DC04279 to K.W.B.). We acknowledge the help of Dr. P. Chen and Q. Ma to improve the manuscript and Drs. P. Chen, H. Zoghbi, N. Segil, D. Ginty, and Q. Ma for providing their mutant mouse lines. This investigation was conducted in a facility constructed with support from Research Facilities Improvement Program Grant Number 1 C06 RR17417-01 from the National Center for Research Resources, National Institutes of Health. We acknowledge the use of the confocal microscope facility of the NCCB, supported by EPSCoR EPS-0346476 (CFD 47.076).

REFERENCES

- Adam J, Myat A, Le Roux I, Eddison M, Henrique D, Ish-Horowitz D, Lewis J. 1998. Cell fate choices and the expression of Notch, Delta and Serrate homologues in the chick inner ear: parallels with *Drosophila* sense-organ development. *Development* 125:4645–4654.
- Akagi T, Inoue T, Miyoshi G, Bessho Y, Takahashi M, Lee JE, Guillemot F, Kageyama R. 2004. Requirement of multiple basic helix-loop-helix genes for retinal neuronal subtype specification. *J Biol Chem* 279:28492–28498.
- Akazawa C, Ishibashi M, Shimizu C, Nakanishi S, Kageyama R. 1995. A mammalian helix-loop-helix factor structurally related to the product of *Drosophila* proneural gene *atonal* is a positive transcriptional regulator expressed in the developing nervous system. *J Biol Chem* 270:8730–8738.
- Amonlirdviman K, Khare NA, Tree DR, Chen WS, Axelrod JD, Tomlin CJ. 2005. Mathematical modeling of planar cell polarity to understand domineering nonautonomy. *Science* 307:423–426.
- Anderson DJ, Groves A, Lo L, Ma Q, Rao M, Shah NM, Sommer L. 1997. Cell lineage determination and the control of neuronal identity in the neural crest. *Cold Spring Harb Symp Quant Biol* 62:493–504.
- Beisel KW, Nelson NC, Delimont DC, Fritzsche B. 2000. Longitudinal gradients of KCNQ4 expression in spiral ganglion and cochlear hair cells correlate with progressive hearing loss in DFNA2. *Brain Res Mol Brain Res* 82:137–149.
- Ben-Arie N, Bellen HJ, Armstrong DL, McCall AE, Gordadze PR, Guo Q, Matzuk MM, Zoghbi HY. 1997. *Math1* is essential for genesis of cerebellar granule neurons. *Nature* 390:169–172.
- Ben-Arie N, Hassan BA, Bermingham NA, Malicki DM, Armstrong D, Matzuk M, Bellen HJ, Zoghbi HY. 2000. Functional conservation of *atonal* and *Math1* in the CNS and PNS. *Development* 127:1039–1048.
- Bermingham NA, Hassan BA, Price SD, Vollrath MA, Ben-Arie N, Eatock RA, Bellen HJ, Lysakowski A, Zoghbi HY. 1999. *Math1*: an essential gene for the generation of inner ear hair cells. *Science* 284:1837–1841.
- Bermingham NA, Hassan BA, Wang VY, Fernandez M, Banfi S, Bellen HJ, Fritzsche B, Zoghbi HY. 2001. Proprioceptor pathway development is dependent on *Math1*. *Neuron* 30:411–422.
- Bertrand N, Castro DS, Guillemot F. 2002. Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* 3:517–530.
- Calegari F, Huttner WB. 2003. An inhibition of cyclin-dependent kinases that lengthens, but does not arrest, neuroepithelial cell cycle induces premature neurogenesis. *J Cell Sci* 116:4947–4955.
- Chen P, Johnson JE, Zoghbi HY, Segil N. 2002. The role of *Math1* in inner ear development: Uncoupling the establishment of the sensory primordium from hair cell fate determination. *Development* 129:2495–2505.
- Daudet N, Lewis J. 2005. Two contrasting roles for Notch activity in chick inner ear development: specification of prosensory patches and lateral inhibition of hair-cell differentiation. *Development* 132:541–551.
- Farinas I, Jones KR, Tessarollo L, Vigers AJ, Huang E, Kirstein M, de Caprona DC, Coppola V, Backus C, Reichardt LF, Fritzsche B. 2001. Spatial shaping of cochlear innervation by temporally regulated neurotrophin expression. *J Neurosci* 21:6170–6180.
- Fritzsche B, Beisel KW. 2003. Molecular conservation and novelties in vertebrate ear development. *Curr Top Dev Biol* 57:1–44.
- Fritzsche B, Beisel KW. 2004. Keeping sensory cells and evolving neurons to connect them to the brain: molecular conservation and novelties in vertebrate ear development. *Brain Behav Evol* 64:182–197.
- Fritzsche B, Farinas I, Reichardt LF. 1997. Lack of neurotrophin 3 causes losses of both classes of spiral ganglion neurons in the cochlea in a region-specific fashion. *J Neurosci* 17:6213–6225.
- Fritzsche B, Beisel KW, Bermingham NA. 2000. Developmental evolutionary biology of the vertebrate ear: conserving mechanoelectric transduction and developmental pathways in diverging morphologies. *Neuroreport* 11:R35–44.
- Fritzsche B, Beisel KW, Jones K, Farinas I, Maklad A, Lee J, Reichardt LF. 2002. Development and evolution of inner ear sensory epithelia and their innervation. *J Neurobiol* 53:143–156.
- Fritzsche B, Tessarollo L, Coppola E, Reichardt LF. 2004. Neurotrophins in the ear: their roles in sensory neuron survival and fiber guidance. *Prog Brain Res* 146:265–278.
- Fritzsche B, Matei VA, Nichols DH, Bermingham N, Jones K, Beisel KW, Wang VY. 2005a. *Atoh1* null mice show directed afferent fiber growth to undifferentiated ear sensory epithelia followed by incomplete fiber retention. *Dev Dyn* 233:570–583.
- Fritzsche B, Piatigorsky J, Tessmar-Raible K, Jekely G, Guy K, Raible F, Wittbrodt J, Arendt D. 2005b. Ancestry of photic and mechanosensation? *Science* 308:1113–1114.
- Gerhart J, Baytion M, Perlman J, Neely C, Heaton B, Nilsen T, Getts R, Kadushin J, George-Weinstein M. 2004. Visualizing the needle in the haystack: *in situ* hybridization with fluorescent dendrimers. *Biol Proceed Online* 6:149–156.
- Gowan K, Helms AW, Hunsaker TL, Collisson T, Ebert PJ, Odom R, Johnson JE. 2001. Crossinhibitory activities of *Ngn1* and *Math1* allow specification of distinct dorsal interneurons. *Neuron* 31:219–232.
- Gu C, Rodriguez ER, Reimert DV, Shu T, Fritzsche B, Richards LJ, Kolodkin AL, Ginty DD. 2003. *Neuropilin-1* conveys semaphorin and VEGF signaling during neural and cardiovascular development. *Dev Cell* 5:45–57.
- Hanashima C, Shen L, Li SC, Lai E. 2002. *Brain factor-1* controls the proliferation and differentiation of neocortical progenitor cells through independent mechanisms. *J Neurosci* 22:6526–6536.
- Hanashima C, Li SC, Shen L, Lai E, Fishell G. 2004. *Foxg1* suppresses early cortical cell fate. *Science* 303:56–59.
- Hellard D, Brosenitsch T, Fritzsche B, Katz DM. 2004. Cranial sensory neuron development in the absence of brain-derived neurotrophic factor in *BDNF/Bax* double null mice. *Dev Biol* 275:34–43.
- Ito H, Nakajima A, Nomoto H, Furukawa S. 2003. Neurotrophins facilitate neuronal differentiation of cultured neural stem cells via induction of mRNA expression of basic helix-loop-helix transcription factors *Mash1* and *Math1*. *J Neurosci Res* 71:648–658.
- Izumikawa M, Minoda R, Kawamoto K, Abrashkin KA, Swiderski DL, Dolan DF, Brough DE, Raphael Y. 2005. Auditory hair cell replacement and hearing improvement by *Atoh1* gene therapy in deaf mammals. *Nat Med* 11:271–276.
- Jones KR, Farinas I, Backus C, Reichardt LF. 1994. Targeted disruption of the *BDNF* gene perturbs brain and sensory neuron development but not motor neuron development. *Cell* 76:989–999.
- Kiernan AE, Pelling AL, Leung KK, Tang AS, Bell DM, Tease C, Lovell-Badge R, Steel KP, Cheah KS. 2005. *Sox2* is re-

- quired for sensory organ development in the mammalian inner ear. *Nature* 434:1031–1035.
- Kim J, Wu HH, Lander AD, Lyons KM, Matzuk MM, Calof AL. 2005. GDF11 Controls the timing of progenitor cell competence in developing retina. *Science* 308:1927–1930.
- Kim WY, Fritzscht B, Serls A, Bakel LA, Huang EJ, Reichardt LF, Barth DS, Lee JE. 2001. NeuroD-null mice are deaf due to a severe loss of the inner ear sensory neurons during development. *Development* 128:417–426.
- Lanford PJ, Shailam R, Norton CR, Gridley T, Kelley MW. 2000. Expression of Math1 and HES5 in the cochleae of wild-type and Jag2 mutant mice. *J Assoc Res Otolaryngol* 1:161–171.
- Li Y, Jia YC, Cui K, Li N, Zheng ZY, Wang YZ, Yuan XB. 2005. Essential role of TRPC channels in the guidance of nerve growth cones by brain-derived neurotrophic factor. *Nature* 434:894–898.
- Liu M, Pereira FA, Price SD, Chu MJ, Shope C, Himes D, Eatock RA, Brownell WE, Lysakowski A, Tsai MJ. 2000. Essential role of BETA2/NeuroD1 in development of the vestibular and auditory systems. *Genes Dev* 14:2839–2854.
- Lumpkin EA, Collisson T, Parab P, Omer-Abdalla A, Haeberle H, Chen P, Doetschhofer A, White P, Groves A, Segil N, Johnson JE. 2003. Math1-driven GFP expression in the developing nervous system of transgenic mice. *Gene Expr Patterns* 3:389–395.
- Ma Q, Chen Z, del Barco Barrantes I, de la Pompa JL, Anderson DJ. 1998. neurogenin1 is essential for the determination of neuronal precursors for proximal cranial sensory ganglia. *Neuron* 20:469–482.
- Ma Q, Fode C, Guillemot F, Anderson DJ. 1999. Neurogenin1 and neurogenin2 control two distinct waves of neurogenesis in developing dorsal root ganglia. *Genes Dev* 13:1717–1728.
- Ma Q, Anderson DJ, Fritzscht B. 2000. Neurogenin 1 null mutant ears develop fewer, morphologically normal hair cells in smaller sensory epithelia devoid of innervation. *J Assoc Res Otolaryngol* 1:129–143.
- Maklad A, Fritzscht B. 1999. Incomplete segregation of endorgan-specific vestibular ganglion cells in mice and rats. *J Vestib Res* 9:387–399.
- Matei VA, Feng F, Pauley S, Beisel K, Nichols M, Fritzscht B. 2005. Two photon laser illuminations photoactivates the fluorescence absorbing X-Gal reaction product to produce a fluorescent signal with superior visibility and stability. *Brain Res Bull* (in press).
- Montcouquiol M, Rachel RA, Lanford PJ, Copeland NG, Jenkins NA, Kelley MW. 2003. Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. *Nature* 423:173–177.
- Ohyama T, Groves AK. 2004. Generation of Pax2-Cre mice by modification of a Pax2 bacterial artificial chromosome. *Genesis* 38:195–199.
- Radde-Gallwitz K, Pan L, Gan L, Lin X, Segil N, Chen P. 2004. Expression of Islet1 marks the sensory and neuronal lineages in the mammalian inner ear. *J Comp Neurol* 477:412–421.
- Rask-Andersen H, Bostrom M, Gerdin B, Kinnefors A, Nyberg G, Engstrand T, Miller JM, Lindholm D. 2005. Regeneration of human auditory nerve. In vitro/in vivo demonstration of neural progenitor cells in adult human and guinea pig spiral ganglion. *Hear Res* 203:180–191.
- Rubel EW, Fritzscht B. 2002. AUDITORY SYSTEM DEVELOPMENT: Primary Auditory Neurons and Their Targets. *Annu Rev Neurosci* 25:51–101.
- Rubel RJ. 1967. Development of the inner ear of the mouse: a radioautographic study of terminal mitoses. *Acta Otolaryngol:Suppl* 220:221–244.
- Ryugo DK. 1992. The auditory nerve; peripheral innervation, cell body morphology, and central projections. In: Webster DB, Popper AN, Fay RR, editors. The mammalian auditory pathway: neuroanatomy. New York: Springer-Verlag. p 23–65.
- Satoh T, Fekete DM. 2005. Clonal analysis of the relationships between mechanosensory cells and the neurons that innervate them in the chicken ear. *Development* 132:1687–1697.
- Seipel K, Yanze N, Schmid V. 2004. Developmental and evolutionary aspects of the basic helix-loop-helix transcription factors Atonal-like 1 and Achaete-scute homolog 2 in the jellyfish. *Dev Biol* 269:331–345.
- Tessarollo L, Coppola V, Fritzscht B. 2004. NT-3 replacement with brain-derived neurotrophic factor redirects vestibular nerve fibers to the cochlea. *J Neurosci* 24:2575–2584.
- Theiler K. 1989. The house mouse. New York: Springer-Verlag. 178 p.
- Wang GX, Poo MM. 2005. Requirement of TRPC channels in netrin-1-induced chemotropic turning of nerve growth cones. *Nature* 434:898–904.
- Wang VY, Hassan BA, Bellen HJ, Zoghbi HY. 2002. Drosophila atonal fully rescues the phenotype of Math1 null mice: new functions evolve in new cellular contexts. *Curr Biol* 12:1611–1616.
- Woods C, Montcouquiol M, Kelley MW. 2004. Math1 regulates development of the sensory epithelium in the mammalian cochlea. *Nat Neurosci* 7:1310–1318.
- Wu HH, Ivkovic S, Murray RC, Jaramillo S, Lyons KM, Johnson JE, Calof AL. 2003. Autoregulation of neurogenesis by GDF11. *Neuron* 37:197–207.
- Xiang M, Maklad A, Pirvola U, Fritzscht B. 2003. Brn3c null mutant mice show long-term, incomplete retention of some afferent inner ear innervation. *BMC Neurosci* 4:2.
- Yang Q, Bermingham NA, Finegold MJ, Zoghbi HY. 2001. Requirement of Math1 for secretory cell lineage commitment in the mouse intestine. *Science* 294:2155–2158.
- Zine A, Aubert A, Qiu J, Therianos S, Guillemot F, Kageyama R, de Ribaupierre F. 2001. Hes1 and Hes5 activities are required for the normal development of the hair cells in the mammalian inner ear. *J Neurosci* 21:4712–4720.
- Zou D, Silvius D, Fritzscht B, Xu PX. 2004. Eya1 and Six1 are essential for early steps of sensory neurogenesis in mammalian cranial placodes. *Development* 131:5561–5572.