



Health Information for Travelers to Indonesia



Travel Notices in Effect

- [Rabies in Bali, Indonesia \(/travel/content/outbreak-notice/rabies-bali-indonesia2008.aspx\)](/travel/content/outbreak-notice/rabies-bali-indonesia2008.aspx) March 01, 2011
- [Update: Dengue, Tropical and Subtropical Regions \(/travel/content/outbreak-notice/dengue-tropical-sub-tropical.aspx\)](/travel/content/outbreak-notice/dengue-tropical-sub-tropical.aspx) January 18, 2011
- [Guidelines and Recommendations: Interim Guidance about Avian Influenza \(H5N1\) for U.S. Citizens Living Abroad \(/travel/content/avian-flu-americans-abroad.aspx\)](/travel/content/avian-flu-americans-abroad.aspx) January 13, 2011
- [Chikungunya Fever in Asia and the Indian Ocean \(/travel/content/outbreak-notice/chikungunya-fever.aspx\)](/travel/content/outbreak-notice/chikungunya-fever.aspx) December 06, 2010
- [2010 Measles Update \(/travel/content/in-the-news/measles.aspx\)](/travel/content/in-the-news/measles.aspx) September 09, 2010
- [Human Infection with Avian Influenza A \(H5N1\) Virus: Advice for Travelers \(/travel/content/avian-flu-asia.aspx\)](/travel/content/avian-flu-asia.aspx) November 04, 2008

[Top of Page \(#top\)](#)

Safety and Security Abroad

- [Registration of Traveler Emergency Contact and Itinerary Information \(/travel/content/register-contact-info.aspx\)](/travel/content/register-contact-info.aspx) January 13, 2011
- [Transportation Security Administration \(http://www.tsa.gov\)](http://www.tsa.gov)
- [U.S. Department of State \(http://travel.state.gov\)](http://travel.state.gov)

[Top of Page \(#top\)](#)

Preparing for Your Trip to Indonesia

Before visiting Indonesia, you may need to get the following vaccinations and medications for vaccine-preventable diseases and other diseases you might be at risk for at your destination: (Note: Your doctor or health-care provider will determine what you will need, depending on factors such as your health and immunization history, areas of the country you will be visiting, and planned activities.)

To have the most benefit, see a health-care provider at least 4–6 weeks before your trip to allow time for your vaccines to take effect and to start taking medicine to prevent malaria, if you need it.

Even if you have less than 4 weeks before you leave, you should still see a health-care provider for needed vaccines, anti-malaria drugs and other medications and information about how to protect yourself from illness and injury while traveling.

CDC recommends that you see a health-care provider who specializes in Travel Medicine. [Find a travel medicine clinic \(http://www.cdc.gov/travel/content/TravelClinics.aspx\)](http://www.cdc.gov/travel/content/TravelClinics.aspx) near you. If you have a medical condition, you should also share your travel plans with any doctors you are currently seeing for other medical reasons.

If your travel plans will take you to more than one country during a single trip, be sure to let your health-care provider know so that you can receive the appropriate vaccinations and information for all of your destinations. Long-term travelers, such as those who plan to work or study abroad, may also need additional vaccinations as required by their employer or school.

Although yellow fever is not a disease risk in Indonesia, the government requires travelers arriving from countries where yellow fever is present (</travel/yellowbook/2010/chapter-2/yellow-fever.aspx#823>) to present proof of yellow fever vaccination. If you will be traveling to one of these countries where yellow fever is present before arriving in Indonesia, this requirement must be taken into consideration.

Be sure your routine vaccinations are up-to-date. Check the links below to see which vaccinations adults and children should get.

Routine vaccines, as they are often called, such as for influenza, chickenpox (or varicella), polio, measles/mumps/rubella (MMR), and diphtheria/pertussis/tetanus (DPT) are given at all stages of life; see the childhood and adolescent immunization schedule (<http://www.cdc.gov/vaccines/recs/schedules/child-schedule.htm>) and routine adult immunization schedule (<http://www.cdc.gov/vaccines/recs/schedules/adult-schedule.htm>).

Routine vaccines are recommended even if you do not travel. Although childhood diseases, such as measles, rarely occur in the United States, they are still common in many parts of the world. A traveler who is not vaccinated would be at risk for infection.

Vaccine recommendations are based on the best available risk information. Please note that the level of risk for vaccine-preventable diseases can change at any time.

| Vaccination or Disease | Recommendations or Requirements for Vaccine-Preventable Diseases |
|--|--|
| Routine (http://www.cdc.gov/vaccines/recs/schedules/default.htm) | Recommended if you are not up-to-date with routine shots such as, measles/mumps/rubella (MMR) vaccine, diphtheria/pertussis/tetanus (DPT) vaccine, poliovirus vaccine, etc. |
| Hepatitis A (/travel/yellowbook/2008/ch4/hep-a.aspx) or immune globulin (IG) | Recommended for all unvaccinated people traveling to or working in countries with an intermediate or high level of hepatitis A virus infection (see <u>map</u> (/travel/yellowbook/2010/chapter-2/hepatitis-a.aspx#761)) where exposure might occur through food or water. Cases of travel-related hepatitis A can also occur in travelers to developing countries with "standard" tourist itineraries, accommodations, and food consumption behaviors. |
| Hepatitis B (/travel/yellowbook/2008/ch4/hep-b.aspx) | Recommended for all unvaccinated persons traveling to or working in countries with intermediate to high levels of endemic HBV transmission (see <u>map</u> (/travel/yellowbook/2010/chapter-2/hepatitis-b.aspx#849)), especially those who might be exposed to blood or body fluids, have sexual contact with the local population, or be exposed through medical treatment (e.g., for an accident). |
| Typhoid (/travel/yellowbook/2008/ch4/typhoid.aspx) | Recommended for all unvaccinated people traveling to or working in Southeast Asia, especially if staying with friends or relatives or visiting smaller cities, villages, or rural areas where exposure might occur through food or water. |
| Japanese encephalitis (/travel/yellowbook/2008/ch4/japanese-encephalitis.aspx) | Recommended if you plan to visit rural farming areas and under special circumstances, such as a known outbreak of Japanese encephalitis, see <u>country-specific information</u> (/travel/yellowbook/2010/chapter-2/japanese-encephalitis.aspx#811). |
| Rabies (/travel/yellowbook/2008/ch4/rabies.aspx) | Recommended for travelers spending a lot of time outdoors, especially in rural areas, involved in activities such as bicycling, camping, or hiking. Also recommended for travelers with significant occupational risks (such as veterinarians), for long-term travelers and expatriates living in areas with a significant risk of exposure, and for travelers involved in any activities that might bring them into direct contact with bats, carnivores, and other mammals. Children are considered at higher risk because they tend to play with animals, may receive more severe bites, or may not report bites. |

Areas of Indonesia with Malaria: Rural areas of Sumatra, Sulawesi, Kalimantan (Borneo) and Nusa Tenggara Barat (includes island of Lombok)

All areas of eastern Indonesia (provinces of Papua Indonesia, Irian Jaya Barat, Nusa Tenggara Timur, Maluku, and Maluku Utara).

None in Jakarta or resort areas of Bali and Java.

Low transmission in rural areas of Java. (more information (</travel/yellowbook/2010/chapter-2/malaria-risk-information-and-prophylaxis/Indonesia.aspx>))

If you will be visiting an area of Indonesia with malaria, you will need to discuss with your doctor the best ways for you to avoid getting sick with malaria. Ways to prevent malaria include the following:

- Taking a prescription antimalarial drug
- Using insect repellent and wearing long pants and sleeves to prevent mosquito bites
- Sleeping in air-conditioned or well-screened rooms or using bednets

All of the following antimalarial drugs are equal options for preventing malaria in Indonesia: Atovaquone/proguanil, doxycycline, or mefloquine. For detailed information about each of these drugs, see Table 2-23: Drugs used in the prophylaxis of malaria (</travel/yellowbook/2010/chapter-2/malaria.aspx#990>). For information that can help you and your doctor decide which of these drugs would be best for you, please see Choosing a

[Drug to Prevent Malaria \(http://www.cdc.gov/malaria/travelers/drugs.html\)](http://www.cdc.gov/malaria/travelers/drugs.html).

Note: Chloroquine is NOT an effective antimalarial drug in Indonesia and should not be taken to prevent malaria in this region.

To find out more information on malaria throughout the world, you can use the [interactive CDC malaria map \(http://www.cdc.gov/malaria/map/index.html\)](http://www.cdc.gov/malaria/map/index.html). You can search or browse countries, cities, and place names for more specific malaria risk information and the recommended prevention medicines for that area.

Malaria Contact for Health-Care Providers

For assistance with the diagnosis or management of suspected cases of malaria, call the CDC Malaria Hotline: **770-488-7788** (M-F, 9 am-5 pm, Eastern time). For emergency consultation after hours, call **770-488-7100** and ask to speak with a CDC Malaria Branch clinician.

A Special Note about Antimalarial Drugs

You should purchase your antimalarial drugs before travel. Drugs purchased overseas may not be manufactured according to United States standards and may not be effective. They also may be dangerous, contain counterfeit medications or contaminants, or be combinations of drugs that are not safe to use.

Halofantrine (marketed as Halfan) is widely used overseas to treat malaria. CDC recommends that you do **NOT** use halofantrine because of serious heart-related side effects, including deaths. You should avoid using antimalarial drugs that are not recommended **unless** you have been diagnosed with life-threatening malaria and no other options are immediately available.

For detailed information about these antimalarial drugs, see [Choosing a Drug to Prevent Malaria \(http://www.cdc.gov/malaria/travelers/drugs.html\)](http://www.cdc.gov/malaria/travelers/drugs.html).

More Information About Malaria

Malaria is always a serious disease and may be a deadly illness. Humans get malaria from the bite of a mosquito infected with the parasite. Prevent this serious disease by seeing your health-care provider for a prescription antimalarial drug and by protecting yourself against mosquito bites ([see below \(#insect\)](#)).

Travelers to malaria risk-areas in Indonesia, including infants, children, and former residents of Indonesia, should take one of the antimalarial drugs listed in the box above.

Symptoms

Malaria symptoms may include

- fever
- chills
- sweats
- headache
- body aches
- nausea and vomiting
- fatigue

Malaria symptoms will occur at least 7 to 9 days after being bitten by an infected mosquito. Fever in the first week of travel in a malaria-risk area is unlikely to be malaria; however, you should see a doctor right away if you develop a fever during your trip.

Malaria may cause anemia and jaundice. Malaria infections with *Plasmodium falciparum*, if not promptly treated, may cause kidney failure, coma, and death. Despite using the protective measures outlined above, travelers may still develop malaria up to a year after returning from a malarious area. You should see a doctor immediately if you develop a fever anytime during the year following your return and tell the physician of your travel.

Medicines you may need:

- **The prescription medicines you take every day.** Make sure you have enough to last during your trip. Keep them in their original prescription bottles and always in your carry-on luggage. [Be sure to follow security guidelines \(http://www.tsa.gov\)](http://www.tsa.gov), if the medicines are liquids.
- Antimalarial drugs, if traveling to a [malaria-risk area \(#mal_risk\)](#) in Indonesia and prescribed by your doctor.
- **Medicine for diarrhea**, usually over-the-counter.

Note: Some drugs available by prescription in the US are illegal in other countries. Check the US Department of State [Consular Information Sheets \(http://travel.state.gov/travel/cis_pa_tw/cis/cis_1765.html\)](http://travel.state.gov/travel/cis_pa_tw/cis/cis_1765.html) for the country(s) you intend to visit or the embassy or consulate for that country(s). If your medication is not allowed in the country you will be visiting, ask your health-care provider to write a letter on office stationery stating the medication has been prescribed for you.

Other items you may need:

- Iodine tablets and portable water filters to purify water if bottled water is not available. See [A Guide to Water Filters \(http://www.cdc.gov/crypto/factsheets/filters.html\)](http://www.cdc.gov/crypto/factsheets/filters.html), [A Guide to Commercially-Bottled Water and Other Beverages \(http://www.cdc.gov/crypto/factsheets/bottled.html\)](http://www.cdc.gov/crypto/factsheets/bottled.html), and [Safe Food and Water \(http://www.cdc.gov/travel/content/safe-food-water.aspx\)](http://www.cdc.gov/travel/content/safe-food-water.aspx) for more detailed information.
- Sunblock and sunglasses for protection from harmful effects of UV sun rays. See [Basic Information about Skin Cancer \(http://www.cdc.gov/cancer/skin/basic_info/index.htm\)](http://www.cdc.gov/cancer/skin/basic_info/index.htm) for more information.
- Antibacterial hand wipes or alcohol-based hand sanitizer containing at least 60% alcohol.
- To prevent insect/mosquito bites, bring:
 - Lightweight long-sleeved shirts, long pants, and a hat to wear outside, whenever possible.
 - Flying-insect spray to help clear rooms of mosquitoes. The product should contain a pyrethroid insecticide; these insecticides quickly kill flying insects, including mosquitoes.
 - Bed nets treated with permethrin, if you will not be sleeping in an air-conditioned or well-screened room and will be in malaria-risk areas. For use and purchasing information, see [Insecticide Treated Bed Nets \(http://www.cdc.gov/malaria/malaria_worldwide/reduction/tn.html\)](http://www.cdc.gov/malaria/malaria_worldwide/reduction/tn.html) on the CDC malaria site. Overseas, permethrin or another insecticide, deltamethrin, may be purchased to treat bed nets and clothes.

See other suggested over-the-counter medications and first aid items for a [travelers' health kit \(http://www.cdc.gov/travel/yellowBookCh2-HealthKit.aspx\)](http://www.cdc.gov/travel/yellowBookCh2-HealthKit.aspx).

Note: Check the [Air Travel section \(http://www.tsa.gov/travelers/airtravel/index.shtm\)](http://www.tsa.gov/travelers/airtravel/index.shtm) of the [Transportation Security Administration \(http://www.tsa.gov/\)](http://www.tsa.gov/) website for the latest information about airport screening procedures and prohibited items.

[Top of Page \(#top\)](#)

Other Diseases Found in Southeast Asia

Risk can vary between countries within this region and also within a country; the quality of in-country surveillance also varies.

The following are disease risks that might affect travelers; this is not a complete list of diseases that can be present. Environmental conditions may also change, and up to date information about risk by regions within a country may also not always be available.

[Dengue \(http://www.cdc.gov/travel/yellowBookCh4-DengueFever.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-DengueFever.aspx), [chikungunya, filariasis \(http://www.cdc.gov/travel/yellowBookCh4-FilariasisLymphatic.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-FilariasisLymphatic.aspx), [Japanese encephalitis \(http://www.cdc.gov/travel/yellowBookCh4-JapaneseEncephalitis.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-JapaneseEncephalitis.aspx), and [plague \(http://www.cdc.gov/travel/yellowBookCh4-Plague.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Plague.aspx) are diseases carried by insects that also occur in this region. Protecting yourself against insect bites ([see below \(#insect\)](#)) will help to prevent these diseases.

Highly pathogenic avian influenza (H5N1) continues to cause outbreaks in domestic and wild bird populations and has caused human cases in several countries in Southeast Asia. In 2006, the virus continued to spread in poultry populations in Indonesia. Avoid all direct contact with birds, including domestic poultry (such as chickens and ducks) and wild birds, and avoid places such as poultry farms and bird markets where live birds are raised or kept. For a current list of countries reporting outbreaks of H5N1 among poultry and/or wild birds, view [updates from the World Organization for Animal Health \(OIE\) \(http://www.who.int/animal-disease/news/2006/06/20060620_h5n1_updates\)](http://www.who.int/animal-disease/news/2006/06/20060620_h5n1_updates), and for total numbers of confirmed human cases of H5N1 virus by country see the [World Health Organization \(WHO\) Avian Influenza website \(http://www.who.int/avian-influenza/\)](http://www.who.int/avian-influenza/).

Do not swim in fresh water (except in well-chlorinated swimming pools) to avoid infection with [schistosomiasis \(http://www.cdc.gov/travel/yellowBookCh4-Schistosomiasis.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Schistosomiasis.aspx). [Leptospirosis \(http://www.cdc.gov/travel/yellowBookCh4-Leptospirosis.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Leptospirosis.aspx), a bacterial infection often contracted through recreational water activities in contaminated water, such as kayaking, is common in tropical areas of Southeast Asia. An outbreak was reported in Borneo among expedition travelers in 2000.

[Measles \(http://www.cdc.gov/travel/yellowBookCh4-Measles.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Measles.aspx) transmission persists in the region, although vaccination coverage is improving in some countries in Southeast Asia. Influenza infections can occur throughout the year in tropical areas.

[Polio \(http://www.cdc.gov/travel/yellowBookCh4-Poliomyelitis.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Poliomyelitis.aspx) resurfaced in Indonesia in 2005. Imported cases in neighboring countries have occasionally occurred.

[Top of Page \(#top\)](#)

Staying Healthy During Your Trip

Many diseases, like [malaria \(http://www.cdc.gov/travel/yellowBookCh4-Malaria.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-Malaria.aspx) and [dengue \(http://www.cdc.gov/travel/yellowBookCh4-DengueFever.aspx\)](http://www.cdc.gov/travel/yellowBookCh4-DengueFever.aspx), are spread through insect bites. One of the best protections is to prevent insect bites by:

- Using insect repellent (bug spray) with 30%-50% DEET. Picaridin, available in 7% and 15% concentrations, needs more frequent application. There is less information available on how effective picaridin is at protecting against all of the types of mosquitoes that transmit malaria.
- Wearing long-sleeved shirts, long pants, and a hat outdoors.

- Remaining indoors in a screened or air-conditioned area during the peak biting period for malaria (dusk and dawn).
- Sleeping in beds covered by nets treated with permethrin, if not sleeping in an air-conditioned or well-screened room.
- Spraying rooms with products effective against flying insects, such as those containing pyrethroid.

For detailed information about insect repellent use, see [Insect and Arthropod Protection \(/travel/contentInsectProtection.aspx\)](/travel/contentInsectProtection.aspx).

Direct contact with animals can spread diseases like rabies or cause serious injury or illness. It is important to prevent animal bites and scratches.

- Be sure you are up to date with tetanus vaccination.
- Do not touch or feed any animals, including dogs and cats. Even animals that look like healthy pets can have rabies or other diseases.
- Help children stay safe by supervising them carefully around all animals.
- If you are bitten or scratched, wash the wound well with soap and water and **go to a doctor right away**.
- After your trip, be sure to tell your doctor or state health department if you were bitten or scratched during travel.

For more information about rabies and travel, see the [Rabies chapter \(/travel/yellowBookCh4-Rabies.aspx\)](/travel/yellowBookCh4-Rabies.aspx) of the [Yellow Book \(/travel/contentYellowBook.aspx\)](/travel/contentYellowBook.aspx) or [CDC's Rabies homepage \(http://www.cdc.gov/rabies/\)](http://www.cdc.gov/rabies/). For more information about how to protect yourself from other risks related to animals, see [Animal-Associated Hazards \(/travel/yellowBookCh6-Animal.aspx\)](/travel/yellowBookCh6-Animal.aspx). Diseases from food and water are the leading cause of illness in travelers. Follow these tips for safe eating and drinking:

- Wash your hands often with soap and water, especially before eating. If soap and water are not available, use an alcohol-based hand gel (with at least 60% alcohol).
- Drink only bottled or boiled water, or carbonated (bubbly) drinks in cans or bottles. Avoid tap water, fountain drinks, and ice cubes. If this is not possible, learn how to [make water safer to drink \(/travel/contentWaterTreatment.aspx\)](/travel/contentWaterTreatment.aspx).
- Do not eat food purchased from street vendors.
- Make sure food is fully cooked.
- Avoid dairy products, unless you know they have been pasteurized.

Diseases from food and water often cause vomiting and diarrhea. Make sure to bring diarrhea medicine with you so that you can treat mild cases yourself.

Car crashes are a leading cause of [injury \(http://wwwnc.cdc.gov/travel/yellowBookCh6-Injuries.aspx\)](http://wwwnc.cdc.gov/travel/yellowBookCh6-Injuries.aspx) among travelers. Protect yourself from these injuries by:

- Not drinking and driving.
- Wearing your seat belt and using car seats or booster seats in the backseat for children.
- Following local traffic laws.
- Wearing helmets when you ride bikes, motorcycles, and motor bikes.
- Not getting on an overloaded bus or mini-bus.
- Hiring a local driver, when possible.
- Avoiding night driving.

- To avoid infections such as HIV and viral hepatitis do not share needles for tattoos, body piercing, or injections.
- To reduce the risk of HIV and other sexually transmitted diseases always use latex condoms.
- To prevent fungal and parasitic infections, keep feet clean and dry, and do not go barefoot, especially on beaches where animals may have defecated.

[Top of Page \(#top\)](#)

After You Return Home

If you are not feeling well, you should see your doctor and mention that you have recently traveled. Also tell your doctor if you were bitten or scratched by an animal while traveling.

If you have visited a malaria-risk area, continue taking your antimalarial drug for 4 weeks (doxycycline or mefloquine) or seven days (atovaquone/proguanil) after leaving the risk area.

Malaria is always a serious disease and may be a deadly illness. If you become ill with a fever or flu-like illness either while traveling in a malaria-risk area or after you return home (for up to 1 year), you should seek **immediate** medical attention and should tell the physician your travel history.

Important Note: This document is not a complete medical guide for travelers to this region. Consult with your doctor for specific information related to your needs and your medical history; recommendations may differ for pregnant women, young children, and persons who have chronic medical conditions.

[Top of Page \(#top\)](#)

***Map Disclaimer** - The boundaries and names shown and the designations used on maps do not imply the expression of any opinion whatsoever on the part of the Centers for Disease Control and Prevention concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Approximate border lines for which there may not yet be full agreement are generally marked.*

- Page last reviewed: January 18, 2011
- Page last updated: March 01, 2011
- Page created: December 22, 2008
- Content source: [Centers for Disease Control and Prevention \(http://www.cdc.gov/\)](http://www.cdc.gov/)
[National Center for Emerging and Zoonotic Infectious Diseases \(NCEZID\) \(http://www.cdc.gov/ncezid/\)](http://www.cdc.gov/ncezid/)
[Division of Global Migration and Quarantine \(DGMQ\) \(http://www.cdc.gov/ncezid/dgmg/index.html\)](http://www.cdc.gov/ncezid/dgmg/index.html)



*Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, 24 Hours/Every Day - cdcinfo@cdc.gov*