

Preventive Interventions for Preterm Children: Effectiveness and Developmental Mechanisms

Michael J. Guralnick, PhD

ABSTRACT: This article provides an integrative review of the effectiveness of and possible developmental mechanisms associated with preventive interventions for preterm children. An analysis of randomized clinical trials carried out within the last 15 years was framed within a contemporary developmental model emphasizing the role of parental adjustments to preterm children's characteristics. Evidence suggested positive outcomes could be understood in terms of improvements in developmental pathways associated with parental sensitive-responsiveness and child participation in intensive intervention-oriented child care. Implications for the critical role of the Medical Home model for preventive interventions for preterm children were discussed.

(*J Dev Behav Pediatr* 33:352–364, 2012) **Index terms:** preterm birth, intervention effectiveness, developmental mechanisms.

The adverse neurodevelopmental consequences of preterm birth remain major concerns worldwide. Despite the identification of risk factors contributing to preterm birth, preventive efforts have not been effective.¹ Moreover, increased survival of very preterm infants in particular has added substantially to this burden as the severity of the impact of preterm birth on children's social and cognitive competence increases in an almost linear fashion with decreasing gestational age.^{2–4} It is also the case that follow-up of children at low biomedical risk (30–34 wk gestational age) without apparent disabilities has revealed an increased risk for many minor but nevertheless developmentally significant problems.^{5,6} Even late preterm children (34–37 wk gestational age) manifest poorer developmental outcomes than do full-term children.^{7–9} Increased risks for sensory and motor problems create additional complexities for these children.³

The diversity of neurodevelopmental outcomes is quite remarkable but can be understood primarily in relation to variations in biomedical risk and the neurobiological mechanisms involved.^{10,11} As manifested at the behavioral level, a wide range of risks to basic developmental processes are evident including visual motor skills, visual memory, spatial processing, language, as

well as more complex and higher order organizational processes including metacognition, executive function, and motivation.^{12–18} Difficulties regulating attention have been noted as well.^{13,19} Correspondingly, numerous socioemotional and emotion regulation concerns have been observed. Early on, preterm children exhibit arousal, regulatory, organizational, and attentional difficulties that often manifest as increased irritability, reduced emotional expression, and lower levels of social initiations.^{13,20–22} Taken together, risks to these and related developmental resources and organizational processes combine to adversely influence children's emerging cognitive and social competence throughout early childhood. Indeed, from a cognitive perspective alone, on average preterm children's IQs are lower by one-half to three-quarters of a SD compared with those born full-term.^{2,23} Correspondingly, academic difficulties become apparent over time as do increased risks for a range of behavioral and social skills problems.^{24–26}

Complementing ongoing biomedical efforts to counter the potential adverse consequences of preterm birth is the wide range of behavioral and developmentally oriented postnatal interventions that have been designed and implemented in an attempt to prevent entirely or at least minimize risks to children's social and cognitive competence. Developmental research has strongly suggested that experientially based environmental influences are closely linked to a preterm child's developmental level.²⁷ When translated into preventive interventions, behavioral/developmental approaches have varied extensively in terms of rationale, timing, comprehensiveness, intensity, professional staff involved, and numerous other factors. Although some interventions have been highly focused, such as those that are oriented toward physiotherapy, the majority have had a broader but common goal, i.e., to assist parents and other caretakers to

From the Center on Human Development and Disability and the Departments of Psychology and Pediatrics, University of Washington, Seattle, WA.

Received November 2011; accepted January 2012.

This research was supported by grant P30 HD02274 from the National Institute of Child Health and Human Development.

Disclosure: The author declares no conflict of interest.

Address for reprints: Michael J. Guralnick, PhD, Center on Human Development and Disability, University of Washington, Box 357920, Seattle, WA 98195-7920; e-mail: mjgural@u.washington.edu.

Copyright © 2012 Lippincott Williams & Wilkins

adjust effectively to a preterm child's developmental and behavioral characteristics to optimize their social and cognitive competence. As will be seen, this is clearly a coregulatory process.

In this article, the possible developmental mechanisms through which preventive interventions emphasizing parental adjustments to preterm children may operate and their effectiveness are examined. This analysis is framed within a contemporary developmental model, the Developmental Systems Approach (DSA),^{28,29} and is summarized following this introduction in the first section of this article. As will be seen, the DSA is organized in terms of the risk and protective factors associated with each of 3 hierarchically arranged, but interrelated levels: (1) children's social and cognitive competence; (2) family patterns of interaction that influence children's competence; and (3) family resources that directly affect family patterns of interaction (see Fig. 1). The second section, "Preventive interventions," provides a summary of the outcomes of recent randomized clinical trials that have focused on assisting parents to

adjust to their child's characteristics. More specifically, these outcomes and their effectiveness are evaluated in terms of the developmental mechanisms described within the framework of the DSA. The implications of this analysis for the design of intervention programs for preterm children from a systems perspective are discussed in the final section.

DEVELOPMENTAL SYSTEMS APPROACH

The DSA framework is designed specifically to address issues related to children at risk for developmental delays and disabilities as well as young children with established disabilities in relation to the design and implementation of early intervention programs. This includes preventive interventions for preterm children. For ease of communication, preventive intervention will be referred to here as intervention in most instances. As illustrated in Figure 1, as children seek to accomplish goals and demonstrate their social and cognitive competence, they rely on a series of developmental resources (fundamental developmental domains of cognitive, lan-




Figure 1. The 3 levels of the Developmental Systems Approach with key components illustrating interrelationships and reciprocal influences including the effects of child-based stressors (adapted from Guralnick³⁰).

guage, motor, socioemotional, and sensory perceptual) as well as organizational processes (executive function, metacognition, social cognition, motivation, and emotion regulation).³⁰ As noted earlier, preterm children are at increased risk with respect to many developmental resources (and components of these resources such as visual memory) as well as organizational processes. Accordingly, these child-specific risks interact with protective factors at the level of child development to establish a child's level of social and cognitive competence at various points in time. For preterm children, these child-specific risk factors often exert an influence sufficient to reduce their overall levels of social and cognitive competence in relation to full-term children.²⁶

Family Patterns of Interaction

As is the case for all children, however, their competence is substantially influenced by the environmental context as primarily established by their families.^{31,32} In most instances, through an array of family patterns of interaction, families are able to adjust to their child's unique and changing developmental patterns to support child development in as optimal a manner as possible. This adjustment process is represented by the dotted arrow from the level of child social and cognitive competence to the level of family patterns of interaction in Figure 1. Three types of family patterns of interaction (parent-child transactions, family-orchestrated child experiences, and health and safety provided by the family) can be identified, and their major components are depicted in the figure. Within the DSA, these 3 components constitute the major developmental pathways directly influencing the level of child competence.

Especially during the first 3 years of the child's life, parent-child transactions are most influential and salient at this level and are emphasized in this article. These parent-child transactions are considered to be relationships necessary for promoting optimal child development that take the following 3 forms: (1) discourse framework, (2) instructional partnership, and (3) socioemotional connectedness. Each relationship creates a psychological state in which both partners (parent and child) have expectations about each others' roles in the transaction.³³ Although it is beyond the scope of this article to discuss details, these relationship processes encompass, for example, active and elaborate "conversations" (discourse framework),^{34,35} scaffolding of tasks (instructional partnership),^{36,37} and the formation of a shared or mutually responsive orientation including a secure attachment (socioemotional connectedness).^{38,39} Parent-child transactions are core features of the DSA and have received widespread conceptual and empirical support.³⁰ It is these 3 relationship processes that are thought to be the mechanisms that mediate many of the effects of family influences on children's competence especially during the child's first 3 years of life.

Most often, however, as will be seen, interventions for preterm children focus not on the 3 relationship

processes themselves but on their building blocks, i.e., sensitive and responsive interactions occurring between parents and children. Measures of parental "sensitive responsiveness" take many forms and include assessments of contingent responsiveness, affective warmth, and intrusiveness of exchanges when interacting with their child. Sensitive-responsiveness is best assessed in different contexts and family routines as well as evaluated in terms of frequency of interactions. That is, parents and children should be engaged with one another to a sufficient extent. High-quality parental sensitive-responsiveness occurring during parent-child interactions represents an awareness of their child's interests, skills, and abilities as well as their emotional and motivational state. It is suggested that the 3 relationship processes emerge over time as a result of ongoing sensitive-responsive exchanges. Moreover, these processes are clearly interrelated but yet sufficiently differentiated to produce varying effects on children's competence.⁴⁰

Stressors to Parent-Child Transactions

For many families, adjustments to create high-quality parent-child transactions are not entirely successful as a result of child-specific characteristics. Within the DSA, these child characteristics are said to constitute stressors (see Fig. 1). Available evidence suggests that, in fact, preterm children's characteristics discussed earlier substantially increase the risk that they will serve as stressors and affect numerous components of a family's pattern of interactions, especially parent-child transactions. In general, mothers' difficulties in adapting to their preterm child to support development in as sensitive and responsive a manner as possible have been well described. Indeed, problems establishing an overall synchronous relationship with their infant are evident even in the neonatal intensive care unit (NICU).^{20,41} Specific maternal behaviors of concern include increased intrusiveness, frequently redirecting their child, and failure to recognize and adjust to their child's signals, among others. These difficulties often continue through various periods of early childhood.^{13,42}

A series of studies by Landry et al²⁷ as well as by other investigators have clearly demonstrated the close association between levels of sensitive-responsiveness and numerous child developmental resources and organizational processes for preterm children. Toy play, language development, and executive function have been especially well studied.^{27,43} Moreover, associations between sensitive-responsiveness and child outcomes closely covary over time, as changes in maternal behaviors at different time points in early childhood are associated with corresponding changes in children's social and cognitive competence.⁴⁴ Of significance, various studies of these interactions indicate that the direction of influence on preterm children's development is from parent to child.^{13,42,45,46}

When stressors are extensive and sensitive-responsiveness is of correspondingly low quality, preterm chil-

dren's competence seems to be more adversely affected than that of full-term children.⁴⁷ Fortunately, higher levels of sensitive-responsiveness may also provide a special benefit for preterm children.⁴⁴ This moderating effect is illustrated in Figure 1 by the dashed line from the level of child social and cognitive competence to the level of family patterns of interaction. Accordingly, interventions that enhance parent sensitive-responsiveness to the extent that improvements in the quality of parent-child transactions occur can be expected to promote children's competence.

Family Resources

Finally, characteristics of preterm children (in terms of both their health and development) can create stressors that can also affect a family's resources (see bottom section of Fig. 1). Among them, the most common effects are unusually high levels of parental distress (a mix of anxiety and depression), especially during the first year of the child's life^{20,48-50} as well as perceptions of child vulnerability that may persist for long periods of time.⁵¹ In turn, these and other stressors affecting the level of family resources can adversely influence 1 or more components of a family's pattern of interactions, especially parent-child transactions (see Fig. 1). For example, increased maternal distress in the child's first year is associated with lower levels of sensitive-responsiveness.^{20,52} Moreover, given the co-occurrence of limited family resources (high environmental risk) and the likelihood of a preterm birth, these preexisting family resource problems are certain to also adversely influence family patterns of interaction over time. Problems associated with families at high environmental risk are often exacerbated by the stressors created by the birth of a preterm child. As discussed later, these "doubly vulnerable" children create unusually complex problems for interventions seeking to improve the quality of parent-child transactions.⁵³

In partial summary, at each of the 3 levels of the DSA (child social and cognitive competence, family patterns of interaction, and family resources), a series of components have been identified each capable of serving as a risk or protective factor for all children, including preterm children. These risk and protective factors interact with one another within each level and also exert influences between levels as illustrated in Figure 1 and described earlier. Optimal child development occurs when children consistently experience high-quality family patterns of interaction. Most families are able to make necessary adjustments to their child's characteristics to achieve sufficient levels of high-quality family patterns of interaction. However, others experience considerable problems. The consequence is to create stressors that elevate risk factors at the level of family patterns of interaction or at the level of family resources.

Especially for the first 3 years of life, a major stressor to family patterns of interaction created by preterm children's characteristics is the ability of parents to en-

gage in sensitive-responsive interactions as effectively as parents of full-term children. It is suggested that this circumstance impairs the formation of relationship processes—discourse framework, instructional partnership, and socioemotional connectedness—processes essential for supporting all children's social and cognitive competence across the early childhood period. Accordingly, successful intervention programs for preterm children will have effectively maximized parent-child transactions and other family patterns of interaction.

PREVENTIVE INTERVENTIONS

In this section, the effectiveness of recent preventive intervention programs for preterm children is reviewed from the perspective of the DSA. As suggested, interventions for preterm children should be most effective if the quality of family patterns of interaction improves, with a primary developmental mechanism being enhanced parent-child transactions. High-quality parent-child transactions that are established early in the child's life may well provide the continuity of relationships necessary to permit adjustments to children's characteristics and to minimize stressors that may emerge over time. As a consequence, a child's development will be maximized in the context of biological constraints. As noted, most intervention studies address the building blocks of those transactions, referred to as parental sensitive-responsiveness. However, these measures serve as useful indices for the 3 key relationship processes and their associations with child outcomes.

In addition to targeting parent-child transactions, another potentially important intervention approach at the level of family patterns of interaction is enrollment of their child in quality child care or preschool programs. Intervention-oriented child development programs may be especially valuable for children at risk as school readiness may be improved and enhanced child competence may contribute to better quality interactions between parents and children. It is in this context that teacher-child relationships can be formed in a manner that parallels parent-child relationships. As discussed later, this suggests the operation of similar developmental mechanisms identified by the DSA functioning with different caregivers in different contexts. This circumstance may also provide a line of continuity needed to establish longer term benefits of early childhood interventions.

Finally, these developmental mechanisms directly addressing the level of family patterns of interaction can be supplemented by interventions utilizing more indirect approaches, i.e., those focused at the level of family resources, including reducing parent distress, providing professional support, or improving parent coping skills (see Fig. 1). The expectation is that components at this level in which intervention successfully reduces risk factors will support higher quality family patterns of interaction and, as a consequence, improved child competence will result. Accordingly, a more complete understanding of the developmental mechanisms that

have been influenced or failed to have been influenced by intervention programs can contribute to a better understanding of the current status of the effectiveness of preventive intervention programs for preterm children and provide a framework for future research and practice.

Organization of This Review

With this background, randomized clinical trials carried out in the last 15 years will be evaluated. Interventions initiated at earlier points are included if warranted by follow-up studies of earlier cohorts. Studies were identified based on a comprehensive search of the literature using standard databases. Only studies that addressed outcomes related to children's social and cognitive competence and provided sufficient detail about the interventions to permit an assessment of possible developmental mechanisms were included.

The review is organized into sections defined by the timing of intervention initiation. Accordingly, the first section describes interventions carried out entirely while the child was in the NICU. The prospect of capitalizing on sensitive periods was central to the rationale for intervening while infants were in the NICU.^{41,54} The second section addresses studies that were initiated in the NICU but were continued into the home setting for various periods of time. The intent here was to facilitate the transition from the NICU to home, often by providing ongoing professional support and by assisting parents to maximize family patterns of interaction. The third section focuses exclusively on interventions that were initiated following discharge from the NICU. By having interventions begin after a period of time has elapsed since discharge from the hospital, parents may be more aware of issues and perhaps more receptive and comfortable with the intervention program. As will be seen, postdischarge interventions, in particular, varied extensively in terms of the intervention length and other characteristics. Please note that children's ages in the studies described in the 3 sections represent a correction for preterm birth.

NICU Interventions

Despite the challenging circumstances for all involved in the NICU environment, a concerted effort was made beginning in the late 1980s to assist infants to organize their behavior and to reduce stress.⁵⁵ The term "developmental care" has been used to characterize this shift in NICU practices that included structural and staffing modifications.⁵⁶ However, central to developmental care were efforts to assist parents to recognize and then adjust to their child's behavioral capacities. The primary approach was to foster sensitive and responsive exchanges to establish a foundation for the development of synchronous parent-child relationships. From the perspective of the DSA, over time these more global relationships would become more differentiated, ultimately supporting all 3 critical forms of parent-child transac-

tions (discourse framework, instructional partnership, and socioemotional connectedness). Accordingly, relationship processes constituted the primary developmental mechanisms intended to produce the hoped for long-term gains in children's social and cognitive competence for this form of intervention.

The most well-known intervention carried out within the developmental care framework is the Neonatal Individualized Developmental Care Program (NIDCAP).⁵⁵ Briefly, although there are a number of variations of this approach, well-trained developmental specialists carry out observations, including those related to the infant's autonomic, motor, and state organization as well as attentional and self-regulating patterns. This information then forms the basis for designing individualized strategies to enhance parent-child transactions.

Als and coworkers implemented this approach in a number of randomized clinical trials yielding positive short-term outcomes (2 wk postterm) for both low and high biological risk preterm children.^{54,57-59} At the level of child competence, available evidence suggests that certain developmental resources (e.g., motor responses) and even the rudiments of organizational processes (e.g., emotion regulation and attentional mechanisms) can be enhanced by the intervention.⁵⁷ These more organized behavioral patterns may well permit infants to attend to the environment and process information more effectively. The result is improved competence, at least in the short term, and perhaps even placing some children on a trajectory that can support longer term effects. Indeed, NIDCAP continued to have a positive effect on children's cognitive development at 9 months.^{54,58} In this context, it should be noted that, although not a randomized trial, rigorously carried out work emphasizing one of the components of the multicomponent NIDCAP intervention, skin-to-skin contact with the mother generally referred to as the Kangaroo Care approach,⁶⁰ has also produced positive sustained effects in comparison with a group for whom this approach was not part of the NICU's standard protocol. Specifically, at 6 months of age, preterm children receiving Kangaroo Care achieved higher scores on cognitive measures and measures of attention and exploration.^{21,61} Improved parental sensitive-responsiveness was associated with positive child outcomes.⁶¹

Alternative or complementary developmental pathways which may have produced these effects are influences that operate at the level of family resources. Indeed, all of the various forms of developmental care provided professional and other forms of social supports to parents while their child was in the NICU. To the extent that this supportive relationship occurred, it can be expected to influence a number of possible components at the level of family resources.⁶²⁻⁶⁴ In fact, available evidence indicates that developmental care interventions do reduce many risk factors associated with family resources. Specifically, as a result of these interventions, parents perceive their child more positively,

experience less overall distress, and consider themselves to be more competent parents.^{57,61,65} As noted earlier, many of these family resource components are at increased risk as a result of preterm birth. Reductions in these risk factors can be expected to have a positive influence on parent-child transactions thereby augmenting any effects of the intervention focusing more directly on these relationships.

Despite these promising results, other investigators have only partially replicated the NICDAP findings, producing complex outcomes and often failing to find sustained effects.⁶⁶⁻⁷⁰ Compounding these inconsistencies are the many methodological problems that have been identified.^{56,71} At present, it can be concluded that interventions in the NICU may well have the potential to assist families to develop more synchronous relationships with their child in the NICU through enhanced sensitive-responsiveness and improved family resources related to reduced parent distress and more confident parenting. However, firm conclusions must await the results of well-designed studies focusing on these particular developmental mechanisms and their impact on children's competence.

Combined NICU and Home Interventions

The transition to home provides a more comfortable and familiar setting for families but also brings about an entirely new set of responsibilities. Moreover, the potential clearly exists for different child-specific risk factors to emerge or now exert a stronger influence on family patterns of interaction and family resources. To ease this transition and to try to maintain any positive benefits that may have occurred in the NICU, a number of interventions have been carried out with the idea of helping families adapt to these new circumstances.

Many contemporary studies followed the approach of an earlier intervention that produced unusually promising effects. Referred to as the Mother-Infant Transaction Program (MITP), this intervention was modeled closely after NICDAP and provided 7 sessions in the NICU during the week before discharge.⁷² Mothers were assisted in identifying child cues to distress and provided with techniques to support their child's self-regulation. Improving parental sensitive-responsiveness was again at the center of this intervention. The 4 home visits that followed discharge from the hospital were designed to provide professional support (e.g., caretaking advice and information on child temperament) and to enhance parental confidence while continuing to encourage effective parent-child exchanges. A major influence on child competence for those participating in the MITP resulted from this modest, 11-session intervention. Specifically, the cognitive development of control group children (and many aspects relevant to their social development) declined over time, whereas the intervention group remained stable and eventually became comparable with a full-term group.^{73,74}

A reasonable interpretation of these findings, and consistent with other measures obtained in this long-term

longitudinal study, is that the MITP intervention provided families, most of whom were not at high environmental risk, with the skills and confidence to continue to adjust parent-child transactions and ultimately other aspects of family patterns of interaction to changing child characteristics over time. That is, stressors to optimal family patterns of interaction were minimized. Of note, other early studies similar to the MITP involving high environmental risk families produced far more modest effects despite an extended home-based component of 1 to 2 years.^{75,76} The ongoing cognitive declines for both intervention and control groups for these high environmental risk families, despite less of a decline for intervention children, emphasize the powerful role of limited family resources including their influence on virtually all components of family patterns of interaction.^{53,77} Again, however, better outcomes for preterm children were associated with higher scores on measures related to sensitive-responsiveness.

The results of contemporary studies using the MITP protocol or variations of this intervention have not been nearly as promising.⁷⁸⁻⁸¹ Either no effects or minor effects on children's cognitive development have been found, despite findings of improved parent sensitive-responsiveness. In fairness, however, children have not been followed for long periods of time. This makes it difficult to evaluate the ultimate effects of these modest interventions as any influences of sensitive-responsiveness that might exist are often not apparent until later periods during early childhood development. Moreover, for both high and low environmental risk samples, those families participating in the interventions were not only found to display higher levels of sensitive-responsiveness but also experienced less child-related stress and considered their children to be less challenging and temperamentally easier.⁸¹⁻⁸⁵ These findings were not consistent across studies but do allow the possibility that benefits to children may arise at later points in time. Although all these studies were based on the MITP, variations in the MITP protocols emphasized by different groups of investigators, sample differences, and the varied training levels and types of professionals implementing the interventions (e.g., nurses, occupational therapists, and physical therapists) make it difficult to find any meaningful patterns in the outcomes for combined NICU and home interventions.

Interventions Initiated in the Home

As discussed later, similar and perhaps even more disappointing outcomes were obtained from a series of studies that emphasized beginning intervention in the home after allowing time for more stable family routines to be established and giving parents an opportunity to identify any child-specific issues that were of concern, especially for interventions that began a few months postdischarge. These interventions initiated in the home were generally of modest intensity, typically consisting of 1- to 1½-hour home visits monthly or twice monthly

for periods of 6 months to as long as 2 years. They also tended to have substantial didactic or educational skills features. Nevertheless, the intervention curricula were quite diverse but sought to build a relationship between home visitors and parents, provide professional support, and make referrals to other service providers as needed. Supportive efforts to promote high-quality parent-child transactions through strategies to improve sensitive-responsiveness were also central to most of the home-initiated interventions. Despite many variations of this general approach, the most consistent result was an absence of effects on children's competence.⁸⁶⁻⁸⁸ Not only did these studies fail to find short-term effects but also longer term follow-ups were equally disappointing.⁸⁹ Unfortunately, limited information with respect to changes in sensitive-responsiveness was obtained, but it is likely that minimal effects occurred.⁸⁸

Playing and Learning Strategies Intervention

The sole exception to the disappointing outcomes of these modest home visiting approaches was the intervention referred to as Playing and Learning Strategies (PALS).²⁷ In contrast to the work noted earlier, this intervention incorporated more contemporary approaches supporting sensitive-responsiveness and related developmental mechanisms and provided extensive information with respect to the possible relationships existing among all 3 levels of the DSA. Accordingly, this intervention is considered in detail.

Initially designed to provide only ten 1½-hour home visits when the child was between 6 and 10 months of age, the intervention was supplemented by an additional 11-week home visiting program when the child was approximately 30 months of age. The detailed curriculum was designed to promote 4 components representing sensitive-responsiveness: (1) contingent responding, (2) emotional affective support, (3) supporting the infant to focus attention (maintaining), and (4) language support consistent with the child's needs. Of importance, this intervention was based on years of careful developmental research identifying the components of sensitive-responsiveness associated with optimal child outcomes most likely to have an impact on the 3 DSA-based relationship processes.

Highly interactive sessions including strategic use of videotapes and alternate caregivers along with techniques to integrate interactive behaviors into everyday routines together contributed to create a highly sophisticated intervention approach. A special challenge for the intervention program was the fact that the children studied were doubly vulnerable, as most families were at high environmental risk. Despite these circumstances, measures obtained from observations of mother-child interactions, independent child play, and interactions with an examiner at different points in time following the end of intervention that began during the child's first year yielded highly encouraging results. Among the positive findings, measures related to child language and cognitive development were higher for children in the

intervention group when compared with a control group assessed 3 months following completion of the initial 10 sessions.⁹⁰ Changes in children's competence were accompanied by corresponding changes in parent sensitive-responsiveness. Of note, sensitive-responsiveness for control group mothers declined over time, likely reflecting the influence of child-specific stressors on parent-child transactions and possibly contributing to elevating risk factors related to other components of family patterns of interaction or components at the level of family resources. Moreover, the cumulative effect of preexisting limited family resources for this high environmental risk sample also likely contributed to the observed decline in sensitive-responsiveness for control families. That the PALS intervention, acting directly at the level of parent-child transactions, was able to overcome these risk factors attests to the quality of this intervention.⁹¹ Adding intervention components at the level of family resources to provide professional supports and assist in accessing community services would be expected to have further positive effects on child competence. Preliminary findings suggest that this is precisely what occurs.⁹²

The second phase of the PALS intervention allowed both a follow-up of children receiving the initial 10 intervention sessions as well as an evaluation of any additional benefits of the second intervention implemented during the preschool period. In this second phase, an 11-week intervention followed the same approach to increasing mother's sensitive-responsiveness but was adjusted to children's characteristics at their current developmental level. Findings were complex, but nevertheless revealed that the second phase made an important additional contribution to key aspects of sensitive-responsiveness, particularly contingent responding and verbal engagement.⁹³ Moreover, child competencies with respect to language development and social engagement were more optimal when families participated in both PALS interventions. Contingent responding was especially important and reflected the ability of mothers to adapt appropriately to their child's changing characteristics. Although other factors such as professional support may have contributed to these outcomes, further analysis revealed that the contingent responding and warm sensitivity components of sensitive-responsiveness were important mediators of the intervention effects on preterm children's competence.

Long-term effects of the PALS intervention have not been examined, but findings for children from both phases of an intervention in the early childhood period that primarily focused on and benefited various components of sensitive-responsiveness are encouraging. It is unclear, however, the extent to which this modest intervention was able to substantially alter any of the 3 relationship processes central to the DSA. Unless these were to occur, only short-term effects would be expected. The added benefits of the second phase during the preschool years did increase prospects for establish-

ing higher quality relationships in the form of parent-child transactions. Accordingly, this suggests the need for some form of periodic assessment process and corresponding interventions as needed over time to maximize long-term outcomes. As discussed earlier, child-specific stressors may emerge over time as preterm children encounter more complex and demanding developmental tasks that now tax developmental resources and organizational processes that are at higher risk.

Infant Health and Development Program

The Infant Health and Development Program (IHDP) is a landmark preventive intervention effort for preterm children which can be characterized by its comprehensiveness, high intensity, and multisite features. The high intensity of the intervention with its many components distinguished it from other intervention programs and is therefore also considered in more detail. Initial results of this well-designed study were reported in 1990,⁹⁴ but many of the nearly 1000 children and their families have been followed for close to 2 decades. Three major intervention components were implemented for a 3-year period beginning soon after discharge from the hospital nursery. Although details have been well described elsewhere,⁹⁵ a central feature of the intervention was a home visiting component in which a home visitor assisted mothers to develop better problem-solving skills related to everyday problems as well as with respect to the care of their preterm child. Parent group meetings were also organized to provide an additional form of social support. In addition, a major focus of the home visiting program was the implementation of 2 formal educational/developmental curricula. The first was implemented soon after the infant was discharged from the hospital and was similar to most other curricula designed to assist mothers to recognize children's cues and to help them self-regulate. The second was adopted from a curriculum for full-term children whose families were at high environmental risk focused on promoting advances in the major developmental domains guided by each child's developmental pattern. Activities were carried out to maximize sensitive-responsiveness and affective warmth and there was a strong "educational skills" emphasis to the curriculum. This same curriculum was implemented by well-trained educational staff for children enrolled in an intervention-oriented child care center operated by the researchers. Families were encouraged to enroll their child in the center during years 2 and 3 of the intervention. Both intervention and control groups received regular pediatric follow-up care, related assessments, and referrals as needed to community specialists. Numerous measures were obtained at various points during and following completion of the intervention.

At the end of the 3-year period, highly positive effects of the intervention were found. Focusing on overall cognitive development, both intervention and control groups showed declines over time, but much less so for children participating in the intervention. In general, these effects were more pronounced for children at

lower biological risk based on birth weight and for mothers at higher environmental risk.⁹⁶ Follow-up of children at later ages revealed some residual positive effects varying with birth weight, but the major differences between the groups were no longer apparent.⁹⁷⁻⁹⁹

Additional analyses were carried out in an effort to identify likely developmental mechanisms and perhaps provide an understanding as to the pattern of short- and long-term results. With respect to short-term benefits, the IHDP intervention appeared to have only minor effects on components assessed at the level of family resources. Specifically, less emotional distress was reported by mothers in the intervention group, but maternal distress did not seem to mediate child outcomes nor were any effects found for maternal coping strategies.^{96,100} Focusing on the level of family patterns of interaction, in view of the home visiting component positive short-term effects may well have been due to direct changes in sensitive-responsiveness, as appeared to be the case for the PALS intervention. Some differences between IHDP intervention and control groups were, in fact, found for relevant measures but seemed to revolve entirely around instructional issues designed to promote their child's development. Specifically, intervention group mothers were observed to provide higher quality assistance in a problem-solving task with their child, although the effects were quite small.¹⁰¹ They also provided more appropriate and stimulating learning materials at home.⁵³ However, other measures relevant to parent-child transactions including a general assessment of sensitive-responsiveness as well as language stimulation did not differ between the groups.⁵³ In addition, no long-term effects of parental style or the provision of developmentally supportive activities in the home were found, but opportunities for earlier employment due to the availability of the child care center may have had a positive effect.¹⁰²

Although the cumulative impact of this comprehensive intervention must be considered, the pattern of findings suggested that the experiences of the child in the child development center were most likely responsible for the between-group improvements in children's competence. Both short-term and the longer term outcomes were closely associated with participation in the center and engagement with the curriculum.^{103,104} Of note, mothers at higher environmental risk for whom the program was most effective seemed to have less of an interest in the educational materials provided by the home visitor component.¹⁰⁵ Accordingly, particularly in instances in which engagement with teachers was high in the child care center, the quality of teacher-child relationships formed in the child development center seem to have created conditions for improving children's social and cognitive competence. Intensive, high-quality child care or preschool programs that foster teacher-child relationships especially for children in high environmental risk families are much more likely to

develop a discourse framework, an instructional partnership, and socioemotional connectedness.^{106,107}

It is clear that many parents of preterm children, particularly those at low environmental risk, were able to adapt to any emerging child-specific risk factors. Family resources such as the ability to cope and utilize their social support networks, including professional supports, were likely among the characteristics of these families which ensured that high-quality family patterns of interaction were provided. The IHDP was apparently not able to enhance this pattern. Moreover, perhaps as a result of the educational skills focus of the intervention, the IHDP did not have major effects on parent-child transactions or family resources, irrespective of environmental risk level. As a consequence, following termination of the intervention, when the child was 3 years of age, many high environmental risk parents likely experienced additional difficulties adapting to their child's changing developmental patterns especially when children encountered more demanding but less supportive situations such as those occurring during preschool or kindergarten programs. The advances that were achieved through participation in the child development center, perhaps including expectations for forming quality relationships with teachers, may have had some long-term benefits.¹⁰⁵ However, for the most part, in the absence of substantial changes in parent-child transactions as reflected by minimal changes in sensitive-responsiveness, any emerging child-specific risk factors (e.g., negative emotionality)¹⁰⁸ were likely to constitute stressors that persisted. These stressors would be compounded further by the many (preexisting) family resource problems that inevitably occurred over time for high-risk families thereby increasing risk levels for the many components of family patterns of interaction and limiting long-term effects.

CONCLUSIONS AND IMPLICATIONS FOR PRACTICE

Despite important medical advances, preterm birth remains a major concern with significant consequences for children's development. Preventive intervention programs have addressed the challenges facing families to assist them to adapt to their child's characteristics and to establish as optimal a developmental environment as possible. These interventions have been initiated at various points throughout the early childhood period and have differed substantially not only in terms of timing but also in duration, approach, intensity, comprehensiveness, and other dimensions. In all instances, however, the expectation was that interventions occurring during the early childhood period will not only produce immediate, short-term benefits but also establish conditions that will create sustained effects over time.

The diverse characteristics of the available studies as well as the complexity and often inconsistent results that have been found do not allow straightforward conclusions as to effectiveness.¹⁰⁹ Nevertheless, sufficient in-

formation is available to suggest that interventions implemented at any point in time during the early childhood period can produce modest short-term effects on children's competence. Positive findings may be related to indirect effects of reduced parental distress, an enhanced support system, or other factors at the level of family resources that can influence parent-child transactions in particular. However, only limited support exists for this developmental pathway. With respect to a different pathway, especially for high environmental risk families, extensive participation in an intervention-oriented child development center can make an important contribution to children's competence at least during the time it is in effect. However, it has been difficult to demonstrate widespread sustained effects even for the high-quality and highly intensive intervention provided by IHDP.

Alternatively, important positive findings have been accompanied by improvements to many components of parental sensitive-responsiveness. When positive effects do occur, they are likely the result of *direct intervention* efforts to improve parents' ability to adapt to their child's characteristics. It is important to emphasize that sensitive-responsiveness is hypothesized to be of significance because it serves as the basis for establishing high-quality relationships (i.e., discourse framework, instructional partnership, and socioemotional connectedness). As suggested by the DSA, it is these parent-child transaction processes that provide the continuity necessary to maintain an optimal developmental environment for the child. Unfortunately, only limited evidence indicates that interventions were sufficient to substantially alter these relationship processes. Despite shorter term effects produced by increases in sensitive-responsiveness, the consequence of this is an absence of continuity of parent-child transactions needed to sustain longer term child social and cognitive competence. Strengthening these relationships constitutes a critical task for future research and practice.

Clearly, so many diverse influences that can affect preterm children's development are beyond our control or current understanding. As discussed, many effects diminish substantially soon after the intervention is complete, often failing to be sustained even during later points in the early childhood period itself. Of note, the evidence for sustained long-term effects may be minimal, but most studies have not carried out the necessary follow-up work. In addition, high levels of variability are common, and many children not receiving intervention can manifest accelerated developmental patterns under favorable circumstances.^{99,110,111} This further diminishes any intervention effects over time. Nevertheless, quality relationships formed during the early childhood period may well provide the level of continuity sufficient to offer at least some protection from the challenges that lie ahead.

Preventive Interventions and the Medical Home

Taken together, this analysis suggests that, to be successful, preventive intervention programs may well require a systems perspective that extends intervention activities across the entire early childhood period. This integrative review has emphasized the centrality of parent-child transactions, but other components noted in Figure 1 must be part of the overall system. Such a system must ensure consistency and continuity over time as well as the ability to integrate and coordinate all the various factors that might be involved. Within the DSA framework, this means organizing a system that is capable of addressing risk and protective factors at all 3 levels: (1) level of child development, (2) level of family patterns of interaction, and (3) level of family resources.

How to accomplish this from a practical perspective is, of course, extraordinarily challenging as systems approaches require levels of coordination, integration, and continuity seldom found in communities. However, the Medical Home is one model that should be considered as a framework for constructing such a comprehensive system. Characteristics of a successful Medical Home model are that care be accessible, family oriented, continuous, comprehensive, coordinated, compassionate, and culturally effective.¹¹² This model is clearly compatible with preventive intervention programs for preterm children that are also likely to be most successful. Specifically, at the level of child development, the follow-up care practices for preterm children for developmental testing¹¹³ and the American Academy of Pediatrics' algorithm for developmental surveillance and screening of young children in the Medical Home¹¹⁴ provide essential guidance. At the next level of the DSA, by developing a family's trust and gaining their confidence, considerable information regarding many components of a family's pattern of interaction can be obtained. Eliciting parent concerns about interactions with their child focusing on relationships or their child's participation in community activities can be incorporated into the components of developmental surveillance.¹¹⁴ Close working relationships with educational programs for children who qualify for special services, with child care or preschool program personnel, or with other community service agencies, can generate additional information with respect to risk and protective factors at the level of family patterns of interaction. These community resources would also be engaged as part of the intervention process. Despite some existing tools,²⁹ feasible measures of the various relationship processes and other components of family patterns of interaction (see Fig. 1) remain to be developed. This constitutes an important future research effort. Nevertheless, the DSA can serve as a common framework for all resources that are involved thereby generating more continuity and intensive efforts to promote quality family patterns of interaction. To help address issues resulting from the limited resources available in many pediatric practices, other community

programs can share or assume greater responsibility for coordinating interventions within this framework at various points in the child's development. Proper coordination may enable a cost-effective and developmentally effective system to emerge. Finally, surveillance and screening within the Medical Home has also been recently recommended for many components at the level of family resources.¹¹⁵ This is certainly not common practice today, but psychosocial screening tools for families are available in many domains including mental health, physical health, substance abuse, and social support. As is the case when child-specific problems are identified, referral to community resources will be necessary. Without question, there are clearly many barriers including time and resources to implementing a system of preventive interventions for preterm children, but for many of these children optimal child development is not likely to occur in its absence.

ACKNOWLEDGMENT

I thank Dr. Curt Bennett for comments on an earlier version of this article.

REFERENCES

1. Simmons LE, Rubens CE, Darmstadt GL, Gravett MG. Preventing preterm birth and neonatal mortality: exploring the epidemiology, causes, and interventions. *Semin Perinatol*. 2010;34:408-415.
2. Aylward GP. Cognitive and neuropsychological outcomes: more than IQ scores. *Ment Retard Dev Disabil Res Rev*. 2002;8:234-240.
3. Behrman RE, Butler AS, eds. *Preterm Birth: Causes, Consequences, and Prevention*. Washington, DC: National Academies Press; 2007.
4. Stromme P, Hagberg G. Aetiology in severe and mild mental retardation: a population-based study of Norwegian children. *Dev Med Child Neurol*. 2000;42:76-86.
5. Caravale B, Tozzi C, Albino G, Vicari S. Cognitive development in low risk preterm infants at 3-4 years of life. *Arch Dis Child Fetal Neonatal Ed*. 2005;90:F474-F479.
6. Lindstrom K, Lindblad F, Hjern A. Preterm birth and attention-deficit/hyperactivity disorder in schoolchildren. *Pediatrics*. 2011;127:858-865.
7. McGowan JE, Alderdice FA, Holmes VA, Johnston L. Early childhood development of late-preterm infants: a systematic review. *Pediatrics*. 2011;127:1111-1124.
8. Petrini JR, Dias T, McCormick MC, Massolo ML, Green NS, Escobar GJ. Increased risk of adverse neurological development for late preterm infants. *J Pediatr*. 2009;154:169-176.
9. Woythaler MA, McCormick MC, Smith VC. Late preterm infants have worse 24-month neurodevelopmental outcomes than term infants. *Pediatrics*. 2011;127:e622-e629.
10. Luciana M. Cognitive development in children born preterm: implications for theories of brain plasticity following early injury. *Dev Psychopathol*. 2003;15:1017-1047.
11. Volpe JJ. Brain injury in premature infants: a complex amalgam of destructive and developmental disturbances. *Lancet Neurol*. 2009;8:110-124.
12. Anderson PJ, Doyle LW; Victorian Infant Collaborative Study Group. Executive functioning in school-aged children who were born very preterm or with extremely low birth weight in the 1990s. *Pediatrics*. 2004;114:50-57.

13. Clark CA, Woodward LJ, Horwood LJ, Moor S. Development of emotional and behavioral regulation in children born extremely preterm and very preterm: biological and social influences. *C Child Dev*. 2008;79:1444-1462.
14. Edgin JO, Inder TE, Anderson PJ, Hood KM, Clark CAC, Woodward LJ. Executive functioning in preschool children born very preterm: relationship with early white matter pathology. *J Int Neuropsychol Soc*. 2008;14:90-101.
15. Landry SH, Smith KE, Swank PR. Environmental effects on language development in normal and high-risk child populations. *Semin Pediatr Neurol*. 2002;9:192-200.
16. Marlow N, Hennessy EM, Bracewell MA, Wolke D; EPICure Study Group. Motor and executive function at 6 years of age after extremely preterm birth. *Pediatrics*. 2007;120:793-804.
17. Roberts G, Lim J, Doyle LW, Anderson PJ. High rates of school readiness difficulties at 5 years of age in very preterm infants compared with term controls. *J Dev Behav Pediatr*. 2011;32:117-124.
18. Rose SA, Feldman JF, Jankowski JJ. Information processing in toddlers: continuity from infancy and persistence of preterm deficits. *Intelligence*. 2009;37:311-320.
19. Landry SH, Chapieski ML. Joint attention and infant toy exploration: effects of Down syndrome and prematurity. *C Child Dev*. 1989;60:103-118.
20. Feldman R. Maternal versus child risk and the development of parent-child and family relationships in five high-risk populations. *Dev Psychopathol*. 2007;19:293-312.
21. Feldman R, Weller A, Sirota L, Eidelman AI. Skin-to-skin contact (Kangaroo care) promotes self-regulation in premature infants: sleep-wake cyclicality, arousal modulation, and sustained exploration. *Dev Psychol*. 2002;38:194-207.
22. Goldberg S, DiVitto B. Parenting children born preterm. In: Bornstein MH, ed. *Handbook of Parenting, Vol 1: Children and Parenting*. 2nd ed. Mahwah, NJ: Lawrence Erlbaum; 2002: 329-354.
23. Aylward G. Neurodevelopmental outcomes of infants born prematurely. *J Dev Behav Pediatr*. 2005;26:427-440.
24. Hack M, Taylor HG, Schluchter M, Andreias L, Drotar D, Klein N. Behavioral outcomes of extremely low birth weight children at age 8 years. *J Dev Behav Pediatr*. 2009;30:122-130.
25. Hebert-Myers H, Guttentag CL, Swank PR, Smith KE, Landry SH. The importance of language, social, and behavioral skills across early and later childhood as predictors of social competence with peers. *Appl Dev Sci*. 2006;10:174-187.
26. Stephens BE, Vohr BR. Neurodevelopmental outcome of the premature infant. *Pediatr Clin North Am*. 2009;56:631-646.
27. Landry SH, Taylor HB, Guttentag C, Smith KE. Responsive parenting: closing the learning gap for children with early developmental problems. In: Glidden LM, ed. *International Review of Research in Mental Retardation*. Vol 36. St. Mary's City, MD: Elsevier; 2008:27-60.
28. Guralnick MJ. A developmental systems model for early intervention. *Infants Young Child*. 2001;14:1-18.
29. Guralnick MJ, ed. *The Developmental Systems Approach to Early Intervention*. Baltimore, MD: Brookes; 2005.
30. Guralnick MJ. Why early intervention works: a systems perspective. *Infants Young Child*. 2011;24:6-28.
31. Bradley RH, Corwyn RF. "Family process" investments that matter for child well-being. In: Kalil A, DeLeire T, eds. *Family Investment in Children's Potential: Resources and Parenting Behaviours That Promote Success*. Mahwah, NJ: Lawrence Erlbaum; 2004:1-32.
32. Collins WA, Maccoby EE, Steinberg L, Hetherington EM, Bornstein MH. Contemporary research on parenting. The case for nature and nurture. *Am Psychol*. 2000;55:218-232.
33. Tomasello M, Carpenter M. Shared intentionality. *Dev Sci*. 2007;10:121-125.
34. Chouinard MM. Children's questions: a mechanism for cognitive development. *Monogr Soc Res Child Dev*. 2007;72:1-112; discussion 113-126.
35. Thompson RA. The development of the person: social understanding, relationships, conscience, self. In: Eisenberg N, Damon W, Lerner RM, eds. *Handbook of Child Psychology: Vol. 3. Social, Emotional, and Personality Development*. 6th ed. Hoboken, NJ: John Wiley & Sons Inc; 2006:24-98.
36. Pratt MW, Kerig P, Cowan PA, Cowan CP. Mothers and fathers teaching 3-year-olds: authoritative parenting and adult scaffolding of young children's learning. *Dev Psychol*. 1988;24: 832-839.
37. Vygotsky LS. *Mind in Society*. Cambridge, MA: Harvard University Press; 1978.
38. Kochanska G. Mutually responsive orientation between mothers and their young children: a context for the early development of conscience. *Curr Dir Psychol Sci*. 2002;11: 191-195.
39. Leerkes EM, Nayena Blankson A, O'Brien M. Differential effects of maternal sensitivity to infant distress and nondistress on social-emotional functioning. *C Child Dev*. 2009;80:762-775.
40. Grusec JE, Davidov M. Integrating different perspectives on socialization theory and research: a domain-specific approach. *C Child Dev*. 2010;81:687-709.
41. Feldman R. Parent-infant synchrony and the construction of shared timing; physiological precursors, developmental outcomes, and risk conditions. *J Child Psychol Psychiatry*. 2007;48:329-354.
42. Forcada-Guex M, Pierrehumbert B, Borghini A, Moessinger A, Muller-Nix C. Early dyadic patterns of mother-infant interactions and outcomes of prematurity at 18 months. *Pediatrics*. 2006;118:e107-e114.
43. Landry SH, Smith KE. Early social and cognitive precursors and parental support for self-regulation and executive function: relations from early childhood into adolescence. In: Sokol BW, Muller U, Carpendale J, Young A, Iarocci G, eds. *Self and Social Regulation: Social Interaction and the Development of Social Understanding and Executive Functions*. New York: Oxford University Press; 2010:386-417.
44. Landry SH, Smith KE, Miller-Loncar CL, Swank PR. The relation of change in maternal interactive styles to the developing social competence of full-term and preterm children. *C Child Dev*. 1998;69:105-123.
45. Feldman R, Eidelman AI. Neonatal state organization, neuromaturation, mother-infant interaction, and cognitive development in small-for-gestational-age premature infants. *Pediatrics*. 2006;118:e869-e878.
46. Landry SH, Smith KE, Swank PR, Miller-Loncar CL. Early maternal and child influences on children's later independent cognitive and social functioning. *C Child Dev*. 2000;71:358-375.
47. Landry SH, Smith KE, Swank PR, Assel MA, Velle S. Does early responsive parenting have a special importance for children's development or is consistency across early childhood necessary? *Dev Psychol*. 2001;37:387-403.
48. Davis L, Edwards H, Mohay H, Wollin J. The impact of very premature birth on the psychological health of mothers. *Ear Hum Dev*. 2003;73:61-70.
49. Eisengart SP, Singer LT, Fulton S, Baley JE. Coping and psychological distress in mothers of very low birth weight young children. *Parent Sci Pract*. 2003;3:49-72.
50. Singer LT, Salvator A, Guo S, Collin M, Lilien L, Baley J. Maternal psychological distress and parenting stress after the birth of a very low-birth-weight infant. *JAMA*. 1999;281:799-805.
51. Miles MS, Holditch-Davis D. Compensatory parenting: how mothers describe parenting their 3-year-old, prematurely born children. *J Pediatr Nurs*. 1995;10:243-253.

52. Muller-Nix C, Forcada-Guex M, Pierrehumbert B, Jaunin L, Borghini A, Ansermet F. Prematurity, maternal stress and mother-child interactions. *Early Hum Dev.* 2004;79:145-158.
53. Bradley RH, Whiteside L, Mundfrom DJ, Casey PH, Kelleher KJ, Pope SK. Early indications of resilience and their relation to experiences in the home environments of low birthweight, premature children living in poverty. *Child Dev.* 1994;65:346-360.
54. Als H, Duffy FH, McAnulty GB, et al. Early experience alters brain function and structure. *Pediatrics.* 2004;113:846-857.
55. Als H. Earliest intervention for preterm infants in the newborn intensive care unit. In: Guralnick MJ, ed. *The Effectiveness of Early Intervention.* Baltimore, MD: Paul H. Brookes; 1997:47-76.
56. Symington AJ, Pinelli J. Developmental care for promoting development and preventing morbidity in preterm infants. *Cochrane Database Syst Rev.* 2006;(2):CD001814.
57. Als H, Gilkerson L, Duffy FH, et al. A three-center, randomized, controlled trial of individualized developmental care for very low birth weight preterm infants: medical, neurodevelopmental, parenting, and caregiving effects. *J Dev Behav Pediatr.* 2003;24:399-408.
58. Als H, Lawhon G, Duffy FH, McAnulty GB, Gibes-Grossman R, Blickman JG. Individualized developmental care for the very low-birth-weight preterm infant. Medical and neurofunctional effects. *JAMA.* 1994;272:853-858.
59. Buehler DM, Als H, Duffy FH, McAnulty GB, Liederman J. Effectiveness of individualized developmental care for low-risk preterm infants: behavioral and electrophysiologic evidence. *Pediatrics.* 1995;96(Pt 1):923-932.
60. Charpak N, Ruiz-Pelaez JG, Figueroa de CZ, Charpak Y. A randomized, controlled trial of kangaroo mother care: results of follow-up at 1 year of corrected age. *Pediatrics.* 2001;108:1072-1079.
61. Feldman R, Eidelman AI, Sirota L, Weller A. Comparison of skin-to-skin (kangaroo) and traditional care: parenting outcomes and preterm infant development. *Pediatrics.* 2002;110(Pt 1):16-26.
62. Crnic K, Stormshak E. The effectiveness of providing social support for families of children at risk. In: Guralnick MJ, ed. *The Effectiveness of Early Intervention.* Baltimore, MD: Paul H. Brookes; 1997:209-225.
63. Thompson RA, Flood MF, Goodvin R. Social support and developmental psychopathology. In: Cicchetti D, Cohen D, eds. *Developmental Psychopathology: Vol. III. Risk, Disorder, and Adaptation.* New York, NY: Wiley; 2006:1-37.
64. Zelkowitz P, Feeley N, Shrier I, et al. The Cues and Care randomized controlled trial of a neonatal intensive care unit intervention: effects on maternal psychological distress and mother-infant interaction. *J Dev Behav Pediatr.* 2011;32:591-599.
65. Melnyk BM, Alpert-Gillis L, Feinstein NF, et al. Improving cognitive development of low-birth-weight premature infants with the COPE program: a pilot study of the benefit of early NICU intervention with mothers. *Res Nurs Healthb.* 2001;24:373-389.
66. Kleberg A, Westrup B, Stjernqvist K. Developmental outcome, child behaviour and mother-child interaction at 3 years of age following Newborn Individualized Developmental Care and Intervention Program (NIDCAP) intervention. *Early Hum Dev.* 2000;60:123-135.
67. Kleberg A, Westrup B, Stjernqvist K, Lagercrantz H. Indications of improved cognitive development at one year of age among infants born very prematurely who received care based on the Newborn Individualized Developmental Care and Assessment Program (NIDCAP). *Early Hum Dev.* 2002;68:83-91.
68. Peters KL, Rosychuk RJ, Henderson L, Cote JJ, McPherson C, Tyebkhan JM. Improvement of short- and long-term outcomes for very low birth weight infants: Edmonton NIDCAP trial. *Pediatrics.* 2009;124:1009-1020.
69. van der Pal S, Maguire CM, Le Cessie S, et al. Parental stress and child behavior and temperament in the first year after the parental Newborn Individualized Developmental Care and Assessment Program. *J Early Interv.* 2008;30:102-115.
70. Westrup B, Böhm B, Lagercrantz H, Stjernqvist K. Preschool outcome in children born very prematurely and cared for according to the Newborn Individualized Developmental Care and Assessment Program (NIDCAP). *Acta Paediatr.* 2004;93:498-507.
71. Jacobs SE, Sokol J, Ohlsson A. The Newborn Individualized Developmental Care and Assessment Program is not supported by meta-analyses of the data. *J Pediatr.* 2002;140:699-706.
72. Rauh VA, Achenbach TM, Nurcombe B, Howell CT, Teti DM. Minimizing adverse effects of low birthweight: four-year results of an early intervention program. *Child Dev.* 1988;59:544-553.
73. Achenbach TM, Howell CT, Aoki MF, Rauh VA. Nine-year outcome of the Vermont intervention program for low birth weight infants. *Pediatrics.* 1993;91:45-55.
74. Achenbach TM, Phares V, Howell CT, Rauh VA, Nurcombe B. Seven-year outcome of the Vermont Intervention Program for Low-Birthweight Infants. *Child Dev.* 1990;61:1672-1681.
75. Resnick MB, Armstrong S, Carter RL. Developmental intervention program for high-risk premature infants: effects on development and parent-infant interactions. *J Dev Behav Pediatr.* 1988;9:73-78.
76. Resnick MB, Eyer FD, Nelson RM, Eitzman DV, Bucciarelli RL. Developmental intervention for low birth weight infants: improved early development outcome. *Pediatrics.* 1987;80:68-74.
77. Evans GW. The environment of childhood poverty. *Am Psychol.* 2004;59:77-92.
78. Glazebrook C, Marlow N, Israel C, et al. Randomised trial of a parenting intervention during neonatal intensive care. *Arch Dis Child Fetal Neonatal Ed.* 2007;92:F438-F443.
79. Johnson S, Whitelaw A, Glazebrook C, et al. Randomized trial of a parenting intervention for very preterm infants: outcome at 2 years. *J Pediatr.* 2009;155:488-494.
80. Koldewijn K, Wolf MJ, van Wassenaer A, et al. The Infant Behavioral Assessment and Intervention Program for very low birth weight infants at 6 months corrected age. *J Pediatr.* 2009;154:33-38. e2.
81. Teti DM, Black MM, Viscardi R, et al. Intervention with African American premature infants. Four-month results of an early intervention program. *J Early Interv.* 2009;31:146-166.
82. Kaarsen PI, Rønning JA, Ulvund SE, Dahl LB. A randomized, controlled trial of the effectiveness of an early-intervention program in reducing parenting stress after preterm birth. *Pediatrics.* 2006;118:e9-e19.
83. Kaarsen PI, Rønning JA, Tunby J, Nordhov SM, Ulvund SE, Dahl LB. A randomized controlled trial of an early intervention program in low birth weight children: outcome at 2 years. *Early Hum Dev.* 2008;84:201-209.
84. Newnham CA, Milgrom J, Skouteris H. Effectiveness of a modified mother-infant transaction program on outcomes for preterm infants from 3 to 24 months of age. *Infant Behav Dev.* 2009;32:17-26.
85. Olafsen KA, Rønning JA, Kaarsen PI, Ulvund SE, Handegard BH, Dahl LB. Joint attention in term and preterm infants at 12 months corrected age: the significance of gender and intervention based on a randomized controlled trial. *Infant Behav Dev.* 2006;29:554-563.
86. Randomised trial of parental support for families with very preterm children. Avon Premature Infant Project. *Arch Dis Child Fetal Neonatal Ed.* 1998;79:F4-F11.
87. Gianni ML, Piccolini O, Ravasi M, et al. The effects of an early developmental mother-child intervention program on

- neurodevelopment outcome in very low birth weight infants: a pilot study. *Early Hum Dev.* 2006;82:691-695.
88. Sajaniemi N, Mäkelä J, Salokorpi T, von Wendt L, Hämäläinen T, Hakamies-Blomqvist L. Cognitive performance and attachment patterns at four years of age in extremely low birth weight infants after early intervention. *Eur Child Adolesc Psychiatry.* 2001;10:122-129.
 89. Johnson S, Ring W, Anderson P, Marlow N. Randomised trial of parental support for families with very preterm children: outcome at 5 years. *Arch Dis Child.* 2005;90:909-915.
 90. Landry SH, Smith KE, Swank PR. Responsive parenting: establishing early foundations for social, communication, and independent problem-solving skills. *Dev Psychol.* 2006;42:627-642.
 91. Guttentag CL, Pedrosa-Josic C, Landry SH, Smith KE, Swank PR. Individual variability in parenting profiles and predictors of change: effects of an intervention with disadvantaged mothers. *J Appl Dev Psychol.* 2006;27:349-369.
 92. Dieterich SE, Landry SH, Smith KE, Swank PR, Hebert HM. Impact of community mentors on maternal behaviors and child outcomes. *J Early Interv.* 2006;28:111-124.
 93. Landry SH, Smith KE, Swank PR, Guttentag C. A responsive parenting intervention: the optimal timing across early childhood for impacting maternal behaviors and child outcomes. *Dev Psychol.* 2008;44:1335-1353.
 94. Enhancing the outcomes of low-birth-weight, premature infants: a multisite, randomized trial. The Infant Health and Development Program. *JAMA.* 1990;263:3035-3042.
 95. Gross RT, Spiker D, Haynes CW. *Helping Low Birth Weight Premature Babies: The Infant Health and Development Program.* Stanford, CA: Stanford University Press; 1997.
 96. McCormick MC, McCarton C, Brooks-Gunn J, Belt P, Gross RT. The Infant Health and Development Program: interim summary. *J Dev Behav Pediatr.* 1998;19:359-370.
 97. Brooks-Gunn J, McCarton CM, Casey PH, et al. Early intervention in low-birth-weight premature infants. Results through age 5 years from the Infant Health and Development Program. *JAMA.* 1994;272:1257-1262.
 98. McCarton CM, Brooks-Gunn J, Wallace IF, et al. Results at age 8 years of early intervention for low-birth-weight premature infants. The Infant Health and Development Program. *JAMA.* 1997;277:126-132.
 99. McCormick MC, Brooks-Gunn J, Buka SL, et al. Early intervention in low birth weight premature infants: results at 18 years of age for the Infant Health and Development Program. *Pediatrics.* 2006;117:771-780.
 100. Klebanov PK, Brooks-Gunn J, McCormick MC. Maternal coping strategies and emotional distress: results of an early intervention program for low birth weight young children. *Dev Psychol.* 2001;37:654-667.
 101. Spiker D, Ferguson J, Brooks-Gunn J. Enhancing maternal interactive behavior and child social competence in low birth weight, premature infants. *Child Dev.* 1993;64:754-768.
 102. Martin A, Brooks-Gunn J, Klebanov P, Buka SL, McCormick MC. Long-term maternal effects of early childhood intervention: findings from the Infant Health and Development Program (IHDP). *J Appl Dev Psychol.* 2008;29:101-117.
 103. Hill JL, Brooks-Gunn J, Waldfogel J. Sustained effects of high participation in an early intervention for low-birth-weight premature infants. *Dev Psychol.* 2003;39:730-744.
 104. Sparling J, Lewis I, Ramey CT, Wasik BH, Bryant DM, LaVange LM. Partners: a curriculum to help premature, low birthweight infants get off to a good start. *Top Early Child Spec Educ.* 1991;11:36-55.
 105. Liaw FR, Meisels SJ, Brooks-Gunn J. The effects of experience of early intervention on low birth weight, premature children: the Infant Health and Development Program. *Early Child Res Q.* 1995;10:405-431.
 106. Dickinson DK, Porche MV. Relation between language experiences in preschool classrooms and children's kindergarten and fourth-grade language and reading abilities. *Child Dev.* 2011;82:870-886.
 107. Howes C, Burchinal M, Pianta R, et al. Ready to learn? Children's pre-academic achievement in pre-Kindergarten programs. *Early Child Res Q.* 2008;23:27-50.
 108. Blair C. Early intervention for low birth weight, preterm infants: the role of negative emotionality in the specification of effects. *Dev Psychopathol.* 2002;14:311-332.
 109. Spittle A, Orton J, Doyle LW, Boyd R. Early developmental intervention programs post hospital discharge to prevent motor and cognitive impairments in preterm infants. *Cochrane Database Syst Rev.* 2007;(2):CD005495.
 110. Luu TM, Vohr BR, Allan W, Schneider KC, Ment LR. Evidence for catch-up in cognition and receptive vocabulary among adolescents born very preterm. *Pediatrics.* 2011;128:313-322.
 111. Ment LR, Vohr B, Allan W, et al. Change in cognitive function over time in very low-birth-weight infants. *JAMA.* 2003;289:705-711.
 112. Medical Home Initiatives for Children With Special Needs Project Advisory Committee. American Academy of Pediatrics. The medical home. *Pediatrics.* 2002;110(Pt 1):184-186.
 113. Johnson S, Marlow N. Developmental screen or developmental testing? *Early Hum Dev.* 2006;82:173-183.
 114. Council on Children With Disabilities; Section on Developmental Behavioral Pediatrics; Bright Futures Steering Committee; Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disorders in the medical home: an algorithm for developmental surveillance and screening. *Pediatrics.* 2006;118:405-420.
 115. Garg A, Dworkin PH. Applying surveillance and screening to family psychosocial issues: implications for the medical home. *J Dev Behav Pediatr.* 2011;32:418-426.