

Medication Reconciliation

National Patient Safety Goal (NPSG) 8: Accurately and completely reconcile medications across the continuum of care.

References

1. Rozich JD, Resar RK. Medication Safety: One organization's approach to the challenge. *JCOM*. 2001;8(10):27-34.
2. Institute for Healthcare Improvement. **100,000 Lives Campaign. How-to Guide: Prevent Adverse Drug Events (Medication Reconciliation)**. <http://www.ihf.org/IHI/Programs/Campaign/>. Last accessed 8/31/05.
3. Whittington J, Cohen H. OSF Healthcare's journey in patient safety. *Quality Management in Health Care*. 2004;13(1):53-59.
4. Michels RD, Meisel S. Program using pharmacy technicians to obtain medication histories. *Am J Health-Sys Pharm*. October 1, 2003;60:1982-1986.
5. Branowicki P. *Sentinel Events: Opportunities for Change*. Presentation at Massachusetts Coalition for the Prevention of Medical Errors Conference, November 18, 2002.
6. Rozich JD, Resar RK, et. al. Standardization as a mechanism to improve safety in health care: impact of sliding scale insulin protocol and reconciliation of medications initiatives. *Joint Commission Journal on Quality and Safety*. 2004;30(1):5-14.

A University of Washington Drug Information Center publication
Distributed monthly by authority of the Pharmacy and Therapeutics Committee
Editor: Nelda A. Murri, Pharm.D. (206) 598-6612 – Asst. Editor: Elizabeth Rudy, D.V.M., R.Ph.
Department of Pharmacy Services / School of Pharmacy

Copyright © 2005 by the University of Washington
Also published on the World Wide Web at <http://uw.pnrx.org/therapyTopics.asp>
No material may be reproduced in whole or in part without written permission from the editor.