

Exercise Effects on Methylation of ASC Gene

Authors

K. Nakajima^{1,7}, M. Takeoka¹, M. Mori², S. Hashimoto³, A. Sakurai⁴, H. Nose⁵, K. Higuchi², N. Itano¹, M. Shiohara⁶, T. Oh⁷, S. Taniguchi¹

Affiliations

Affiliation addresses are listed at the end of the article

Key words

- interval walking
- pyrosequencing
- age

Abstract

▼
Chronic moderate exercise has been reported to reduce pro-inflammatory cytokines. To analyze the molecular mechanisms by which training exerts these effects, the epigenetic influences of age and exercise on the ASC gene, which is responsible for IL-1 β and IL-18 secretion, were investigated by ASC gene methylation. Further, the relationship between carcinogenesis and exercise, and methylation of the *p15* tumor suppressive gene was also analyzed. High-intensity interval walking exercise, consisting of 3 min low-intensity walking at 40% of peak aerobic capacity followed by a 3 min high-intensity walking period above 70% of peak aerobic capacity, was continued for 6 months. Peripheral blood

DNA extracts from young control (n=34), older control (n=153), and older exercise (n=230) groups were then analyzed by pyrosequencing for DNA methylation. Methylation of ASC decreased significantly with age (young control vs. older control, $p < 0.01$), which is indicative of an age-dependent increase in ASC expression. Compared to the older control group, the degree of ASC methylation was higher in the older exercise group (older control vs. older exercise: $p < 0.01$), and presumably lower ASC expression. Neither exercise nor age affected the methylation of the *p15*. In summary, chronic moderate exercise appears to attenuate the age-dependent decrease in ASC methylation, implying suppression of excess pro-inflammatory cytokines through reduction of ASC expression.

Introduction

▼
The relationship between exercise and susceptibility to infection has been modeled in the form of a “J”-shaped [18] or “S”-shaped curve [13]. These models imply that although moderate activity may enhance immune function, excessive amounts of prolonged and highly intensive exercise may actually impair function. Other anti-inflammatory effects mediated through the suppression of pro-inflammatory cytokines were also reported to be possible benefits of moderate exercise.

Chronic inflammation caused by excess pro-inflammatory cytokines, such as TNF- α , IL-1 β and IL-6, plays a prominent role in the development and progression of several age- and inactivity-related diseases. The relationship between chronic moderate exercise and reduced inflammation has been proven in several reports, which also identified some of the molecular mechanisms involved.

Cytosolic pathogen receptors have also garnered attention in the field of inflammation and innate

immunity. One such molecule, ASC (apoptosis-associated speck-like protein containing a caspase recruitment domain), is an adapter molecule consisting of a caspase recruit domain (CARD) and a pyrin domain (PYD) that was first revealed in our laboratory as a proapoptotic protein [15,24]. ASC functions as a mediator in the cytosol-type inflammatory signaling pathway, as well as a downstream molecule in toll-like receptors (TLRs) signaling. It has been established to activate procaspase-1 [22] and process proIL-1 β and proIL-18 to IL-1 β and IL-18, respectively [14,28] and ultimately lead to the initiation of innate immunity. ASC gene expression is recognized to be modified epigenetically, i.e. is down regulated by the methylation of its CpG island surrounding exon 1, and the methylation was inversely correlated with ASC protein expression [11,29]. Here, we investigated how ASC activity is epigenetically influenced by age or chronic moderate exercise by assessing the methylation of the CpG islands in the ASC gene using bisulfate genomic sequencing methods.

accepted after revision
December 06, 2009

Bibliography

DOI <http://dx.doi.org/10.1055/s-0029-1246140>
Published online: 2010
Int J Sports Med
© Georg Thieme
Verlag KG Stuttgart · New York
ISSN 0172-4622

Correspondence

Dr. Shun'ichiro Taniguchi
Department of Molecular
Oncology, (Institute on Aging
and Adaptation)
Shinshu University Graduate
School of Medicine
3-1-1 Asahi
390-8621 Matsumoto
Japan
Tel.: +81/263-37-2679
Fax: +81/263-37-2724
stangch@shinshu-u.ac.jp

Table 1 Clinical data of subjects.

	Young control group	Older control group	Older exercise group
number (n)	34	153	230
age range (years)	18–22**	40–87	41–86
age (years)	19.4±0.9	64.8±7.5	66.2±7.1
height (cm)	165.3±7.6	156.8±7.4	155.8±7.4
weight	58.4±11.3	58.9±9.7	57.8±8.9
BMI	21.1±3.2**	24.0±2.3	23.8±1.9
body fat (%)	19.2±8.0**	29.7±6.5	30.7±6.7
HDL cholesterol (mg/dl)	–	69.3±15.9	69.0±16.6
LDL cholesterol (mg/dl)	–	131.9±29.5	132.4±28.3
triglycerides (mg/dl)	–	108.2±58.9	102.9±49.4
VO _{2max} (ml/kg/min)	41.5±8.5**	23.8±4.3	23.8±5.1
glucose (mg/dl)	–	101.9±12.8	105.9±19.9*

Data are written as mean ± SD. * $p < 0.05$, ** $p < 0.01$ vs. Older control group

Another representative gene whose expression is epigenetically changed by methylation is the tumor suppressor gene *p15* [1]. Since the relationship between exercise and cancer has also been reported [27], *P15* was added to the present analysis. Taken together, the main purpose of this study was to investigate whether such epigenetic changes as methylation of the *ASC* and *p15* genes are influenced by age or chronic moderate exercise, namely a 6-month high-intensity interval walking regimen [17], in older subjects.

Materials and Methods

Subjects and exercise application

A total of 436 (142 male and 294 female) healthy subjects in two groups who participated in The Health-promotion Program for Elderly People in Matsumoto (Jukunen Taiiku Daigaku) was the object of our present research. One was the older control group ($n = 162$, age 40–87, 64.9 ± 7.4 years) and the other was the older exercise group ($n = 274$, age 41–86, 65.7 ± 7.4 years). For the young control group, 37 university students were recruited from Matsumoto University (age 18–22, 19.4 ± 0.9 years).

Subjects following the high-intensity interval walking regimens were instructed to repeat several sets of 3 min low-intensity walking at 40% of peak aerobic capacity followed by a 3 min high-intensity walking period above 70% of peak aerobic capacity for 6 months. Exercise intensity and time were supervised by accelerometry (Jukudai mate, Kissei Comtec Co., Ltd, Nagano). Mean walking time was 3.9 ± 1.2 days/week, 52.2 ± 18.5 min/day. The lowest acceptable training frequency was 2 days/week and training time was 26 min/day. Close monitoring was maintained for the first month of the regimen until subjects could learn high-intensity interval walking. Every two weeks, the subjects visited a local office to transfer the data from their tracking devices to a central server at the administrative center for automatic analysis and reporting. Trainers used these reports to track daily walking intensity and other parameters and to instruct subjects on how to best achieve exercise target levels. Dropouts were excluded from the beginning. Clinical data of the subjects are shown in **Table 1**.

This study has been approved by the Committee on the Use of Human Research Subjects of Shinshu University School of Medicine and Matsumoto University, and also performed in accordance with the ethical standards of the IJSM [12]. Participant's consent was received.




Fig. 1 Schematic view of the *ASC* gene region analyzed in this study. Seven CpG sites in the CpG island were designated for pyrosequencing analysis based on the effective region from Conway et al. [5].

Bisulfite treatment

Genomic DNA was extracted from the peripheral blood of control groups and subjects after a 6-month exercise period using a QIAamp DNA Mini Kit (QIAGEN, MD, USA). Blood sampling was performed preprandially at approximately 8:30 in the morning around one week after the last session of interval walking. From that, 1 µg of DNA was treated with sodium bisulfite using a CpGenome DNA Modification Kit (Chemicon, Temecula, CA, USA) according to manufacturer's instructions.

PCR and pyrosequencing

Bisulfite-modified DNA was resuspended in a total volume of 50 µl elution buffer and 0.3 µl, for *ASC*, amplified by a 1st-round of PCR (Go Taq DNA polymerase, Promega, Madison, WI, USA) and then adjusted to a final volume of 20 µl with TE buffer. The 2nd-round (nested) amplification was performed using 1 µl of the first-round PCR product, and then the 2nd-round PCR product was adjusted to a final volume of 50 µl with TE buffer. For *p15*, 5 µl bisulfite modified DNA was amplified by PCR and then adjusted to a final volume of 70 µl with TE buffer. Seven sites in the CpG island of *ASC* designated by Conway et al. [5] (**Fig 1**) and 6 sites in *p15* cited by Bollati et al. [1] were selected for pyrosequencing analysis. The primers and PCR conditions used are shown in **Table 2**.

DNA methylation was quantified using bisulfite-PCR and pyrosequencing of 50 µl or 70 µl of PCR products [7] (Biotage, Summit

Table 2 Primers and conditions used for ASC and p15 PCR.

Gene	Forward primer (5'to 3')	Reverse primer (5'to 3')	Sequencing primer (5'to 3')	PCR conditions
ASC				
1 st (423bp)	TGTATTAGTTGGTGAAG	CACACCCACAACAAC		95 °C-5m, (95 °C-30s; 50 °C-30s, 72 °C-30s)×3, (95 °C-30s, 58 °C-30s, 72 °C-30s)×47, 72 °C-5m
nested (237bp)	AGGGGATTAAGGGTG	Biotin -CTCCTCCACCAT	GGGATTTGGAG	95 °C-5m, (95 °C-30s; 58 °C-30s, 72 °C-20s)×50, 72 °C-5m
P15 (133bp)	GTTTTTTTGTAGAAG	Biotin-CCTTCTACRACT	GTTAGGAAAAGTT	95 °C-5m, (95 °C-30s; 50 °C-30s, 72 °C-30s)×50, 72 °C-5m

Pharmaceuticals Intl. Corp., Tokyo, Japan). Individual ASC gene pyrosequencing data, whose all 7 sites were judged as successful, were subjected to statistical analysis. After adjustments for technical error, the size of the older control group, older exercise group, and young control group used for statistic analysis was 153, 230 and 34, respectively.

Statistical analysis

The Student's t-test and ANOVA followed by the post-hoc test were used to test for differences in methylation levels among categorical variables. An F-test for ANOVA was performed to evaluate for normal distribution of test data. Correlation of clinical data with ASC methylation was estimated using Pearson's correlation coefficient, and association of data with ASC methylation was evaluated by both stepwise (backward elimination) and forced entry multivariable linear regression analysis with SPSS version 11.0J software (SPSS Japan Inc., Japan). A *P* value of less than 0.05 was considered statistically significant. Values were expressed as mean±standard deviation (SD) of the mean for characteristic data and mean±standard error (SE) of the mean for methylation data.

Results



Clinical data of subjects

The values of body mass index (BMI) and body fat were significantly lower ($p < 0.01$), and VO_{2max} was higher ($p < 0.01$) in young control group than in older control group. As far as within older groups, there was a significant difference in glucose value ($p < 0.05$) between the older control group and older exercise group (Table 1). When pre- (no methylation data) and post-exercise clinical data in the older exercise group were compared, only VO_{2max} was found to be significantly increased by exercise (21.9 ± 4.3 to 23.8 ± 5.1 ml/kg/min, $p < 0.01$).

ASC findings

A schematic view of the ASC gene region analyzed in this study is indicated in Fig. 1. ASC methylation in the older control group ($5.33 \pm 0.14\%$) was significantly lower than that in the young control group ($6.73 \pm 0.38\%$, $p < 0.01$, Fig 2), indicating that expression of ASC increased age-dependently. Analysis using Pearson's correlation coefficient showed that age ($p < 0.001$), BMI ($p < 0.005$), body fat ($p < 0.001$), and VO_{2max} ($p < 0.001$) among clinical data in Table 1 were correlated with ASC methylation between young control and older control groups. Since only age (standardized multivariate regression coefficient beta -0.422 , $p < 0.001$; $R^2 = 0.178$, $p < 0.001$) remained by stepwise with backward elimination multivariable linear regression analysis for the association with ASC methylation, forced entry multivariable linear regression analysis was then applied. The analysis also




Fig. 2 Methylation of ASC and age. Difference in the mean methylation values of 7 sites between the young control group ($n = 34$) and the older control group ($n = 153$). Values are written as mean ± SE. ** $p < 0.01$.

Table 3 Association of data with ASC methylation was evaluated by forced entry multivariable linear regression analysis.

	B	SE	Beta	p-Value
age	0.042	0.012	-0.396	0.001
BMI (%)	0.005	0.052	0.009	0.920
body fat (%)	0.043	0.023	-0.175	0.059
VO_{2max} (ml/kg/min)	0.012	0.030	-0.045	0.696

$R^2 = 0.196$, $p < 0.001$

revealed that ASC methylation was associated with age (beta -0.396 , $p < 0.001$), but not BMI (beta 0.009 , $p < 0.920$), body fat (beta -0.175 , $p < 0.059$), or VO_{2max} (beta -0.045 , $p < 0.696$, Table 3), indicating that age was the strongest contributor to ASC methylation.

Next, the influence of chronic moderate exercise on ASC methylation was investigated in older subjects. As shown in Fig. 3 Left, methylation was higher in the exercise group than in the control group at all 7 sites in the CpG island. At 6 of these sites, the difference was significant ($p < 0.05$ or $p < 0.01$). The mean methylation value of the 7 sites together was significantly higher in the older exercise group ($6.29 \pm 0.26\%$) than in the older control group ($5.33 \pm 0.14\%$, $p < 0.01$, Fig. 3 Right), clearly indicative of a decrease in ASC expression by chronic moderate exercise. Examination of data using Pearson's correlation coefficient showed that exercise days ($p < 0.030$) was correlated with ASC methylation between older control and older exercise groups, whereas other clinical data in Table 1, including glucose ($p < 0.083$), were not.

The relationship between ASC methylation, age, and exercise is summarized in Fig. 4. The mean methylation value was significantly higher in the older exercise group than in the older control group and approximated a value closer to that of the




Fig. 3 Methylation of ASC and exercise in older groups. Left: Methylation rates of all 7 targeted sites. Open circles: older control group (n=153). Closed circles: older exercise group (n=230), after 6-month interval walking regimen. Right: Difference in mean methylation values of the 7 sites between the older control group and the older exercise group. Values are written as mean \pm SE. * $p < 0.05$, ** $p < 0.01$




Fig. 4 Relationship between methylation of ASC, age, and exercise. Dashed arrows indicate the schematically estimated age of youth restoration caused by the exercise. Values are written as mean \pm SE. * $p < 0.05$, ** $p < 0.01$. Significance is versus older control group.

young control group, indicating that the methylation state of ASC was markedly improved by the exercise.

p15 findings

The methylation rate did not change age-dependently in the case of *p15*, nor was it affected by the 6-month exercise regimen (data not shown).

Discussion

In the present study, the methylation of ASC was found to decrease in an age-dependent fashion and implied that ASC expression increased with age. Higher ASC methylation in the older exercise group than in the control group indicates that moderate exercise may restore ASC methylation to comparably younger levels. Neither moderate exercise nor age affected methylation of the *p15* tumor suppressor gene.

BMI and body fat were lower and VO_{2max} was higher in the young control group than in the older control group, therefore we analyzed the association of age and those clinical data with ASC methylation by using forced entry multivariable linear regression analysis following Pearson's correlation coefficient estimation, and found that age was the main factor associated with ASC methylation between young and older control groups. There was a difference in glucose values between the older control and

older exercise groups, however, exercise was primarily associated with ASC methylation in those two groups.

Adequate amounts of cytokines are required for innate immune function, however, elevation of TNF- α , IL-1 β and/or IL-6 is observed in type 2 diabetes [21], rheumatoid arthritis [4] and atherosclerosis patients [2]. TNF- α , IL-6 levels [19] and IL-1, TNF- α levels [3,6] have been reported to increase age-dependently as well. Thus, the development and progression of several age-related diseases are associated with chronic inflammation. Chronic training has been shown to reduce the expression of pro-inflammatory cytokines, including TNF- α , IL-1 β , and IL-6 [8,9,26]. According to Greiwe et al. [10], expression of TNF- α in skeletal muscle is higher in elderly subjects compared to young subjects, but decreases after 3 months of resistance exercise. Our present results concerning ASC methylation are consistent with these studies. Since skeletal muscles are reported to produce and express cytokines regulated by exercise [20], ASC methylation should be measured not only in blood cells, but also in voluntary muscle. However, obtaining informed consent was quite difficult in volunteer subjects, which constitutes a challenge in future studies.

ASC is reported to be required for the induction not only of IL-1 β , IL-18, but also of IL-6, IL-8, IL-10, and TNF through both caspase-1-dependent and -independent mechanisms [25]. Chronic moderate exercise reduces IL-6, TLR4 [23] and IL-6, IL-1 β , TNF- α , and TLR4 expression [16]. Since TLR4 is one of the upstream receptors of ASC [14], ASC together with TLR4 might reduce inflammatory cytokines by chronic moderate exercise.

Our findings lack the measurement of cytokines, such as IL-1 β and IL-6. Although the measurement of them is important, such studies with volunteer participants are usually complicated by difficulties in obtaining consent for larger amounts of blood. The same scenario applies to the measurement of ASC protein changes associated with ASC methylation. However, we have already been able to confirm that a decrease in ASC protein expression is associated with increased methylation [11,29]. Also, the exercise period in this study was 6 months, but a one year regime may be more revealing of the effects of interval walking since other unknown factors may have affected ASC methylation in the current experiment.

Lastly, the *P15* gene was selected to investigate the influence of exercise on carcinogenesis. No influence was observed, however, *P15* became a methodologically good reference result. From this finding, discrepancies among groups in terms of season, classification, exercise quantity, and so on could be excluded.

As we described in Materials, the exercise group had just finished its 6-month exercise regimen when we collected blood from older subjects. Ideally, DNA samples from the exercise group should also have been collected prior to starting exercise, however, such DNA samples for methylation analysis were unobtainable at the time, since our experimental protocol was under consideration by the University's ethical committee. To strengthen the usefulness of our data, we are planning to compare the methylation data from a control group with the data after 6 months of exercise in a future study. Nonetheless, we believe that the present results are still useful as a novel finding which may impact the development of molecular research in sports medicine. As far as we have examined, the differences in methylation of the ASC gene were attributed to no other apparent factors besides training.

In summary, chronic moderate exercise, such as intense interval walking, can increase ASC methylation and thereby may suppress excess pro-inflammatory cytokine expression.

Acknowledgements

▼
We thank the members of Shinshu University Genetic Research Consortium (Shinshu University) and Ms S. Nakajima (Matsumoto University) for technical supports and valuable discussions. This work was supported by the Shinshu University Partnership Project between Shinshu University, Jukunen Taiiku Daigaku Research Center: JTRC, and the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Affiliations

- ¹ Department of Molecular Oncology, Institute on Aging and Adaptation, Shinshu University Graduate School of Medicine, Matsumoto, Japan
- ² Department of Aging Biology, Institute on Aging and Adaptation, Shinshu University Graduate School of Medicine, Matsumoto, Japan
- ³ Preventive Medical Center, Shinshu University Hospital, Matsumoto, Japan
- ⁴ Department of Medical Genetics, Shinshu University School of Medicine, Matsumoto, Japan
- ⁵ Department of Sports Medical Sciences, Institute on Aging and Adaptation, Shinshu University Graduate School of Medicine, Matsumoto, Japan
- ⁶ Department of Pediatrics, Shinshu University School of Medicine, Matsumoto, Japan
- ⁷ Department of Sports and Health Science, Faculty of Human Health Science, Matsumoto University, Matsumoto, Japan

References

- 1 Bollati V, Baccarelli A, Hou L, Bonzini M, Fustinoni S, Cavallo D, Byun HM, Jiang J, Marinelli B, Sestori AC, Bertazzi PA, Yang AS. Changes in DNA methylation patterns in subjects exposed to low-dose benzene. *Cancer Res* 2007; 67: 876–880
- 2 Calabró P, Willerson JT, Yeh ET. Inflammatory cytokines stimulated C-reactive protein production by human coronary artery smooth muscle cells. *Circulation* 2003; 108: 1930–1932
- 3 Casolini P, Catalani A, Zuena AR, Angelucci L. Inhibition of COX-2 reduces the age-dependent increase of hippocampal inflammatory markers, corticosterone secretion, and behavioral impairments in the rat. *J Neurosci Res* 2002; 68: 337–343
- 4 Choy EH, Panayi GS. Cytokine pathways and joint inflammation in rheumatoid arthritis. *N Engl J Med* 2001; 344: 907–916
- 5 Conway KE, McConnell BB, Bowring CE, Donald CD, Warren ST, Vertino PM. TMS1, a novel proapoptotic caspase recruitment domain protein, is a target of methylation-induced gene silencing in human breast cancers. *Cancer Res* 2000; 60: 6236–6242
- 6 Dayan M, Segal R, Globerson A, Habut B, Shearer GM, Mozes E. Effect of aging on cytokine production in normal and experimental systemic lupus erythematosus-afflicted mice. *Exp Gerontol* 2000; 35: 225–236
- 7 England R, Pettersson M. Pyro Q-CpGTM: quantitative analysis of methylation in multiple CpG sites by Pyrosequencing®. *Nat Methods* 2005; 2: 1–2
- 8 Flynn MG, McFarlin BK, Phillips MD, Stewart LK, Timmerman KL. Toll-like receptor 4 and CD14 mRNA expression are lower in resistive exercise-trained elderly women. *J Appl Physiol* 2003; 95: 1833–1842
- 9 Gielen S, Adams V, Möbius-Winkler S, Linke A, Erbs S, Yu J, Kempf W, Schubert A, Schuler G, Hambrecht R. Anti-inflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. *J Am Coll Cardiol* 2003; 42: 861–868
- 10 Greive JS, Cheng B, Rubin DC, Yarasheski KE, Semenkovich CF. Resistance exercise decreases skeletal muscle tumor necrosis factor alpha in frail elderly humans. *FASEB J* 2001; 15: 475–482
- 11 Guan X, Sagara J, Yokoyama T, Koganehira Y, Oguchi M, Saida T, Taniguchi S. ASC/TMS1, a caspase-1 activating adaptor, is downregulated by aberrant methylation in human melanoma. *Int J Cancer* 2003; 107: 202–208
- 12 Harriss DJ, Atkinson G. International Journal of Sports Medicine – Ethical standards in sport and exercise science research. *Int J Sports Med* 2009; 30: 701–702
- 13 Malm C. Susceptibility to infections in elite athletes: the S-curve. *Scand J Med Sci Sports* 2006; 16: 4–6
- 14 Mariathasan S, Monack DM. Inflammasome adaptors and sensors: intracellular regulators of infection and inflammation. *Nat Rev Immunol* 2007; 7: 31–40
- 15 Masumoto J, Taniguchi S, Ayukawa K, Sarvatham H, Kishino T, Niikawa N, Hidaka E, Katsuyama T, Higuchi T, Sagara J. ASC, a novel 22-kDa protein, aggregates during apoptosis of human promyelocytic leukemia HL-60 cells. *J Biol Chem* 1999; 274: 33835–33838
- 16 McFarlin BK, Flynn MG, Campbell WW, Craig BA, Robinson JP, Stewart LK, Timmerman KL, Coen PM. Physical activity status, but not age, influences inflammatory biomarkers and toll-like receptor 4. *J Gerontol A Biol Sci Med Sci* 2006; 61: 388–393
- 17 Nemoto K, Gen-no H, Masuki S, Okazaki K, Nose H. Effects of high-intensity interval walking training on physical fitness and blood pressure in middle-aged and older people. *Mayo Clin Proc* 2007; 82: 803–811
- 18 Nieman DC. Exercise, upper respiratory tract infection, and the immune system. *Med Sci Sports Exerc* 1994; 26: 128–139
- 19 O'Mahony L, Holland J, Jackson J, Feighery C, Hennessy TP, Mealy K. Quantitative intracellular cytokine measurement: age-related changes in proinflammatory cytokine production. *Clin Exp Immunol* 1998; 113: 213–219
- 20 Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev* 2008; 88: 1379–1406
- 21 Spranger J, Kroke A, Möhlig M, Hoffmann K, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF. Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *Diabetes* 2003; 52: 812–872
- 22 Stehlik C, Lee SH, Dorfleitner A, Stassinopoulos A, Sagara J, Reed JC. Apoptosis-associated speck-like protein containing a caspase recruitment domain is a regulator of procaspase-1 activation. *J Immunol* 2003; 171: 6154–6163
- 23 Stewart LK, Flynn MG, Campbell WW, Craig BA, Robinson JP, McFarlin BK, Timmerman KL, Coen PM, Felker J, Talbert E. Influence of exercise training and age on CD14+ cell-surface expression of toll-like receptor 2 and 4. *Brain Behav Immun* 2005; 19: 389–397
- 24 Taniguchi S, Sagara J. Regulatory molecules involved in inflammasome formation with special reference to a key mediator protein, ASC. *Semin Immunopathol* 2007; 29: 231–238
- 25 Taxman DJ, Zhang J, Champagne CB, Bergstralh DT, Iocca HA, Lich JD, Ting JP. Cutting edge: ASC mediates the induction of multiple cytokines by Porphyromonas gingivalis via caspase-1- dependent and -independent pathways. *J Immunol* 2006; 177: 4252–4256
- 26 Venkatraman JT, Rowland JA, Denardin E, Horvath PJ, Pendergast D. Influence of the level of dietary lipid intake and maximal exercise on the immune status in runners. *Med Sci Sports Exerc* 1997; 29: 333–344
- 27 Wang JS, Chang CY, Chow SE, Chen YW, Yang CM. Exercise modulates platelet-nasopharyngeal carcinoma cell aggregation and subsequent tissue factor and matrix metalloproteinase activities. *J Appl Physiol* 2007; 103: 763–770
- 28 Yamamoto M, Yaginuma K, Tsutsui H, Sagara J, Guan X, Seki E, Yasuda K, Yamamoto M, Akira S, Nakanishi K, Noda T, Taniguchi S. ASC is essential for LPS-induced activation of procaspase-1 independently of TLR-associated signal adaptor molecules. *Genes Cells* 2004; 9: 1055–1067
- 29 Yokoyama T, Sagara J, Guan X, Masumoto J, Takeoka M, Komiyama Y, Miyata K, Higuchi K, Taniguchi S. Methylation of ASC/TMS1, a proapoptotic gene responsible for activating procaspase-1, in human colorectal cancer. *Cancer Lett* 2003; 202: 101–108