

# Collaborative Care for Perinatal Depression Among Socioeconomically Disadvantaged Women: Adverse Neonatal Birth Events and Treatment Response

Amritha Bhat, M.B.B.S., M.D., Nancy K. Grote, Ph.D., Joan Russo, Ph.D., Mary Jane Lohr, M.S., Hyunzee Jung, Ph.D., Caroline E. Rouse, M.D., Elaine C. Howell, B.A., Jennifer L. Melville, M.D., M.P.H., Kathy Carson, B.S.N., Wayne Katon, M.D.

**Objective:** The study examined the effectiveness of a perinatal collaborative care intervention in moderating the effects of adverse neonatal birth events on risks of postpartum depressive symptoms and impaired functioning among women of lower socioeconomic status with antenatal depression.

**Methods:** A randomized controlled trial with blinded outcome assessments was conducted in ten public health centers, comparing MOMCare (choice of brief interpersonal psychotherapy, pharmacotherapy, or both) with intensive maternity support services (MSS-Plus). Participants had probable diagnoses of major depressive disorder or dysthymia during pregnancy. Generalized estimating equations estimated differences in depression and functioning measures between groups with and without adverse birth events within the treatment arms. A total of 160 women, 43% of whom experienced at least one adverse birth event, were included in the analyses.

**Results:** For women who received MOMCare, postpartum depression scores (measured with the Symptom Checklist-20) did not differ by whether or not they experienced an adverse birth event (mean  $\pm$  SD scores of  $.86 \pm .51$  for mothers with an adverse birth event and  $.83 \pm .56$  for mothers with no event;  $p = .78$ ). For women who received MSS-Plus, having an adverse birth event was associated with persisting depression in the postpartum period (mean scores of  $1.20 \pm .061$  for mothers with an adverse birth event and  $.93 \pm .52$  for mothers without adverse birth event;  $p = .04$ ). Similar results were seen for depression response rates and functioning.

**Conclusions:** MOMCare mitigated the risk of postpartum depressive symptoms and impaired functioning among women of low socioeconomic status who had antenatal depression and who experienced adverse birth events.

*Psychiatric Services in Advance* (doi: 10.1176/appi.ps.201600002)

Adverse birth events are more common among women who have depression during pregnancy (1). Antenatal depressive symptoms increase the relative risk of preterm birth (<37 weeks' gestation) by 39%, low birth weight (LBW) (<2,500 g) by 49%, and intrauterine growth retardation (below the tenth percentile for gestational age) by 45% (2). This risk is higher among women of lower socioeconomic class (2). Elevated depression scores at 32 weeks' gestation (3) and major depression diagnosed at any time during pregnancy (4) are also related to adverse birth events, with greater duration of antenatal depression associated with higher odds of admission to the neonatal intensive care unit (NICU) (5-7). Not only are adverse birth events more common among women with antenatal depression, but obstetric complications and adverse birth events are also associated with rates of postpartum depressive symptoms as high as 40% (8). Mothers with LBW babies continue to be at increased risk of depression at one year postpartum (9). For mothers with low socioeconomic status this period of risk extends to two years (10).

According to the stress-diathesis model (11), for a person with predisposing factors for depression (family history or past history), experiencing a significant stressor is more likely to lead to a depressive episode, compared with a person without this predisposition (12,13). Antenatal depression is one of the strongest predisposing factors for postpartum depression, even after normal full-term deliveries (14,15). Adverse birth events constitute a significant and stressful life event (16). Thus having an adverse birth event could increase the risk of postpartum depression (6) among women with antenatal depression.

Infants who experienced adverse birth events, such as prematurity or LBW, are at high risk of long-term negative outcomes, such as lower educational achievement and attention-deficit hyperactivity disorder (17), anxiety disorders, and autism spectrum disorder (18). In addition to prematurity, environmental factors, such as parental psychological distress, contribute to this set of risks for the infant. Mothers who have depression can be less responsive

and less sensitively attuned to their infants (19). Of note, maternal responsivity moderates the effects of LBW on hyperkinetic and internalizing behaviors among LBW babies (20,21). In other words, maternal postpartum depression, by reducing maternal responsivity, can magnify the risk of poor outcomes for preterm or LBW infants.

Thus any effort to prevent poor infant outcomes is incomplete without attention to maternal psychological distress, especially because maternal depression has been shown to mediate the remoteness seen in parenting interactions between mothers and their preterm babies (21). To that end, various interventions to reduce mothers' psychological distress after adverse birth events have been tested, including educational-behavioral programs (22), anxiety reduction (23), and sensitivity training (24). These programs have components of child development or parenting support; however, none addresses maternal depression specifically. Overall, although studies reported decreased stress (22,24) and increased rates of breastfeeding (25), most studies found no improvement in maternal depression (22–24).

With the exception of a single study (26), research has not addressed the population of women who were depressed during pregnancy and who experienced adverse birth events and, therefore, were at higher risk of postpartum depression (27). Postpartum support can improve maternal mental health in high-risk populations (28), and parents' coping after an adverse birth event has been reported to be influenced by the availability of social support (29). The literature suggests that general social support and parenting support may reduce risk of postpartum depression among mothers without antenatal depression who experience adverse birth events (22,30). However, mothers who have had antenatal depression and also experience adverse birth events may require more specialized support to reduce depression during the perinatal period (26).

We recently reported that the MOMCare collaborative care intervention for perinatal depression was more effective on average in reducing depression severity and increasing remission rates from pregnancy to 15 months postpartum than a comparison condition of intensive public health maternity support services (MSS-Plus) (31). The purpose of the analysis reported here was to determine to what extent MOMCare, compared with MSS-Plus, reduced postpartum depression severity for mothers who experienced one or more adverse birth events. That is, for the study sample of women who had antenatal depression, to what extent did the intervention moderate the negative effect of experiencing an adverse birth event on postpartum depression severity?

We hypothesized that despite having an adverse birth event, women with antenatal depression would have less postpartum depression severity if they received the MOMCare intervention rather than MSS-Plus. We also hypothesized that despite experiencing an adverse birth event, mothers in the MOMCare intervention would have better functioning in the postpartum period compared with mothers in MSS-Plus.

## METHODS

In a multisite randomized controlled trial with blinded assignment, we evaluated the MOMCare collaborative care intervention for perinatal depression compared with MSS-Plus in ten Seattle–King County public health centers from July 2009 to January 2011. Participants were recruited who met criteria for a probable major depressive disorder on the basis of the nine-item Patient Health Questionnaire (PHQ-9) (32) (at least five symptoms scored as  $\geq 2$ , with one cardinal symptom; and functional impairment noted on the PHQ-9) or a probable diagnosis of dysthymia on the Mini-International Neuropsychiatric Interview (33). Study procedures were approved by the University of Washington Institutional Review Board and monitored by a Data Safety Monitoring Board. All participants signed study consent and HIPAA release forms, allowing the project to request their hospital labor, delivery, and birth information; 98% of hospital records were obtained.

### Participants

The original study sample included 168 Medicaid-enrolled pregnant women; 83 were randomly assigned to the MOMCare intervention and 85 to the MSS-Plus comparison group. Randomization was by an adaptive block scheme, stratified on initial depression severity via the 20-item Symptom Checklist (SCL-20) (34) ( $\leq 2.0$  or  $> 2.0$ ) and gestational age ( $< 22$  weeks or  $\geq 22$  weeks). The analyses reported here included 160 women; it excluded those for whom hospitals failed to provide medical records ( $N=4$ ) and those who dropped out of the study prior to the baseline interview ( $N=4$ ).

### MSS-Plus

Maternity support services (MSS) is the usual standard of care in the public health system of Seattle–King County for Medicaid-enrolled pregnant women, delivered by a multidisciplinary team of public health social workers, nurses, and nutritionists, who screen at least once for depression from pregnancy to two months postpartum. Goals of usual MSS include services to promote healthy pregnancies and positive birth and parenting outcomes, case management services, and facilitating contact with an obstetric provider. Pregnant women with PHQ-9 scores  $\geq 10$  were eligible for intensive MSS-Plus services, entailing more frequent and longer visits from their multidisciplinary team from pregnancy to two months postpartum. MSS-Plus providers referred depressed patients for mental health treatment in the community or from their obstetric providers. The MSS-Plus providers did not provide evidence-based depression care, systematic outreach, or stepped care.

### MOMCare

The MOMCare intervention was added onto MSS-Plus and extended from pregnancy to 12 months postpartum (35). MOMCare was delivered by depression care specialists (DCSSs) in consultation with the principal investigator

**TABLE 1. Comparison of key components of MOMCare and Maternity Support Services–Plus (MSS-Plus)<sup>a</sup>**

Component	MOMCare	MSS-Plus
Study procedures	MOMCare DCS delivers psychoeducation about depression and depression self-help book in one-hour appointment. MOMCare DCS notifies MSS social worker of client's intervention condition.	MOMCare depression care specialist (DCS) delivers psychoeducation about depression and depression self-help book in one-hour appointment. MOMCare DCS notifies MSS-Plus social worker of client's intervention condition.
Pretreatment Acute interventions if PHQ-9 score is >10	Engagement session by MOMCare DCS and collaborative care team (PI, study psychiatrist, and OB provider) provide acute brief IPT or pharmacotherapy or both. MSS social worker or nurse provides maternity support services from pregnancy to 2 months postpartum.	No engagement session MSS-Plus social worker makes a referral for depression care in the community or from an OB provider. MSS-Plus social worker or nurse provides MSS-Plus from pregnancy to 2 months postpartum.
Maintenance treatment for depression Treatment tracking	Up to 1 year postpartum Weekly, monthly, or bimonthly assessments on PHQ-9 by DCS	None No assessments by MSS-Plus social worker
Supervision for depression care management	Weekly group and individual supervision of DCS by team PI and psychiatrist	No supervision

<sup>a</sup> Abbreviations: DCS, depression care specialist; IPT, interpersonal therapy; MSS, maternity support services (MSS-Plus is the intensive standard of care in the public health system of Seattle–King County for Medicaid-enrolled pregnant women scoring >10 on the PHQ-9); OB, obstetric; PHQ-9, 9-item Patient Health Questionnaire; PI, principal investigator

(NKG) and the study psychiatrist (WK). The DCSs collaborated with their patients' obstetric providers, providing updates on progress, and collaborating on medication management if indicated. Intervention services were provided in the public health centers, by phone, in community settings, and at home. The DCSs specifically provided depression care management, and the MSS-Plus providers did not (Table 1).

MOMCare was designed to decrease mental health treatment disparities in access to and quality of care (35). Novel components include a pretherapy engagement session to help resolve practical, psychological, and cultural barriers to care; patient choice of brief interpersonal psychotherapy (IPT), pharmacotherapy, or both; telephone sessions in addition to in-person visits; outreach for women who miss sessions; and DCSs who deliver both the intervention and case management to meet basic needs (35).

### Measures

The primary outcome measure was the SCL-20, a reliable and valid measure of depression (34). It is sensitive to change (36) and has good internal consistency (37) and good convergent validity with the items on the PHQ-9 (38). Scores range from 0 to 4, with higher scores indicating more severe symptoms. The mean score was used to measure overall depression severity and to calculate depression treatment response ( $\geq 50\%$  reduction from baseline depression severity) and remission (SCL-20 score  $<.5$ ).

Functional impairment was assessed by the Work and Social Adjustment Scale (WSAS) (39), a five-item, self-report scale that measures functional impairment attributable to an identified problem. It is reliable and valid and has high correlation with clinician reports. Questions were asked in reference to "emotional difficulties."

The Peripartum Events Scale (PES) (40) has established reliability and validity and is used to quantify labor, delivery, and birth outcome events obtained from hospital records. Two coders (one primary coder [HJ] and one reliability coder [CER]) were trained by the project's obstetrics and gynecology (OB/GYN) consultant physician. The primary coder had a doctoral degree in social work, and the reliability coder was a resident OB/GYN physician. The first ten cases were coded by both coders to establish reliability. Subsequently, 20% of the medical records were coded by both coders, and interrater reliability averaged 99% agreement. The project's OB/GYN consultant physician (JLM) and a resident physician provided assistance in coding when needed. The PES produces 11 subscales assessing a variety of stressful events associated with childbearing. For the purpose of this analysis, we defined an adverse neonatal birth event as any event endorsed on the PES infant outcome subscale (ten items) or whether the infant was placed in the NICU or special care nursery.

The Post-Traumatic Stress Disorder (PTSD) Checklist–Civilian Version is a 17-item questionnaire that assesses intrusive symptoms, avoidance, and arousal associated with traumatic events (41). A continuous PTSD severity score was calculated.

### Statistical Analyses

We first calculated descriptive statistics for the entire sample. Because previous analyses showed significantly reduced depression severity for mothers in the MOMCare intervention compared with those in MSS-Plus (31), we tested postpartum depression symptom levels between women with and without an adverse birth event within each intervention arm separately. Because the adverse birth event groups were

not based on randomization, we examined baseline differences between variables, including obstetric complications, parity (number of births after gestational age 20 weeks), and social support in these groups within each intervention arm by using chi-square analyses and t tests. We found no differences, and the variables were not utilized as adjustments in our models. However, because baseline PTSD symptom severity was found to have a substantial impact on depression symptom levels in our initial primary outcome analyses (31), we included it in our models.

We used generalized estimating equations (GEEs) to estimate differences in depression measures and functioning between women with and without adverse birth events within the treatment arm cohorts. The GEE longitudinal model allowed inclusion of all women, including those with missing assessments. All GEE models were fitted with robust error estimation and an unstructured covariance matrix. The dependent variables were SCL-20 depression severity and response and remission at six, 12, and 18 months post-baseline (that is, postpartum depression for a year after birth). Each model contained main effects for adverse birth event and time, adjusted for baseline depression severity and PTSD severity level. For the analysis of functioning, baseline WSAS level was used in the model in lieu of baseline depression. Means and standard errors were estimated from the GEE models.

**RESULTS**

There were 160 women in the analysis sample; 68 (43%) had one or more adverse birth events and 92 (58%) had no adverse event. Data on baseline demographic variables are presented in Table 2. There was no association between having an adverse birth event and treatment group; 47% (N=37) of the MSS-Plus group experienced an adverse birth event, compared with 40% (N=32) of the MOMCare group. Within the MSS-Plus and MOMCare groups, all baseline variables between those with and without an eventual adverse birth event were similar. Frequencies of events considered to be adverse are presented in Table 3.

**Adverse Birth Events and Postpartum Depression Severity**

Time was not a significant main effect or interaction in any of the models (Figure 1). In the MOMCare group, the model for continuous SCL-20 depression severity scores indicated a nonsignificant main effect of adverse birth outcome. The average depression score (average of scores at six, 12, and 18 months postbaseline) was .86 for the mothers with an adverse birth event, compared with .83 for those with no adverse birth event (Table 4). In contrast, in the MSS-Plus group, there was a significant main effect of adverse birth event (Wald  $\chi^2=4.16$ ,  $df=1$ ,  $p=.04$ ), with average depression severity scores of 1.20 for mothers with an adverse birth event, compared with .93 for those without an adverse birth event.

**TABLE 2. Baseline demographic and clinical variables of 160 study participants**

Variable	N	%
Race-ethnicity		
White	66	41
African American	38	24
Latina	37	23
Asian/Pacific Islander	11	7
American Indian/Alaska native	8	5
Nonwhite	94	59
Marital status		
Married	48	30
Living with a partner	54	34
Partner (not living with)	20	13
No partner	38	24
Education		
Less than high school	33	21
High school degree or GED	32	20
Some college or vocational school	77	48
College degree or higher	18	11
Employment		
Full- or part-time	56	35
Unemployed	104	65
Income $\leq$ \$10,000 <sup>a</sup>	65	42
Homeless	22	14
Past pregnancy	113	71
Previous postpartum depression	59	53
Unplanned pregnancy	115	72
Positive about the pregnancy (quite a bit or very)	100	63
Depressive disorder		
Major depression <sup>b</sup>	154	96
Dysthymia <sup>c</sup>	36	23
Depression and dysthymia	33	21
Any previous depressive episode	132	83
Age (range 18–44) (M $\pm$ SD)	27.10 $\pm$ 6.01	
Gestational age (M $\pm$ SD weeks)	23.61 $\pm$ 6.32	
N of previous depressive episodes for those reporting $\geq$ 1 (M $\pm$ SD)	5.55 $\pm$ 5.51	
Baseline functioning		
SCL-20 depression score (M $\pm$ SD) <sup>d</sup>	1.78 $\pm$ .59	
PTSD severity score <sup>e</sup>	48.65 $\pm$ 11.31	
Work and social functioning <sup>f</sup>	21.19 $\pm$ 8.74	

<sup>a</sup> Data available for 156 participants  
<sup>b</sup> Assessed by the 9-item Patient Health Questionnaire  
<sup>c</sup> Assessed by the Mini-International Neuropsychiatric Interview  
<sup>d</sup> 20-item Symptom Checklist. Possible scores range from 0 to 4; a score of  $\geq$ .5 indicates possible depression.  
<sup>e</sup> Assessed by the Post-Traumatic Stress Disorder Checklist–Civilian Version. Possible scores range from 17 to 85, with higher scores indicating more severe symptoms.  
<sup>f</sup> Assessed by the Work and Social Adjustment Scale. Possible scores range from 0 to 45, with higher scores indicating more symptoms or greater impairment.

Depression response results ( $\geq$ 50% reduction from baseline severity) were similar for MOMCare; the model showed a nonsignificant main effect of adverse birth event. The average depression response (average at six, 12, and 18 months postbaseline) was 51% for mothers with an adverse birth event, compared with 55% for those without an adverse birth event (Table 4). In contrast, in the MSS-Plus group, there was a significant main effect of birth event

**TABLE 3. Adverse birth events among the 160 study participants**


Event <sup>a</sup>	N	%
Preterm (<37 weeks)	14	9
Postterm (>41 weeks)	6	4
Intrauterine growth retardation, small for gestational age, or large for gestational age	23	14
1-minute Apgar score <6	9	6
5-minute Apgar score <8	5	3
Need for transfusion or exchange transfusion	1	1
Hypoglycemia	5	3
Hyperbilirubinemia	9	6
Treatment for sepsis	13	8
Meconium aspiration pneumonitis	1	1
Baby sent to any NICU or special care nursery <sup>b</sup>	25	16
Other <sup>c</sup>	23	14
Other life threatening	2	1

<sup>a</sup> Assessed by the Peripartum Events Scale. A total of 68 participants had one or more adverse birth events.

<sup>b</sup> NICU, neonatal intensive care unit

<sup>c</sup> Other events include congenital malformations, such as patent ductus arteriosus, ventricular septal defect, hydronephrosis, and hypospadias; metabolic disorders, such as galactosemia; and hematologic disorders, such as bone marrow suppression and hemoglobinopathies.

(Wald  $\chi^2=4.04$ ,  $df=1$ ,  $p=.04$ ), with an average depression response of 36% for the mothers with an adverse birth event, compared with 55% for mothers without an adverse event. Remission from depression (SCL-20 score <.5) was not

**FIGURE 1. Depression severity by intervention group over 18-month follow-up for women with major depression with and without adverse birth events<sup>a</sup>**

<sup>a</sup> Scores on the 20-item Symptom Checklist (SCL-20). Possible scores range from 0 to 4; a score of  $\geq .5$  indicates possible depression. Six months is three months postpartum, 12 months is nine months postpartum, and 18 months is 15 months postpartum. MSS, maternity support services. MSS-Plus is the intensive standard of care in the public health system of Seattle–King County for Medicaid-enrolled pregnant women scoring  $>10$  on the nine-item Patient Health Questionnaire.

significantly different between those with and without adverse events for either MOMCare or MSS-Plus.

### Adverse Birth Events and Postpartum Functioning

In the MOMCare cohort, the mothers with and without adverse birth events did not differ on WSAS functioning scores (Table 4). However, in the MSS-Plus group, those with an adverse birth event reported significantly more impairment (Wald  $\chi^2=4.29$ ,  $df=1$ ,  $p=.04$ ). The mean WSAS score was 15.22 for MSS-Plus group mothers with an adverse birth event, compared with 11.65 for those without an adverse birth event.

## DISCUSSION AND CONCLUSIONS

In this study, we examined the effectiveness of collaborative care for perinatal depression among mothers with antenatal depression who were enrolled in Medicaid and who experienced an adverse birth event such as preterm delivery, intrauterine growth retardation, “small for gestational age,” low Apgar score, or admission of the infant to a progressive care nursery or NICU. To our knowledge, this is the first study testing the effectiveness of an intervention in reducing postpartum depression among women with antenatal depression and low incomes who have experienced an adverse birth event.

The rate of adverse birth events of 43% in our sample is comparable to previous reports of 46% in other pregnant populations (26). Corresponding to the stress-diathesis hypothesis, women with antenatal depression (42) and women of low socioeconomic status (43) are vulnerable to postpartum depression. For these women, the stress of events, including premature birth and infant illness (44), can increase the risk of postpartum depression. Our intervention, the patient’s choice of IPT, pharmacotherapy, or both, appears to have mitigated the risk of worsening maternal postpartum depression severity and impaired functioning associated with adverse birth events (45).

For mothers who received the MOMCare intervention, postpartum depression scores and functioning did not differ among those who did or did not experience an adverse birth event. By contrast, for women who received the MSS-Plus comparison condition, having an adverse birth event was associated with persistent distress in the postpartum period. That is, without the additional, longer-lasting support of MOMCare, those in MSS-Plus fared worse in terms of postpartum depression severity and impaired functioning after an adverse birth event.

Adverse birth events such as preterm labor have also been associated with a perception of the child as vulnerable to medical illness—more so in women of lower socioeconomic status (46). Both postpartum depression and perceived child vulnerability can lead to disturbance of the mother-child interaction (47) and contribute to an increased risk of behavioral and emotional disorders’ persisting into adolescence (48). Thus interventions such as MOMCare that

**TABLE 4. Postpartum clinical outcomes for MOMCare and MSS-Plus participants with and without adverse birth events**

Measure	MOMCare Intervention (N=81)							MSS-Plus (N=79)						
	Adverse event (N=32)		No adverse event (N=49)		Between-group difference	95% CI	p	Adverse event (N=37)		No adverse event (N=42)		Between-group difference	95% CI	p
M	SD	M	SD	M				SD	M	SD	M			
SCL-20 <sup>a</sup>	.86	.51	.83	.56	-.04	-.28 to .21	.78	1.20	.61	.93	.52	-.27	-.52 to -.01	.04 <sup>b</sup>
WSAS <sup>c</sup>	10.55	6.84	11.54	7.35	-.99	-4.16 to 2.18	.54	15.22	6.87	11.65	7.39	3.57	.43 to 6.72	.03 <sup>b</sup>
	%	SD	%	SD	Between-group difference	95% CI	p	%	SD	%	SD	Between-group difference	95% CI	p
Response <sup>d</sup>	51	42	55	40	4	14 to 23	.64	36	38	55	41	19	10 to 36	.04 <sup>b</sup>
Remission <sup>e</sup>	33	40	43	41	10	-8 to 28	.30	21	33	31	38	10	-5 to 26	.22

<sup>a</sup> 20-item Symptom Checklist. Possible scores range from 0 to 4; a score of ≥.5 indicates possible depression.

<sup>b</sup> Significant main effect for group at 6-, 12-, and 18-month follow-ups, on average, with control for baseline values and PTSD severity

<sup>c</sup> Work and Social Adjustment Scale. Possible scores range from 0 to 45, with higher scores indicating more symptoms or greater impairment.

<sup>d</sup> ≥50% decrease in SCL-20 score from baseline

<sup>e</sup> SCL-20 score <.5

reduce risk of maternal postpartum depression severity after adverse birth events can have especially far-reaching benefits for high-risk mother-baby dyads.

A strength of the study was that it was conducted in a sample of socioeconomically disadvantaged women, a population at higher risk of both postpartum depression and adverse birth events. We have previously reported on the effectiveness of MOMCare in reducing depression among women of lower socioeconomic status with perinatal depression (31), and we also reported that for women with comorbid depression and PTSD, MOMCare performed better than MSS-Plus in terms of depression severity improvement (49). The analysis reported here adds to these findings, in finding that for mothers with the vulnerability of low socioeconomic status, antenatal depression, and at least one adverse birth event, receipt of the MOMCare intervention prevented continuing postpartum depression. We did not find a reduction in adverse birth events among women who received MOMCare; however, randomization was not based on risk of adverse birth events, and thus the study was not designed or powered to examine this question.

Randomization was not based on adverse events, which was a limitation. However, on comparison of multiple variables, the two groups (mothers with and without adverse birth events) were strikingly similar. Reports of psychiatric outcomes are limited to probable major depression and dysthymia and were based on rating scales and not diagnostic assessments. However, in the perinatal period, even subsyndromal symptoms of anxiety and depression can have implications for mother and child and are no less important than diagnoses (50,51). The PES does not specifically include the adverse event of LBW, which is an important outcome among women with perinatal depression. However, it does include “small for gestational age,” which factors in LBW. Although improvement in child outcomes is increasingly emphasized as a paired index of successful treatment of maternal depression, we did not have objective measures for this.

The United States continues to have high rates of adverse birth events, such as preterm birth (9.7% in 2014) and LBW (8% in 2014), with the highest rates found among women from racial-ethnic minority groups and women of low socioeconomic status (52). Given that low socioeconomic status and adverse birth events contribute to accumulation of risk for postpartum depressive symptoms, there is a compelling need to implement stepped care interventions, such as MOMCare, that are effective for socioeconomically disadvantaged women and other high-risk groups. IPT is effective in the treatment of postpartum depression (53), and as we report, it was effective in mitigating the risk of postpartum depressive severity after adverse birth events. In other words, for women with antenatal depression, MOMCare can reduce the risk of persistence of depression in the postpartum period despite the additional risk of an adverse birth event.

Providers caring for mothers with complicated deliveries and adverse birth events, for preterm LBW neonates, or for pregnant and postpartum women with mental health conditions are all well placed to identify persistent depression in the postpartum period. It is important for them to be aware of available effective interventions, such as MOMCare, that can be implemented in the context of caring for vulnerable infants.

**AUTHOR AND ARTICLE INFORMATION**

Dr. Bhat and Dr. Russo are with the Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, where the late Dr. Katon was affiliated (e-mail: amritha@uw.edu). Dr. Grote, Ms. Lohr, Dr. Jung, and Ms. Howell are with the School of Social Work, University of Washington, Seattle. Dr. Rouse is with the Department of Obstetrics and Gynecology, Brigham and Women’s Hospital, Boston. Dr. Melville is with Northwest Women’s HealthCare, Swedish Medical Center, Seattle. Ms. Carson is with the Seattle–King County Department of Public Health, Seattle.

This research was supported by grant R01 MH084897 from the National Institute of Mental Health (NIMH) to Dr. Grote. Dr. Bhat was supported by a postdoctoral fellowship from NIMH (T32 MH20021 Psychiatry–Primary Care Fellowship Program training grant).

The authors report no financial relationships with commercial interests. Received January 5, 2016; revision received March 25, 2016; accepted April 25, 2016; published online October 1, 2016.

## REFERENCES

- Accortt EE, Cheadle AC, Dunkel Schetter C: Prenatal depression and adverse birth outcomes: an updated systematic review. *Maternal and Child Health Journal* 19:1306–1337, 2015
- Grote NK, Bridge JA, Gavin AR, et al: A meta-analysis of depression during pregnancy and the risk of preterm birth, low birth weight, and intrauterine growth restriction. *Archives of General Psychiatry* 67:1012–1024, 2010
- Chung TK, Lau TK, Yip AS, et al: Antepartum depressive symptomatology is associated with adverse obstetric and neonatal outcomes. *Psychosomatic Medicine* 63:830–834, 2001
- Misri S, Oberlander TF, Fairbrother N, et al: Relation between prenatal maternal mood and anxiety and neonatal health. *Canadian Journal of Psychiatry* 49:684–689, 2004
- Latendresse G: The interaction between chronic stress and pregnancy: preterm birth from a biobehavioral perspective. *Journal of Midwifery and Women's Health* 54:8–17, 2009
- Stewart DE, Robertson E, Dennis C-L, et al: Postpartum Depression: Literature Review of Risk Factors and Interventions. Toronto, University Health Network Women's Health Program for Toronto Public Health, 2003
- Da Costa D, Larouche J, Dritsa M, et al: Psychosocial correlates of prepartum and postpartum depressed mood. *Journal of Affective Disorders* 59:31–40, 2000
- Collins NL, Dunkel-Schetter C, Lobel M, et al: Social support in pregnancy: psychosocial correlates of birth outcomes and postpartum depression. *Journal of Personality and Social Psychology* 65:1243–1258, 1993
- Vigod SN, Villegas L, Dennis CL, et al: Prevalence and risk factors for postpartum depression among women with preterm and low-birth-weight infants: a systematic review. *BJOG* 117: 540–550, 2010
- Poehlmann J, Schwichtenberg AJ, Bolt D, et al: Predictors of depressive symptom trajectories in mothers of preterm or low birth weight infants. *Journal of Family Psychology* 23:690–704, 2009
- Robins CJ, Block P: Cognitive theories of depression viewed from a diathesis-stress perspective: evaluations of the models of Beck and of Abramson, Seligman, and Teasdale. *Cognitive Therapy and Research* 13:297–313, 1989
- Monroe SM, Simons AD: Diathesis-stress theories in the context of life stress research: implications for the depressive disorders. *Psychological Bulletin* 110:406–425, 1991
- Abela JR, Aydin C, Auerbach RP: Operationalizing the “vulnerability” and “stress” components of the hopelessness theory of depression: a multi-wave longitudinal study. *Behaviour Research and Therapy* 44:1565–1583, 2006
- Graff LA, Dyck DG, Schallow JR: Predicting postpartum depressive symptoms: a structural modeling analysis. *Perceptual and Motor Skills* 73:1137–1138, 1991
- Rich-Edwards JW, Kleinman K, Abrams A, et al: Sociodemographic predictors of antenatal and postpartum depressive symptoms among women in a medical group practice. *Journal of Epidemiology and Community Health* 60:221–227, 2006
- Arizmendi TG, Affonso DD: Stressful events related to pregnancy and postpartum. *Journal of Psychosomatic Research* 31:743–756, 1987
- Bhutta AT, Cleves MA, Casey PH, et al: Cognitive and behavioral outcomes of school-aged children who were born preterm: a meta-analysis. *JAMA* 288:728–737, 2002
- Hack M, Taylor HG, Schluchter M, et al: Behavioral outcomes of extremely low birth weight children at age 8 years. *Journal of Developmental and Behavioral Pediatrics* 30:122–130, 2009
- Weinberg MK, Tronick EZ: The impact of maternal psychiatric illness on infant development. *Journal of Clinical Psychiatry* 59(suppl 2):53–61, 1998
- Korja R, Savonlahti E, Ahlqvist-Björkroth S, et al: Maternal depression is associated with mother-infant interaction in preterm infants. *Acta Paediatrica* 97:724–730, 2008
- Agostini F, Neri E, Dellabartola S, et al: Early interactive behaviours in preterm infants and their mothers: influences of maternal depressive symptomatology and neonatal birth weight. *Infant Behavior and Development* 37:86–93, 2014
- Melnyk BM, Crean HF, Feinstein NF, et al: Maternal anxiety and depression after a premature infant's discharge from the neonatal intensive care unit: explanatory effects of the creating opportunities for parent empowerment program. *Nursing Research* 57:383–394, 2008
- Zelkowitz P, Feeley N, Shrier I, et al: The Cues and Care Trial: a randomized controlled trial of an intervention to reduce maternal anxiety and improve developmental outcomes in very low birth-weight infants. *BMC Pediatrics* 8:38, 2008
- Newnham CA, Milgrom J, Skouteris H: Effectiveness of a modified Mother-Infant Transaction Program on outcomes for preterm infants from 3 to 24 months of age. *Infant Behavior and Development* 32:17–26, 2009
- Ravn IH, Smith L, Smeby NA, et al: Effects of early mother-infant intervention on outcomes in mothers and moderately and late preterm infants at age 1 year: a randomized controlled trial. *Infant Behavior and Development* 35:36–47, 2012
- Hunker DF, Patrick TE, Albrecht SA, et al: Is difficult childbirth related to postpartum maternal outcomes in the early postpartum period? *Archives of Women's Mental Health* 12:211–219, 2009
- Murdoch MR, Franck LS: Gaining confidence and perspective: a phenomenological study of mothers' lived experiences caring for infants at home after neonatal unit discharge. *Journal of Advanced Nursing* 68:2008–2020, 2012
- Shaw E, Levitt C, Wong S, et al: Systematic review of the literature on postpartum care: effectiveness of postpartum support to improve maternal parenting, mental health, quality of life, and physical health. *Birth* 33:210–220, 2006
- Lau R, Morse CA: Parents' coping in the neonatal intensive care unit: a theoretical framework. *Journal of Psychosomatic Obstetrics and Gynaecology* 22:41–47, 2001
- Cutrona CE, Troutman BR: Social support, infant temperament, and parenting self-efficacy: a mediational model of postpartum depression. *Child Development* 57:1507–1518, 1986
- Grote NK, Katon WJ, Russo JE, et al: Collaborative care for perinatal depression in socioeconomically disadvantaged women: a randomized controlled trial. *Depression and Anxiety* 32:821–834, 2015
- Spitzer RL, Williams JB, Kroenke K, et al: Validity and utility of the PRIME-MD patient health questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *American Journal of Obstetrics and Gynecology* 183:759–769, 2000
- Sheehan DV, Lecrubier Y, Sheehan KH, et al: The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 59(suppl 20): 22–33, 1998
- Derogatis LR: SCL-90-R: Symptom Checklist-90-R: Administration, Scoring, and Procedures Manual. San Antonio, Tex, NCS Pearson, 1996
- Grote NK, Katon WJ, Lohr MJ, et al: Culturally relevant treatment services for perinatal depression in socio-economically disadvantaged women: the design of the MOMCare study. *Contemporary Clinical Trials* 39:34–49, 2014
- Schmitz N, Hartkamp N, Franke GH: Assessing clinically significant change: application to the SCL-90-R. *Psychological Reports* 86:263–274, 2000

37. Katon W, Robinson P, Von Korff M, et al: A multifaceted intervention to improve treatment of depression in primary care. *Archives of General Psychiatry* 53:924–932, 1996
38. Unützer J, Katon W, Callahan CM, et al: Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA* 288:2836–2845, 2002
39. Mundt JC, Marks IM, Shear MK, et al: The Work and Social Adjustment Scale: a simple measure of impairment in functioning. *British Journal of Psychiatry* 180:461–464, 2002
40. O'Hara MW, Varner MW, Johnson SR: Assessing stressful life events associated with childbearing: the Peripartum Events Scale. *Journal of Reproductive and Infant Psychology* 4:85–98, 1986
41. Blanchard EB, Jones-Alexander J, Buckley TC, et al: Psychometric properties of the PTSD Checklist (PCL). *Behaviour Research and Therapy* 34:669–673, 1996
42. Dennis CL, Janssen PA, Singer J: Identifying women at-risk for postpartum depression in the immediate postpartum period. *Acta Psychiatrica Scandinavica* 110:338–346, 2004
43. Mayberry LJ, Horowitz JA, Declercq E: Depression symptom prevalence and demographic risk factors among US women during the first 2 years postpartum. *Journal of Obstetric, Gynecologic, and neonatal nursing* 36:542–549, 2007
44. Miles MS, Holditch-Davis D, Schwartz TA, et al: Depressive symptoms in mothers of prematurely born infants. *Journal of Developmental and Behavioral Pediatrics* 28:36–44, 2007
45. Logsdon MC, Wisner K, Sit D, et al: Depression treatment and maternal functioning. *Depression and Anxiety* 28:1020–1026, 2011
46. Burger J, Horwitz SM, Forsyth BWC, et al: Psychological sequelae of medical complications during pregnancy. *Pediatrics* 91:566–571, 1993
47. Beck CT: The effects of postpartum depression on maternal-infant interaction: a meta-analysis. *Nursing Research* 44:298–304, 1995
48. Betts KS, Williams GM, Najman JM, et al: Maternal depressive, anxious, and stress symptoms during pregnancy predict internalizing problems in adolescence. *Depression and Anxiety* 31:9–18, 2014
49. Grote NK, Katon WJ, Russo JE, et al: A randomized trial of collaborative care for perinatal depression in socio-economically disadvantaged women: the impact of comorbid PTSD. *Journal of Clinical Psychiatry*, in press
50. O'Connor TG, Heron J, Golding J, et al: Maternal antenatal anxiety and children's behavioural/emotional problems at 4 years: report from the Avon Longitudinal Study of Parents and Children. *British Journal of Psychiatry* 180:502–508, 2002
51. Martin RP, Noyes J, Wisenbaker J, et al: Prediction of early childhood negative emotionality and inhibition from maternal distress during pregnancy. *Merrill-Palmer Quarterly* 45:370–391, 1999
52. Hamilton B, Martin J, Osterman M, et al: Births: preliminary data for 2014; in *National Vital Statistics Reports*. Hyattsville, Md, National Center for Health Statistics, 2015
53. Mulcahy R, Reay RE, Wilkinson RB, et al: A randomised control trial for the effectiveness of group Interpersonal Psychotherapy for postnatal depression. *Archives of Women's Mental Health* 13: 125–139, 2010