



NORTHWEST AIDS EDUCATION AND TRAINING CENTER

2012 CROI Update: Selected Studies

David Spach, MD

Clinical Director, Northwest AETC

Professor of Medicine, Division of Infectious Diseases

University of Washington

Presentation Prepared by:

David Spach, MD

Last Updated: March 27, 2012

2012 CROI Update: Selected Studies

- Telaprevir in HCV Treatment Naïve
- Drug Interactions with Boceprevir and PIs
- Boceprevir in HCV Treatment Naïve
- Quad Pill versus Atripla in Treatment Naïve
- Dolutegravir in Treatment Naïve

Telaprevir in HCV Treatment Naive Study 110

Telaprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection

Study 110: Design

Protocol Features for Study 110

- Phase 2a trial; randomized, placebo-controlled
- Chronic HCV; Genotype 1; HCV- treatment naïve
- Randomized to Telaprevir + PegIFN+ RBV versus PegIFN + RBV
 - Part A: No Antiretroviral Rx
 - Part B: Antiretroviral Rx
- Antiretroviral Regimens in Part B:
 - (1) Tenofovir-Emtricitabine-Efavirenz
 - (2) Tenofovir + (Emtricitabine or Lamivudine) + Ritonavir + Atazanavir
- Drug Dosing: Telaprevir = 750 mg tid (1125 mg tid with Efavirenz)

Telaprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection Study 110: Design


		Week	0	12	48	60	72
						SVR 12	SVR 24
Part A No Current ART	N=7	T12 PR48	Telaprevir				
			Peginterferon + Ribavirin				
	N=6	PR48	Placebo				
			Peginterferon + Ribavirin				
Part B Stable ART	N=31	T12 PR48	Telaprevir				
			Peginterferon + Ribavirin				
	N=16	PR48	Placebo				
			Peginterferon + Ribavirin				

Source: Dieterich D, et al. 19th CROI. 2012; Abstract 46.




Telaprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection Study 110: Design

Week 12 Post Treatment (SVR-12)



Telaprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection Study 110: Design

Week 12 Post Treatment (SVR-12)



Boceprevir Interactions with Ritonavir-Boosted PIs

Interaction Between Boceprevir and Ritonavir-Boosted Protease Inhibitors in Healthy Volunteers

Impact of Boceprevir on Levels of Ritonavir-Boosted Protease Inhibitors

Protease Inhibitor	Cmin (trough)	Mean AUC	Cmax (peak)
Atazanavir	↓ 49%	↓ 35%	↓ 25%
Darunavir	↓ 59%	↓ 44%	↓ 36%
Lopinavir	↓ 43%	↓ 34%	↓ 30%



[Home](#)

[Food](#)

[Drugs](#)

[Medical Devices](#)

[Vaccines, Blood & Biologics](#)

[Animal & Veterinary](#)

[Cosmetics](#)

[Radiation-Emitting Products](#)

[Tobacco](#)

For Consumers

[Home](#) [For Consumers](#) [Consumer Information by Audience](#) [For Patients and Patient Advocates](#)

Consumer Information by Audience

[For Patients and Patient Advocates](#)

[Hepatitis B and C](#)

[Patient Involvement](#)

[Speeding Access to Important New Therapies](#)

[Participating in Clinical Trials](#)

[Access to Investigational Drugs](#)

[HIV and AIDS Activities](#)

[Cancer Liaison Program](#)

Victrelis (boceprevir) and Ritonavir-Boosted HIV Protease Inhibitor Drugs - Drug Interactions

The U.S. Food and Drug Administration (FDA) is notifying healthcare professionals and patients that drug interactions between the hepatitis C virus (HCV) protease inhibitor Victrelis (boceprevir) and certain ritonavir-boosted human immunodeficiency virus (HIV) protease inhibitors (atazanavir, lopinavir, darunavir) can potentially reduce the effectiveness of these medicines when they are used together.

Patients should not stop taking any of their medicines without talking to their healthcare professional. Patients should contact their healthcare professional if they have any questions or concerns.

Healthcare professionals who have started patients infected with both chronic HCV and HIV on Victrelis and antiretroviral therapy containing a ritonavir-boosted protease inhibitor should closely monitor patients for HCV treatment response and for potential HCV and HIV virologic rebound.

A drug interaction study showed that taking boceprevir (Victrelis) with ritonavir (Norvir) in combination with atazanavir (Reyataz) or darunavir (Prezista), or with Kaletra (lopinavir/ritonavir) reduced the blood levels of the HIV medicines and boceprevir in the body (see Data Summary below). FDA will be updating the Victrelis drug label to include information about these drug interactions.

Merck and Company has issued a [Dear Healthcare Professional letter \(PDF - 67KB\)](#) with information about this drug interaction study.

Facts about Victrelis (boceprevir) and HIV protease inhibitors

- Victrelis is a hepatitis C virus (HCV) protease inhibitor used with the medicines peginterferon alfa and ribavirin to treat chronic (long-lasting) hepatitis C infection in adults who have not been treated before or who have failed previous treatment.
- HIV protease inhibitors are a class of anti-viral drugs used to treat HIV infection.
- Ritonavir is an HIV protease inhibitor used to "boost" other HIV protease inhibitors, increasing their levels in the blood and making them more effective.



Boceprevir in HCV Treatment Naïve



Boceprevir with Antiretroviral Therapy Study : Design

Protocol Features

- N = 100 HIV/HCV coinfecting, Age 18-65
- Phase 2a trial; randomized, placebo-controlled
- Chronic HCV; Genotype 1; HCV- treatment naïve
- CD4 \geq 200 cells/mm³; HIV RNA < 50 copies/ml on ARV therapy
- Some ARVs Not Allowed: NNRTI, unboosted PI, ZDV, d4T
- Randomized: (1:2) and all had 4-week lead in with PegIFN+ Ribavirin
 - (1x) Peginterferon + Ribavirin (n = 34)
 - (2x) Boceprevir + Peginterferon + Ribavirin (n = 64)


Boceprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection Study: Design

Week	0	4	48	60	72
	Lead In			SVR 12	SVR 24



Boceprevir plus Peginterferon/Ribavirin in HIV/HCV Coinfection Study: Results

Treatment Response



HIV Breakthrough Viremia

- HIV Breakthrough Viremia (HIV RNA > 50 copies/ml on 2 consecutive visits)
 - Overall: 7 (7%) of 98 patients
 - Control Group (PR): 4 (12%) of 34
 - Boceprevir Group (B + PR): 3 (5%) of 64

Elvitegravir in “Quad Pill” versus *Atripla* Study 102

Elvitegravir-Cobicistat-TDF-FTC versus Efavirenz-TDF-FTC

Study 102: Design

Study Design

Protocol

- N = 700 HIV-infected Adults
- Randomized, Double-blind, Phase 3 trial
- Antiretroviral naive
- No baseline NRTI, NNRTI mutations
- HIV RNA \geq 5,000 copies/ml
- Creatinine Clearance \geq 70 ml/min

***Elvitegravir-Cobicistat-
Tenofovir-Emtricitabine**

(n = 348)

**^Efavirenz-
Tenofovir-Emtricitabine**

(n = 352)


*Dosing: Elvitegravir (150 mg); Cobicistat (150 mg); Tenofovir (300 mg); Emtricitabine (200mg)

^Dosing: Efavirenz (600 mg); Tenofovir (300 mg); Emtricitabine (200mg)

Elvitegravir-Cobicistat-TDF-FTC versus Efavirenz-TDF-FTC

Study 102: Results

Week 48: Virologic Response (ITT-TLOVR*)




*ITT-TLOVR = Intention to Treat-Time to Loss of Virologic Response

Source: Sax P, et al. 19th CROI. 2012; Abstract 101.

Dolutegravir (“572”) vs. Efavirenz
SPRING-1 Study: 96 Week Data


Dolutegravir vs. Efavirenz in ARV-Naive SPRING-1: Study Design



*2 NRTIs: Tenofovir-Emtricitabine + (67%); Abacavir-Lamivudine (33%)

Dolutegravir vs. Efavirenz in ARV-Naive SPRING-1: Study Results

96 Week Data: Virologic Response (TLOVR)



All regimens included 2 NRTIs: Tenofovir-Emtricitabine or Abacavir-Lamivudine

End
