

- Alptekin, K., Akvardar, Y., Kivircik Akdede, B. B., Dumlu, K., Isik, D., Pirincci, F., et al. (2005). Is quality of life associated with cognitive impairment in schizophrenia? *Prog Neuropsychopharmacol Biol Psychiatry*, 29(2), 239-244.
- Amir, M., & Lev-Wiesel, R. (2003). Time does not heal all wounds: Quality of life and psychological distress of people who survived the holocaust as children 55 years later. *J Trauma Stress*, 16(3), 295-299.
- Amir, M., Lewin-Epstein, N., Becker, G., & Buskila, D. (2002). Psychometric properties of the sf-12 (hebrew version) in a primary care population in israel. *Med Care*, 40(10), 918-928.
- Awadalla, A. W., Ohaeri, J. U., Salih, A. A., & Tawfiq, A. M. (2005). Subjective quality of life of community living sudanese psychiatric patients: Comparison with family caregivers' impressions and control group. *Qual Life Res*, 14(8), 1855-1867.
- Bedi, G. S., Gupta, N., Handa, R., Pal, H., & Pandey, R. M. (2005). Quality of life in indian patients with rheumatoid arthritis. *Qual Life Res*, 14(8), 1953-1958.
- Berlim, M. T., Pavanello, D. P., Caldieraro, M. A., & Fleck, M. P. (2005). Reliability and validity of the whoqol bref in a sample of brazilian outpatients with major depression. *Qual Life Res*, 14(2), 561-564.
- Bittencourt, Z. Z., Alves Filho, G., Mazzali, M., & Santos, N. R. (2004). [quality of life in renal transplant patients: Impact of a functioning graft]. *Rev Saude Publica*, 38(5), 732-734.
- Bolsche, F., Hasenbein, U., Reissberg, H., Schlotte, A., & Wallesch, C. W. (2003). [results of in- vs outpatient post-stroke rehabilitation over 6 months]. *Fortschr Neurol Psychiatr*, 71(9), 458-468.
- Bowman, S. J., Booth, D. A., & Platts, R. G. (2004). Measurement of fatigue and discomfort in primary sjogren's syndrome using a new questionnaire tool. *Rheumatology (Oxford)*, 43(6), 758-764.
- Brieger, P., Rottig, S., & Marneros, A. (2004). [quality of life in unipolar depressive and bipolar affective patients]. *Psychiatr Prax*, 31(6), 304-309.
- Cankurtaran, E. S., Ulug, B., Saygi, S., Tiryaki, A., & Akalan, N. (2005). Psychiatric morbidity, quality of life, and disability in mesial temporal lobe epilepsy patients before and after anterior temporal lobectomy. *Epilepsy Behav*, 7(1), 116-122.
- Chan, G. W., Ungvari, G. S., Shek, D. T., & Leung Dagger, J. J. (2003). Hospital and community-based care for patients with chronic schizophrenia in hong kong--quality of life and its correlates. *Soc Psychiatry Psychiatr Epidemiol*, 38(4), 196-203.

- Chand, P. K., Mattoo, S. K., & Sharan, P. (2004). Quality of life and its correlates in patients with bipolar disorder stabilized on lithium prophylaxis. *Psychiatry Clin Neurosci*, 58(3), 311-318.
- Chandra, P. S., Deepthivarma, S., Jairam, K. R., & Thomas, T. (2003). Relationship of psychological morbidity and quality of life to illness-related disclosure among hiv-infected persons. *J Psychosom Res*, 54(3), 199-203.
- Chapin, M. H., Miller, S. M., Ferrin, J. M., Chan, F., & Rubin, S. E. (2004). Psychometric validation of a subjective well-being measure for people with spinal cord injuries. *Disabil Rehabil*, 26(19), 1135-1142.
- Chopra, P. K., Couper, J. W., & Herrman, H. (2004). The assessment of patients with long-term psychotic disorders: Application of the who disability assessment schedule ii. *Aust NZ J Psychiatry*, 38(9), 753-759.
- Cieza, A., & Stucki, G. (2005). Content comparison of health-related quality of life (hrqol) instruments based on the international classification of functioning, disability and health (icf). *Qual Life Res*, 14(5), 1225-1237.
- Clayton, W. J., Lipton, M., Elford, J., Rustin, M., & Sherr, L. (2005). A randomized controlled trial of laser treatment among hirsute women with polycystic ovary syndrome. *Br J Dermatol*, 152(5), 986-992.
- da Silva Lima, A. F., Fleck, M., Pechansky, F., de Boni, R., & Sukop, P. (2005). Psychometric properties of the world health organization quality of life instrument (whoqol-bref) in alcoholic males: A pilot study. *Qual Life Res*, 14(2), 473-478.
- Development of the world health organization whoqol-bref quality of life assessment. The whoqol group. (1998). *Psychol Med*, 28(3), 551-558.
- Dogan, S., Dogan, O., Tel, H., Coker, F., Polatoz, O., & Dogan, F. B. (2004). Psychosocial approaches in outpatients with schizophrenia. *Psychiatr Rehabil J*, 27(3), 279-282.
- Edwards, B., & O'Connell, B. (2003). Internal consistency and validity of the stroke impact scale 2.0 (sis 2.0) and sis-16 in an australian sample. *Qual Life Res*, 12(8), 1127-1135.
- Eryilmaz, M. M., Ozdemir, C., Yurtman, F., Cilli, A., & Karaman, T. (2005). Quality of sleep and quality of life in renal transplantation patients. *Transplant Proc*, 37(5), 2072-2076.
- Fang, C. T., Hsiung, P. C., Yu, C. F., Chen, M. Y., & Wang, J. D. (2002). Validation of the world health organization quality of life instrument in patients with hiv infection. *Qual Life Res*, 11(8), 753-762.

- Fellinger, J., Holzinger, D., Dobner, U., Gerich, J., Lehner, R., Lenz, G., et al. (2005). Mental distress and quality of life in a deaf population. *Soc Psychiatry Psychiatr Epidemiol*, 40(9), 737-742.
- Fleck, M. P., Louzada, S., Xavier, M., Chachamovich, E., Vieira, G., Santos, L., et al. (2000). [application of the portuguese version of the abbreviated instrument of quality life whoqol-bref]. *Rev Saude Publica*, 34(2), 178-183.
- Gazalle, F. K., Andreazza, A. C., Cereser, K. M., Hallal, P. C., Santin, A., & Kapczinski, F. (2005). Clinical impact of late diagnose of bipolar disorder. *J Affect Disord*, 86(2-3), 313-316.
- Golimbet, V., & Trubnikov, V. (2001). Evaluation of the dementia carers situation in russia. *Int J Geriatr Psychiatry*, 16(1), 94-99.
- Gorna, K., Jaracz, K., & Rybakowski, F. (2005). Objective and subjective quality of life in schizophrenic patients after a first hospitalization. *Rocz Akad Med Bialymst*, 50 Suppl 1, 225-227.
- Haltbakk, J., Hanestad, B. R., & Hunskaar, S. (2005a). How important are men's lower urinary tract symptoms (luts) and their impact on the quality of life (qol)? *Qual Life Res*, 14(7), 1733-1741.
- Haltbakk, J., Hanestad, B. R., & Hunskaar, S. (2005b). Relevance and variability of the severity of incontinence, and increased daytime and night-time voiding frequency, associated with quality of life in men with lower urinary tract symptoms. *BJU Int*, 96(1), 83-87.
- Hanestad, B. R., Rustoen, T., Knudsen, O., Jr., Lerdal, A., & Wahl, A. K. (2004). Psychometric properties of the whoqol-bref questionnaire for the norwegian general population. *J Nurs Meas*, 12(2), 147-159.
- Hasanah, C. I., Naing, L., & Rahman, A. R. (2003). World health organization quality of life assessment: Brief version in bahasa malaysia. *Med J Malaysia*, 58(1), 79-88.
- Heald, A. H., Ghosh, S., Bray, S., Gibson, C., Anderson, S. G., Buckler, H., et al. (2004). Long-term negative impact on quality of life in patients with successfully treated cushing's disease. *Clin Endocrinol (Oxf)*, 61(4), 458-465.
- Herrman, H., Hawthorne, G., & Thomas, R. (2002). Quality of life assessment in people living with psychosis. *Soc Psychiatry Psychiatr Epidemiol*, 37(11), 510-518.
- Hinterseher, I., Saeger, H. D., Koch, R., Bloomenthal, A., Ockert, D., & Bergert, H. (2004). Quality of life and long-term results after ruptured abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg*, 28(3), 262-269.

- Horng, Y. S., Hwang, Y. H., Wu, H. C., Liang, H. W., Mhe, Y. J., Twu, F. C., et al. (2005). Predicting health-related quality of life in patients with low back pain. *Spine, 30*(5), 551-555.
- Hsiung, P. C., Fang, C. T., Chang, Y. Y., Chen, M. Y., & Wang, J. D. (2005). Comparison of whoqol-bref and sf-36 in patients with hiv infection. *Qual Life Res, 14*(1), 141-150.
- Hsu, C., Wang, J. D., Hwang, J. S., Tien, H. F., Chang, S. M., Cheng, A. L., et al. (2003). Survival-weighted health profile for long-term survivors of acute myelogenous leukemia. *Qual Life Res, 12*(5), 503-517.
- Hwang, H. F., Liang, W. M., Chiu, Y. N., & Lin, M. R. (2003). Suitability of the whoqol-bref for community-dwelling older people in taiwan. *Age Ageing, 32*(6), 593-600.
- Izutsu, T., Tsutsumi, A., Islam, A., Matsuo, Y., Yamada, H. S., Kurita, H., et al. (2005). Validity and reliability of the bangla version of whoqol-bref on an adolescent population in bangladesh. *Qual Life Res, 14*(7), 1783-1789.
- Jang, Y., Hsieh, C. L., Wang, Y. H., & Wu, Y. H. (2004). A validity study of the whoqol-bref assessment in persons with traumatic spinal cord injury. *Arch Phys Med Rehabil, 85*(11), 1890-1895.
- Kao, S., Lai, K. L., Lin, H. C., Lee, H. S., & Wen, H. C. (2005). Whoqol-bref as predictors of mortality: A two-year follow-up study at veteran homes. *Qual Life Res, 14*(6), 1443-1454.
- Khanna, S., Pal, H., Pandey, R. M., & Handa, R. (2004). The relationship between disease activity and quality of life in systemic lupus erythematosus. *Rheumatology (Oxford), 43*(12), 1536-1540.
- Kilian, R., Matschinger, H., Loeffler, W., Roick, C., & Angermeyer, M. C. (2002). A comparison of methods to handle skew distributed cost variables in the analysis of the resource consumption in schizophrenia treatment. *J Ment Health Policy Econ, 5*(1), 21-31.
- Kim, D. Y., Lee, S. H., Lee, H. Y., Lee, H. J., Chang, S. B., Chung, S. K., et al. (2005). Validation of the korean version of the oswestry disability index. *Spine, 30*(5), E123-127.
- Kuehner, C., & Buerger, C. (2005). Determinants of subjective quality of life in depressed patients: The role of self-esteem, response styles, and social support. *J Affect Disord, 86*(2-3), 205-213.
- Lai, K. L., Tzeng, R. J., Wang, B. L., Lee, H. S., Amidon, R. L., & Kao, S. (2005). Health-related quality of life and health utility for the institutional elderly in taiwan. *Qual Life Res, 14*(4), 1169-1180.

- Lapinski, T., Bien, B., & Wilmanska, J. (2004). [health-related quality of life in the elderly after coronary artery bypass grafting]. *Przegl Lek*, 61(12), 1351-1355.
- Law, C. W., Chen, E. Y., Cheung, E. F., Chan, R. C., Wong, J. G., Lam, C. L., et al. (2005). Impact of untreated psychosis on quality of life in patients with first-episode schizophrenia. *Qual Life Res*, 14(8), 1803-1811.
- Leung, K. F., Wong, W. W., Tay, M. S., Chu, M. M., & Ng, S. S. (2005). Development and validation of the interview version of the hong kong chinese whoqol-bref. *Qual Life Res*, 14(5), 1413-1419.
- Lin, M. R., Huang, W., Huang, C., Hwang, H. F., Tsai, L. W., & Chiu, Y. N. (2002). The impact of the chi-chi earthquake on quality of life among elderly survivors in taiwan--a before and after study. *Qual Life Res*, 11(4), 379-388.
- Liou, H. H., Chen, R. C., Chen, C. C., Chiu, M. J., Chang, Y. Y., & Wang, J. D. (2005). Health related quality of life in adult patients with epilepsy compared with a general reference population in taiwan. *Epilepsy Res*, 64(3), 151-159.
- Lo Coco, G., Lo Coco, D., Cicero, V., Oliveri, A., Lo Verso, G., Piccoli, F., et al. (2005). Individual and health-related quality of life assessment in amyotrophic lateral sclerosis patients and their caregivers. *J Neurol Sci*, 238(1-2), 11-17.
- Lung, F. W., Huang, Y. L., Shu, B. C., & Lee, F. Y. (2004). Parental rearing style, premorbid personality, mental health, and quality of life in chronic regional pain: A causal analysis. *Compr Psychiatry*, 45(3), 206-212.
- Michelone Ade, P., & Santos, V. L. (2004). [quality of life of cancer patients with and without an ostomy]. *Rev Lat Am Enfermagem*, 12(6), 875-883.
- Min, S. K., Kim, K. I., Lee, C. I., Jung, Y. C., Suh, S. Y., & Kim, D. K. (2002). Development of the korean versions of who quality of life scale and whoqol-bref. *Qual Life Res*, 11(6), 593-600.
- Mond, J. M., Hay, P. J., Rodgers, B., Owen, C., & Beumont, P. J. (2005). Assessing quality of life in eating disorder patients. *Qual Life Res*, 14(1), 171-178.
- Mooney, M., Hannon, F., Barry, M., Friel, S., & Kelleher, C. (2002). Perceived quality of life and mental health status of irish female prisoners. *Ir Med J*, 95(8), 241-243.
- Muller, K., Schwesig, R., Leuchte, S., & Riede, D. (2001). [coordinative treatment and quality of life - a randomised trial of nurses with back pain]. *Gesundheitswesen*, 63(10), 609-618.
- Nasermoaddeli, A., Sekine, M., Hamanishi, S., & Kagamimori, S. (2003). Associations between sense of coherence and psychological work characteristics with changes in quality of life in japanese civil servants: A 1-year follow-up study. *Ind Health*, 41(3), 236-241.

- Naumann, V. J., & Byrne, G. J. (2004). Whoqol-bref as a measure of quality of life in older patients with depression. *Int Psychogeriatr*, 16(2), 159-173.
- Niu, S. F., & Li, I. C. (2005). Quality of life of patients having renal replacement therapy. *J Adv Nurs*, 51(1), 15-21.
- Noerholm, V., Groenvold, M., Watt, T., Bjorner, J. B., Rasmussen, N. A., & Bech, P. (2004). Quality of life in the danish general population--normative data and validity of whoqol-bref using rasch and item response theory models. *Qual Life Res*, 13(2), 531-540.
- Norholm, V., & Bech, P. (2001). The who quality of life (whoqol) questionnaire: Danish validation study. *Nord J Psychiatry*, 55(4), 229-235.
- O'Carroll, R. E., Smith, K., Couston, M., Cossar, J. A., & Hayes, P. C. (2000). A comparison of the whoqol-100 and the whoqol-bref in detecting change in quality of life following liver transplantation. *Qual Life Res*, 9(1), 121-124.
- Ogunseitan, O. A. (2005). Topophilia and the quality of life. *Environ Health Perspect*, 113(2), 143-148.
- Ohaeri, J. U., Olusina, A. K., & Al-Abassi, A. H. (2004). Factor analytical study of the short version of the world health organization quality of life instrument. *Psychopathology*, 37(5), 242-248.
- Phunggrassami, T., Katikarn, R., Watanaarepornchai, S., & Sangtawan, D. (2004). Quality of life assessment in radiotherapy patients by whoqol-bref-thai: A feasibility study. *J Med Assoc Thai*, 87(12), 1459-1465.
- Radosevic-Vidacek, B., Macan, J., & Koscec, A. (2004). [stress and allergy]. *Arh Hig Rada Toksikol*, 55(2-3), 205-211.
- Rose, M., Kohler, K., Kohler, F., Sawitzky, B., Fliege, H., & Klapp, B. F. (2005). Determinants of the quality of life of patients with congenital heart disease. *Qual Life Res*, 14(1), 35-43.
- Ruesch, P., Graf, J., Meyer, P. C., Rossler, W., & Hell, D. (2004). Occupation, social support and quality of life in persons with schizophrenic or affective disorders. *Soc Psychiatry Psychiatr Epidemiol*, 39(9), 686-694.
- Saupe, R. S., Nietche, E. A., Cestari, M. E., Giorgi, M. D., & Krahl, M. (2004). [quality of life of nursing students]. *Rev Lat Am Enfermagem*, 12(4), 636-642.
- Saxena, S., Carlson, D., & Billington, R. (2001). The who quality of life assessment instrument (whoqol-bref): The importance of its items for cross-cultural research. *Qual Life Res*, 10(8), 711-721.

- Saxena, S., Chandiramani, K., & Bhargava, R. (1998). Whoqol-hindi: A questionnaire for assessing quality of life in health care settings in india. World health organization quality of life. *Natl Med J India*, 11(4), 160-165.
- Schmidt, S., Muhlan, H., & Power, M. (2005). The eurohis-qol 8-item index: Psychometric results of a cross-cultural field study. *Eur J Public Health*.
- Sim, K., Mahendran, R., Siris, S. G., Heckers, S., & Chong, S. A. (2004). Subjective quality of life in first episode schizophrenia spectrum disorders with comorbid depression. *Psychiatry Res*, 129(2), 141-147.
- Skevington, S. M., Lotfy, M., & O'Connell, K. A. (2004). The world health organization's whoqol-bref quality of life assessment: Psychometric properties and results of the international field trial. A report from the whoqol group. *Qual Life Res*, 13(2), 299-310.
- Taylor, W. J., Myers, J., Simpson, R. T., McPherson, K. M., & Weatherall, M. (2004). Quality of life of people with rheumatoid arthritis as measured by the world health organization quality of life instrument, short form (whoqol-bref): Score distributions and psychometric properties. *Arthritis Rheum*, 51(3), 350-357.
- Toprak, A. B., Eser, E., Guler, C., Baser, F. E., & Mayali, H. (2005). Cross-validation of the turkish version of the 25-item national eye institute visual functioning questionnaire (nei-vfq 25). *Ophthalmic Epidemiol*, 12(4), 259-269.
- Trompenaars, F. J., Masthoff, E. D., Van Heck, G. L., Hodiamont, P. P., & De Vries, J. (2005a). Content validity, construct validity, and reliability of the whoqol-bref in a population of dutch adult psychiatric outpatients. *Qual Life Res*, 14(1), 151-160.
- Trompenaars, F. J., Masthoff, E. D., Van Heck, G. L., Hodiamont, P. P., & De Vries, J. (2005b). Relationships between demographic variables and quality of life in a population of dutch adult psychiatric outpatients. *Soc Psychiatry Psychiatr Epidemiol*, 40(7), 588-594.
- Tzeng, D. S., Lung, F. W., & Chang, Y. Y. (2004). Comparison of quality of life for people with schizophrenia and mental health of caregivers beteen commn-based and hospital-based services. *Kaohsiung J Med Sci*, 20(9), 443-451.
- Umansky, R., Amir, M., Fridmann, M., Zidon, E., Chen, D., & Nemetz, B. (2003). Was it a good move? Improvement in quality of life among chronic mental patients moving from a mental hospital to a hostel in the community. *Isr J Psychiatry Relat Sci*, 40(4), 248-257.
- van de Willige, G., Wiersma, D., Nienhuis, F. J., & Jenner, J. A. (2005). Changes in quality of life in chronic psychiatric patients: A comparison between euroqol (eq-5d) and whoqol. *Qual Life Res*, 14(2), 441-451.

- Wang, X., Gao, L., Zhang, H., Zhao, C., Shen, Y., & Shinfuku, N. (2000). Post-earthquake quality of life and psychological well-being: Longitudinal evaluation in a rural community sample in northern china. *Psychiatry Clin Neurosci*, 54(4), 427-433.
- Weatherall, M., McPherson, K., Taylor, W., & Simpson, R. (2004). Avoiding pitfalls of correlation coefficients in the assessment of measurement instruments in rehabilitation research. *Clin Rehabil*, 18(2), 186-194.
- Wegener, S., Redoblado-Hodge, M. A., Lucas, S., Fitzgerald, D., Harris, A., & Brennan, J. (2005). Relative contributions of psychiatric symptoms and neuropsychological functioning to quality of life in first-episode psychosis. *Aust N Z J Psychiatry*, 39(6), 487-492.
- Wik, K. E., Lindegaard, K. F., Brønborg, B., Bjørk, I. T., & Ruland, C. (2005). [life following acute subarachnoid haemorrhage]. *Tidsskr Nor Laegeforen*, 125(2), 152-154.
- Wray, J., Al-Ruzzeh, S., Mazrani, W., Nakamura, K., George, S., Ilsley, C., et al. (2004). Quality of life and coping following minimally invasive direct coronary artery bypass (midcab) surgery. *Qual Life Res*, 13(5), 915-924.
- Yang, S. C., Kuo, P. W., Wang, J. D., Lin, M. I., & Su, S. (2005). Quality of life and its determinants of hemodialysis patients in taiwan measured with whoqol-bref(tw). *Am J Kidney Dis*, 46(4), 635-641.
- Yao, G., Chung, C. W., Yu, C. F., & Wang, J. D. (2002). Development and verification of validity and reliability of the whoqol-bref taiwan version. *J Formos Med Assoc*, 101(5), 342-351.
- Yao, G., & Wu, C. H. (2005). Factorial invariance of the whoqol-bref among disease groups. *Qual Life Res*, 14(8), 1881-1888.
- Yen, C. F., Tsai, J. J., Lu, P. L., Chen, Y. H., Chen, T. C., Chen, P. P., et al. (2004). Quality of life and its correlates in hiv/aids male outpatients receiving highly active antiretroviral therapy in taiwan. *Psychiatry Clin Neurosci*, 58(5), 501-506.
- Yu, C. L., Fielding, R., Chan, C. L., Tse, V. K., Choi, P. H., Lau, W. H., et al. (2000). Measuring quality of life of chinese cancer patients: A validation of the chinese version of the functional assessment of cancer therapy-general (fact-g) scale. *Cancer*, 88(7), 1715-1727.