

3. Apply the central limit theorem to the random walk in 4, comp. 4.7. Compare the results with an explicit calculation as above.

Since the total displacement is a sum of identical, independent steps we can apply the central limit theorem to the displacement after r steps. Let X_i represent the i th step, for all X_i we have $\langle X_i \rangle = 0$ and $\langle X_i^2 \rangle = 1$. Let $Y = \sum_{i=1}^r X_i$ and $Z = Y/\sqrt{r}$. Then by the CLT

$$P_Z \approx \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}},$$

which means that we can approximate $p_n(r) = 1/2^r \binom{r}{\frac{r-n}{2}}$ by

$$P_Y \approx \frac{1}{\sqrt{2\pi r}} e^{-\frac{y^2}{2r}}.$$

This approximation allows for displacements y larger than the number of steps, which is not admissible by the construction of the random walk. As r grows large we need to show that this probability,

$$P_Y(|y| > r) = \frac{2}{\sqrt{2\pi r}} \int_r^\infty e^{-\frac{y^2}{2r}},$$

vanishes. By examining that this expression note that this is equivalent to asking the probability of being $r/\sqrt{r} = \sqrt{r}$ standard deviations away from the mean, which vanishes as r grows.

5. A random set of dots on $(0, \infty)$ is constructed according to the following recipe. The probability for the first dot to lie in $(\tau_1, \tau_1 + d\tau_1)$ is $w(\tau_1)d\tau_1$, where w is a given non-negative function with

$$\int_0^\infty w(\tau) d\tau = \frac{1}{\gamma} < 1.$$

The probability density for the second dot is $w(\tau_2 - \tau_1)$ and so on. Calculate Q_s .

The probability of the first event NOT happening is $Q_0 = 1 - \frac{1}{\gamma}$. The probability of only one event happening is the same as the second event not happening and thus $Q_1(\tau_1) = w(\tau_1)(1 - \frac{1}{\gamma})$. Following this line of reasoning and making use of the fact that joint PDFs for independent random variables are simply the products of the individual PDFs we get,

$$Q_s(\tau_1, \tau_2, \dots, \tau_s) = (1 - \frac{1}{\gamma}) \prod_{i=1}^s w(\tau_i - \tau_{i-1}),$$

where $\tau_0 = 0$. It remains to show that 1.2 holds and that Q is indeed a probability density function for random dots and sums to 1. The construction already assumes that $\tau_1 < \tau_2 < \tau_3 \dots$, so

$$\begin{aligned} & Q_0 + \int_{-\infty}^\infty Q_1(\tau_1) d\tau_1 + \int_{-\infty}^\infty \int_{\tau_1}^\infty Q_2(\tau_1, \tau_2) d\tau_2 d\tau_1 + \int_{-\infty}^\infty \int_{\tau_1}^\infty \int_{\tau_2}^\infty Q_3(\tau_1, \tau_2, \tau_3) d\tau_3 d\tau_2 d\tau_1 + \dots \\ &= (1 - \frac{1}{\gamma}) \left(1 + \int_{-\infty}^\infty w(\tau_1) d\tau_1 + \int_{-\infty}^\infty \int_{\tau_1}^\infty w(\tau_1) w(\tau_2 - \tau_1) d\tau_2 d\tau_1 + \int_{-\infty}^\infty \int_{\tau_1}^\infty \int_{\tau_2}^\infty w(\tau_1) w(\tau_2 - \tau_1) w(\tau_3 - \tau_2) d\tau_3 d\tau_2 d\tau_1 + \dots \right) \\ &= (1 - \frac{1}{\gamma}) \sum_{n=0}^\infty \left(\frac{1}{\gamma} \right)^n = (1 - \frac{1}{\gamma}) \frac{1}{1 - \frac{1}{\gamma}} = 1, \end{aligned}$$

as required.