
 CALCULUS & ANALYTIC GEOMETRY I

The Derivative as a Function

Problem. Find the slope of the function's graph at the given point. Then find the equation for the line tangent to the graph there.

1. $y = \frac{x-1}{x+1}, \quad x = 0$


Working Backwards. What derivative is represented by each of the following expressions?

$$\lim_{x \rightarrow 5} \frac{2^x - 32}{x - 5} \qquad \lim_{h \rightarrow 0} \frac{\cos(\pi + h) + 1}{h}$$

Derivative as a function. Up to now, we have looked at the derivative as a *number*. It gives us information about a function at a *point*. But the numerical value of derivative varies from point to point, and these new values can also be considered as the values of a new function—the derivative function—with its own graph. Viewed in this way the derivative is a global object.

All functions and their derivatives are related in the same way.

Function	\leftrightarrow	Derivative
increasing	\leftrightarrow	
decreasing	\leftrightarrow	
horizontal	\leftrightarrow	
steep (rising or falling)	\leftrightarrow	
gradual (rising or falling)	\leftrightarrow	
straight	\leftrightarrow	




Notation. Many ways to denote the derivative of the function $y = f(x)$:

$$f'(y) = y' = \frac{dy}{dx} = \frac{df}{dx} = \frac{d}{dx}f(x) = D(f)(x) = D_x f(x).$$


To indicate the value of the derivative at a specified point $x = a$:

$$f'(a) = \left. \frac{dy}{dx} \right|_{x=a} = \left. \frac{df}{dx} \right|_{x=a} = \left. \frac{d}{dx}f(x) \right|_{x=a}.$$

Illustration. For the graph below, sketch its derivative.



Now make a sketch of a graph which would have the original graph as its derivative.



As time permits: For each of the elementary functions below, first make a rough sketch of the function itself, then, based on the correspondences between functions and their derivatives, sketch the derivative of the function (as best you can.)

1. $f(x) = 3x$


2. $g(x) = 3x + 1$

3. $h(x) = x^2$

4. $i(x) = e^x$

5. $j(x) = \sin x$

Definition. A function f is *differentiable at a* if $f'(a)$ exists. It is *differentiable on an open interval (a, b)* if it is differentiable at every number in the interval. A function is *differentiable* if it is differentiable at every point in its domain.




© Thomson Higher Education

For your consideration

[T/F] If f is differentiable at a , then f is continuous at a .

[T/F] If f is continuous at a , the f is differentiable at a .

Higher Derivatives Since the derivative of a function f gives a new function f' , there is nothing stopping us from analyzing the rate of change of f' , denoted f'' or d^2f/dy^2 . What about the rate of change of f'' ?



© Thomson Higher Education