

1. One process undoes what the other one does. The precise version of this statement is given by the Fundamental Theorem of Calculus. See the statement of this theorem and the paragraph that follows it on page 387.

2. (a)  $g(x) = \int_0^x f(t) dt$ , so  $g(0) = \int_0^0 f(t) dt = 0$ .

$$g(1) = \int_0^1 f(t) dt = \frac{1}{2} \cdot 1 \cdot 1 \quad [\text{area of triangle}] = \frac{1}{2}.$$

$$g(2) = \int_0^2 f(t) dt = \int_0^1 f(t) dt + \int_1^2 f(t) dt \quad [\text{below the } x\text{-axis}] \\ = \frac{1}{2} - \frac{1}{2} \cdot 1 \cdot 1 = 0.$$

$$g(3) = g(2) + \int_2^3 f(t) dt = 0 - \frac{1}{2} \cdot 1 \cdot 1 = -\frac{1}{2}.$$


$$g(4) = g(3) + \int_3^4 f(t) dt = -\frac{1}{2} + \frac{1}{2} \cdot 1 \cdot 1 = 0.$$

$$g(5) = g(4) + \int_4^5 f(t) dt = 0 + 1.5 = 1.5.$$

$$g(6) = g(5) + \int_5^6 f(t) dt = 1.5 + 2.5 = 4.$$

(b)  $g(7) = g(6) + \int_6^7 f(t) dt \approx 4 + 2.2$  [estimate from the graph]  $= 6.2$ .

(c) The answers from part (a) and part (b) indicate that  $g$  has a minimum at  $x = 3$  and a maximum at  $x = 7$ . This makes sense from the graph of  $f$  since we are subtracting area on  $1 < x < 3$  and adding area on  $3 < x < 7$ .



7.  $f(t) = \frac{1}{t^3+1}$  and  $g(x) = \int_1^x \frac{1}{t^3+1} dt$ , so by FTC1,  $g'(x) = f(x) = \frac{1}{x^3+1}$ . Note that the lower limit, 1, could be any real number greater than  $-1$  and not affect this answer.

12.  $G(x) = \int_x^1 \cos \sqrt{t} dt = - \int_1^x \cos \sqrt{t} dt \Rightarrow G'(x) = - \frac{d}{dx} \int_1^x \cos \sqrt{t} dt = - \cos \sqrt{x}$

17. Let  $w = 1 - 3x$ . Then  $\frac{dw}{dx} = -3$ . Also,  $\frac{dy}{dx} = \frac{dy}{dw} \frac{dw}{dx}$ , so

$$y' = \frac{d}{dx} \int_{1-3x}^1 \frac{u^3}{1+u^2} du = \frac{d}{dw} \int_w^1 \frac{u^3}{1+u^2} du \cdot \frac{dw}{dx} = - \frac{d}{dw} \int_1^w \frac{u^3}{1+u^2} du \cdot \frac{dw}{dx} = - \frac{w^3}{1+w^2} (-3) = \frac{3(1-3x)^3}{1+(1-3x)^2}$$

19.  $\int_{-1}^2 (x^3 - 2x) dx = \left[ \frac{x^4}{4} - x^2 \right]_{-1}^2 = \left( \frac{2^4}{4} - 2^2 \right) - \left( \frac{(-1)^4}{4} - (-1)^2 \right) = (4 - 4) - \left( \frac{1}{4} - 1 \right) = 0 - \left( -\frac{3}{4} \right) = \frac{3}{4}$

23.  $\int_0^1 x^{4/5} dx = \left[ \frac{5}{9} x^{9/5} \right]_0^1 = \frac{5}{9} - 0 = \frac{5}{9}$

31.  $\int_0^{\pi/4} \sec^2 t dt = [\tan t]_0^{\pi/4} = \tan \frac{\pi}{4} - \tan 0 = 1 - 0 = 1$

36.  $\int_0^1 10^x dx = \left[ \frac{10^x}{\ln 10} \right]_0^1 = \frac{10}{\ln 10} - \frac{1}{\ln 10} = \frac{9}{\ln 10}$

39.  $\int_{-1}^1 e^{u+1} du = [e^{u+1}]_{-1}^1 = e^2 - e^0 = e^2 - 1$  [or start with  $e^{u+1} = e^u e^1$ ]

43.  $f(x) = x^{-4}$  is not continuous on the interval  $[-2, 1]$ , so FTC2 cannot be applied. In fact,  $f$  has an infinite discontinuity at  $x = 0$ , so  $\int_{-2}^1 x^{-4} dx$  does not exist.