

---

 CALCULUS & ANALYTIC GEOMETRY I
 

---

 The Theorems—Most Especially the MVT
 

---

**The Intermediate Value Theorem.** A function  $y = f(x)$  that is continuous on  $[a, b]$  takes on every value between  $f(a)$  and  $f(b)$ .

**Rolle's Theorem.** If a function  $y = f(x)$  is

- continuous on  $[a, b]$
- differentiable on  $(a, b)$
- $f(a) = f(b)$


then there exists a  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

**The Mean Value Theorem.** If a function  $y = f(x)$  is

- continuous on  $[a, b]$
- differentiable on  $(a, b)$

then there exists a  $c$  in  $(a, b)$  such that  $f'(c) = \frac{f(b) - f(a)}{b - a}$ .


---



**Proof of Rolle's Theorem.**

**Application of Rolle's Theorem.** Show that the function  $f(x) = \frac{1}{1-t} + \sqrt{1+t} - 3.1$  has exactly one real zero in the interval  $(-1, 1)$ .

### Proof of the MVT.



Let  $h(x) = f(x) - (\text{equation of secant line})$ .

**Corollary 1.** If  $f'(x) = 0$  at each point of  $(a, b)$ , then  $f(x)$  is a constant function on the interval  $(a, b)$ . (i.e.  $f(x) = C$  for some real number  $C$ .)

**Corollary 2.** If  $f'(x) = g'(x)$  at each point of  $(a, b)$ , then  $f(x)$  and  $g(x)$  differ by a constant function on the interval  $(a, b)$ . (i.e.  $f(x) = g(x) + C$  for some real number  $C$ .)

### Problems.

1. Find all possible functions  $f(x)$  such that  $f'(x) = x^3$ .
2. Find the function  $g(t)$  such that  $g'(t) = e^{2t}$  and  $g(0) = \frac{3}{2}$ .
3. Suppose the acceleration of a body is measure as  $9.8 \text{ m/s}^2$  and we know that  $v(0) = -3 \text{ m/s}$ ,  $s(0) = 5 \text{ m}$ . Find an equation to describe the body's position at time  $t$ .

## CALCULUS &amp; ANALYTIC GEOMETRY I

## The Derivative Tests

**Question.** How do you know where a function is increasing or decreasing?


**Question.** How can a function behave at a transition point as it changes from increasing to decreasing or vice versa?

**Question.** How does this help us identify local extrema?

Consider a function whose derivative is  $f'(x) = x^{-1/2}(x - 3)$ . Where is the function increasing? decreasing? Identify inputs that give local extremes.

Knowing how quickly the derivative of a function is changing can also give important information about local extremes.

**Definition.** The graph of a differentiable function  $y = f(x)$  is *concave up* on an open interval  $I$  if  $f'$  is increasing on  $I$  and *concave down* if  $f'$  is decreasing on  $I$ .



**Definition.** A point where the graph of a function has a tangent line and where the concavity changes is called a *point of inflection*.

**First Derivative Test.** Suppose that  $c$  is a critical point of a continuous function  $f$ , and that  $f$  is differentiable at every point in some interval containing  $c$  except possibly  $c$  itself. Moving across  $c$  from left to right

- if  $f'$  changes from negative to positive at  $c$ , the  $f$  has a local \_\_\_\_\_ at  $c$ ;
- if  $f'$  changes from positive to negative at  $c$ , the  $f$  has a local \_\_\_\_\_ at  $c$ ;
- if  $f'$  does not change sign at  $c$ , then  $f$  has no local extremum at  $c$ .

**Second Derivative Test.** Suppose  $f''$  is continuous on an open interval that contains  $x = v$ .

- If  $f'(c) = 0$  and  $f'' < 0$ , then  $f$  has a local \_\_\_\_\_ at  $x = c$ ;
- If  $f'(c) = 0$  and  $f'' > 0$ , then  $f$  has a local \_\_\_\_\_ at  $x = c$ ;
- If  $f'(c) = 0$  and  $f'' = 0$ , then the test fails. The function  $f$  may have a local maximum, a local minimum or neither.

The rest of today and tomorrow, we will focus on sketching curves using these two tests.

**Problem.** Sketch the general shape of a curve satisfying the given information

interval:	$x < 0$	$0 < x < 2$	$2 < x < 3$	$3 < x$
sign of $f'$ :	-	-	-	+
sign of $f''$ :	+	-	+	+

**Problem.** Find all local extrema of  $f(x) = -2 \cos x - \cos^2 x$  on the interval  $-\pi \leq x \leq \pi$ .