

 CALCULUS & ANALYTIC GEOMETRY II

The Fundamental Theorem(s) of Calculus

Accumulation function. Suppose that $f(x)$ is a continuous function on $[a, b]$, let

$$g(x) = \int_a^x f(t)dt.$$

g is *accumulating* the signed area under f between a and x .



Example 1.

Use the graph above to compute the following:

$$\int_1^3 f(t)dt$$

$$\int_1^5 f(t)dt$$

$$\int_1^1 f(t)dt$$

$$\int_1^0 f(t)dt$$

The Fundamental Theorem of Calculus Part 1 (FTC1) explains the rate of change of an accumulation function.

Fundamental Theorem of Calculus I (FTC1). If f is a continuous function on $[a, b]$, then the accumulation function g defined by

$$g(x) = \int_a^x f(t)dt \quad a \leq x \leq b$$

is continuous on $[a, b]$, differentiable on (a, b) , and $g'(x) = f(x)$.

Illustration. Let $g(x) = \int_1^x f(t)dt$ with f as in Example 1. Compute

$$g'(3) = \left. \frac{dg}{dx} \right|_{x=5} = \left. \frac{d}{dx} \int_1^x f(t)dt \right|_{x=0} =$$

Let's prove the FTC1 for a specific function.

$$f(t) =$$

Draw a rough sketch of $f(t)$.

$$\text{Let } g(x) = \int_a^x f(t) dt =$$

Find $g'(x)$ via the definition of derivative.

$$g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} =$$

Find the maximum and minimum values of f on $[x, x+h]$ and use the **Squeeze Theorem** to estimate

$$\leq \frac{1}{h} \int_x^{x+h} \text{_____} dt \leq$$

Conclusion. So $g'(x) =$

Question. How does our argument change if we move from a specific function like $f(t) = e^{-2t}$ to an arbitrary function $f(t)$? (See page 382 of your book.)

Practice Problems. Compute the following derivatives:

$$\frac{d}{dx} \int_5^x \frac{dt}{\ln t}$$

$$\frac{d}{dz} \int_{-4}^z 27 dt$$

$$\frac{d}{dx} \int_1^{x^2} \tan(t) \csc(t) dt$$

Fundamental Theorem of Calculus II (FTC2). If f is a continuous function on $[a, b]$ and F is any antiderivative of f , then

$$\int_a^b f(x)dx = F(b) - F(a).$$

Proof. This proof uses FTC1 and exploits the relationship between antiderivatives...

Let $g(x) = \int_a^x f(t)dt$. Then FTC1 implies that

$$g'(x) =$$

If F is any antiderivative of f , what is the relationship between F and g ?

Now compute $F(b) - F(a)$.

Section 5.4 restates FTC2 as follows:

The Net Change Theorem. The (definite) integral of a rate of change on an interval $[a, b]$ is the net change on the interval.

Restating the Net Change Theorem more mathematically, let $F(x)$ be a continuous function on $[a, b]$...

Quick check. If $w(t)$ is the weight of a child in lbs, what does $\int_2^5 w'(t)dt$ represent?

More Practice Problems. Compute the following definite integrals:

$$\int_5^8 \frac{dt}{t}$$

$$\int_{-4}^4 27dt$$

$$\int_{\pi/4}^{\pi/3} \tan(t) \sec(t) dt$$

IMPORTANT! Know the distinction between

$$\int_a^b f(t)dt$$

$$\int_a^x f(t)dt$$

$$\int f(t)dt$$

Clearly, being able to calculate antiderivatives is going to be important. We already know:

$$\int k dx =$$
$$\int x^n = \begin{cases} & n \neq 1 \\ & n = 1 \end{cases}$$

$$\int e^x dx = \qquad \int a^x dx =$$

$$\int \sin x dx = \qquad \int \cos x dx =$$

$$\int \sec^2 x dx = \qquad \int \csc^2 x dx =$$

$$\int \sec x \tan x dx = \qquad \int \csc x \cot x dx =$$

$$\int \frac{1}{x^2 + 1} dx = \qquad \int \frac{dx}{\sqrt{1 - x^2}} =$$

If you are facing an integral and you don't immediately know the antiderivative, rewrite, simplify, guess, and check.

1. $\int x^{-3}(x + 1) dx$

2. $\int (1 + \tan^2 \theta) d\theta$

3. $\int \cos \theta (\tan \theta + \sec \theta) d\theta$

4. $\int \frac{\csc x}{\csc x - \sin x} dx$