

---

 CALCULUS & ANALYTIC GEOMETRY II
 

---

## What to do about troublesome integrals

**Warm-up.** Find an antiderivative of  $\frac{e^x}{x}$ .

One possible answer is  $F(x) = \int_0^x \frac{e^t}{t} dt$ .

The majority of elementary functions don't have elementary antiderivatives. *Stewart, p. 488*

$$e^{x^2}$$



$$\sin(x^2)$$

$$\cos(e^x)$$

$$\frac{1}{\ln x}$$

But we still might want to calculate definite integrals involving these functions. What should we do?

**Big idea—Numerical Integration** Partition the interval of integration, replace troublesome function by a *closely fitting* polynomial on each subinterval, integrate polynomial, add results.



---


 CALCULUS & ANALYTIC GEOMETRY II
 

---

## Trapezoidal Approximations

Find the area of a trapezoid with base  $\Delta x$  and heights  $y_{i-1}$  and  $y_i$ .

Let  $T$  be the approximation of  $\int_a^b f(x)dx$  by  $n$  trapezoids where  $a = x_0, x_1, x_2, \dots, x_n = b$  are the subdivision points and  $y_i = f(x_i)$  for  $i = 1, 2, \dots, n$ . Then



**The Trapezoid Rule** To approximate  $\int_a^b f(x)dx$ , use

$$T = \frac{\Delta x}{2} (y_0 + 2y_1 + 2y_2 + \dots + 2y_{n-1} + y_n).$$

The  $y_i$ s are the values of  $f$  at the partition points  $x_0, x_1, \dots, x_n$  where  $\Delta x = \frac{(b-a)}{n}$ .

**Example 1.** Use the Trapezoid Rule with  $n = 4$  to estimate  $\int_1^2 x^2 dx$ . Compare this estimate with the exact solution.

*Ans: 75/32, 7/3  
% error: 0.00446*

---

 CALCULUS & ANALYTIC GEOMETRY II
 


---

 Simpson's Rule: Approximations Using Parabolas
 

---

**Warm-up.** Any three points define a parabola. Find the parabola that passes through the points  $(-1, 4)$ ,  $(0, 1)$ , and  $(1, 2)$ .

Find the area under the parabola  $y = Ax^2 + Bx + C$  passing through the points  $(-h, y_0)$ ,  $(0, y_1)$ , and  $(h, y_2)$ . Express your answer in terms of  $y_0$ ,  $y_1$ , and  $y_2$ .



How would the area change if the parabola were shifted horizontally?

So what is the area under the parabola  $y = Ax^2 + Bx + C$  passing through the points  $(x_2, y_2)$ ,  $(x_3, y_3)$ ,  $(x_4, y_4)$ ?

Computing the area under all the parabolas and adding the results gives the approximation

$$\int_a^b f(x) dx \approx \frac{h}{3}(y_0 + 4y_1 + y_2) +$$

**Simpson's Rule.** To approximate  $\int_a^b f(x)dx$ , use

$$S = \frac{\Delta x}{3}(y_0 + 4y_1 + 2y_2 + 4y_3 + \cdots + 2y_{n-2} + 4y_{n-1} + y_n).$$

The  $y_i$ s are the values of  $f$  at the partition points  $x_0, x_1, \dots, x_n$  where  $\Delta x = \frac{(b-a)}{n}$ .

**Example 2.** Use Simpson's Rule with  $n = 4$  to approximate  $\int_0^2 5x^4 dx$  and compare your result to the exact solution.

*Ans:  $32\frac{1}{12}$ , less than 3/10 %*

**Error Estimates in the Trapezoidal and Simpson's Rules.** If  $f''$  is continuous and  $M$  is any upper bound for the values of  $|f''|$  on  $[a, b]$ , the the error  $E_T$  in the trapezoidal approximation of the integral of  $f$  from  $a$  to  $b$  for  $n$  steps satisfies the inequality

$$|E_T| \leq \frac{M(b-a)^3}{12n^2}. \quad \text{Trapezoid}$$

If  $f^{(4)}$  is continuous and  $M$  is any upper bound for the values of  $|f^{(4)}|$  on  $[a, b]$ , then the error  $E_S$  in the Simpson's Rule approximation of the integral of  $F$  from  $a$  to  $b$  for  $n$  steps satisfies the inequality

$$|E_S| \leq \frac{M(b-a)^5}{180n^4}. \quad \text{Simpson's}$$

**Example 2 continued.** Approximate the error in example 2 using the above estimate.

**Example 3.** How large should  $n$  be to guarantee that Simpson's Rule approximation to  $\int_0^1 e^{x^2} dx$  is accurate to within .00001?

$$f^{(4)} = (12 + 48x^2 + 16x^4)e^{x^2}$$

*Ans:  $n \geq 20$ .*