


1. (a) $\Delta x = (b - a)/n = (4 - 0)/2 = 2$

$$L_2 = \sum_{i=1}^2 f(x_{i-1}) \Delta x = f(x_0) \cdot 2 + f(x_1) \cdot 2 = 2[f(0) + f(2)] = 2(0.5 + 2.5) = 6$$

$$R_2 = \sum_{i=1}^2 f(x_i) \Delta x = f(x_1) \cdot 2 + f(x_2) \cdot 2 = 2[f(2) + f(4)] = 2(2.5 + 3.5) = 12$$

$$M_2 = \sum_{i=1}^2 f(\bar{x}_i) \Delta x = f(\bar{x}_1) \cdot 2 + f(\bar{x}_2) \cdot 2 = 2[f(1) + f(3)] \approx 2(1.6 + 3.2) = 9.6$$

(b)



L_2 is an underestimate, since the area under the small rectangles is less than the area under the curve, and R_2 is an overestimate, since the area under the large rectangles is greater than the area under the curve. It appears that M_2 is an overestimate, though it is fairly close to I . See the solution to Exercise 45 for a proof of the fact that if f is concave down on $[a, b]$, then the Midpoint Rule is an overestimate of $\int_a^b f(x) dx$.

(c) $T_2 = (\frac{1}{2} \Delta x)[f(x_0) + 2f(x_1) + f(x_2)] = \frac{2}{2}[f(0) + 2f(2) + f(4)] = 0.5 + 2(2.5) + 3.5 = 9.$

This approximation is an underestimate, since the graph is concave down. Thus, $T_2 = 9 < I$. See the solution to Exercise 45 for a general proof of this conclusion.

(d) For any n , we will have $L_n < T_n < I < M_n < R_n$.


3. $f(x) = \cos(x^2)$, $\Delta x = \frac{1-0}{4} = \frac{1}{4}$

(a) $T_4 = \frac{1}{4 \cdot 2} [f(0) + 2f(\frac{1}{4}) + 2f(\frac{2}{4}) + 2f(\frac{3}{4}) + f(1)] \approx 0.895759$

(b) $M_4 = \frac{1}{4} [f(\frac{1}{8}) + f(\frac{3}{8}) + f(\frac{5}{8}) + f(\frac{7}{8})] \approx 0.908907$

The graph shows that f is concave down on $[0, 1]$. So T_4 is an underestimate and M_4 is an overestimate. We can conclude that

$$0.895759 < \int_0^1 \cos(x^2) dx < 0.908907.$$



9. $f(x) = \frac{\ln x}{1+x}$, $\Delta x = \frac{2-1}{10} = \frac{1}{10}$

(a) $T_{10} = \frac{1}{10 \cdot 2} [f(1) + 2f(1.1) + 2f(1.2) + \dots + 2f(1.8) + 2f(1.9) + f(2)] \approx 0.146879$

(b) $M_{10} = \frac{1}{10} [f(1.05) + f(1.15) + \dots + f(1.85) + f(1.95)] \approx 0.147391$

(c) $S_{10} = \frac{1}{10 \cdot 3} [f(1) + 4f(1.1) + 2f(1.2) + 4f(1.3) + 2f(1.4) + 4f(1.5) + 2f(1.6) + 4f(1.7) + 2f(1.8) + 4f(1.9) + f(2)]$

$$\approx 0.147219$$

17. $f(y) = \frac{1}{1+y^5}$, $\Delta y = \frac{3-0}{6} = \frac{1}{2}$

(a) $T_6 = \frac{1}{2 \cdot 2} [f(0) + 2f(\frac{1}{2}) + 2f(\frac{2}{2}) + 2f(\frac{3}{2}) + 2f(\frac{4}{2}) + 2f(\frac{5}{2}) + f(3)] \approx 1.064275$

(b) $M_6 = \frac{1}{2} [f(\frac{1}{4}) + f(\frac{3}{4}) + f(\frac{5}{4}) + f(\frac{7}{4}) + f(\frac{9}{4}) + f(\frac{11}{4})] \approx 1.067416$

(c) $S_6 = \frac{1}{2 \cdot 3} [f(0) + 4f(\frac{1}{2}) + 2f(\frac{2}{2}) + 4f(\frac{3}{2}) + 2f(\frac{4}{2}) + 4f(\frac{5}{2}) + f(3)] \approx 1.074915$

19. $f(x) = \cos(x^2)$, $\Delta x = \frac{1-0}{8} = \frac{1}{8}$

(a) $T_8 = \frac{1}{8 \cdot 2} \{f(0) + 2[f(\frac{1}{8}) + f(\frac{2}{8}) + \dots + f(\frac{7}{8})] + f(1)\} \approx 0.902333$

$M_8 = \frac{1}{8} [f(\frac{1}{16}) + f(\frac{3}{16}) + f(\frac{5}{16}) + \dots + f(\frac{15}{16})] = 0.905620$

(b) $f(x) = \cos(x^2)$, $f'(x) = -2x \sin(x^2)$, $f''(x) = -2 \sin(x^2) - 4x^2 \cos(x^2)$. For $0 \leq x \leq 1$, sin and cos are positive, so $|f''(x)| = 2 \sin(x^2) + 4x^2 \cos(x^2) \leq 2 \cdot 1 + 4 \cdot 1 \cdot 1 = 6$ since $\sin(x^2) \leq 1$ and $\cos(x^2) \leq 1$ for all x ,

and $x^2 \leq 1$ for $0 \leq x \leq 1$. So for $n = 8$, we take $K = 6$, $a = 0$, and $b = 1$ in Theorem 3, to get

$|E_T| \leq 6 \cdot 1^3 / (12 \cdot 8^2) = \frac{1}{128} = 0.0078125$ and $|E_M| \leq \frac{1}{256} = 0.00390625$. [A better estimate is obtained by noting from a graph of f'' that $|f''(x)| \leq 4$ for $0 \leq x \leq 1$.]

(c) Take $K = 6$ [as in part (b)] in Theorem 3. $|E_T| \leq \frac{K(b-a)^3}{12n^2} \leq 0.0001 \Leftrightarrow \frac{6(1-0)^3}{12n^2} \leq 10^{-4} \Leftrightarrow$

$\frac{1}{2n^2} \leq \frac{1}{10^4} \Leftrightarrow 2n^2 \geq 10^4 \Leftrightarrow n^2 \geq 5000 \Leftrightarrow n \geq 71$. Take $n = 71$ for T_n . For E_M , again take $K = 6$ in

Theorem 3 to get $|E_M| \leq 10^{-4} \Leftrightarrow 4n^2 \geq 10^4 \Leftrightarrow n^2 \geq 2500 \Leftrightarrow n \geq 50$. Take $n = 50$ for M_n .

20. $f(x) = e^{1/x}$, $\Delta x = \frac{2-1}{10} = \frac{1}{10}$

(a) $T_{10} = \frac{1}{10 \cdot 2} [f(1) + 2f(1.1) + 2f(1.2) + \dots + 2f(1.9) + f(2)] \approx 2.021976$

$M_{10} = \frac{1}{10} [f(1.05) + f(1.15) + f(1.25) + \dots + f(1.95)] \approx 2.019102$

(b) $f(x) = e^{1/x}$, $f'(x) = -\frac{1}{x^2} e^{1/x}$, $f''(x) = \frac{2x+1}{x^4} e^{1/x}$. Now f'' is decreasing on $[1, 2]$, so let $x = 1$ to take $K = 3e$.

$|E_T| \leq \frac{3e(2-1)^3}{12(10)^2} = \frac{e}{400} \approx 0.006796$. $|E_M| \leq \frac{|E_T|}{2} = \frac{e}{800} \approx 0.003398$.

(c) Take $K = 3e$ [as in part (b)] in Theorem 3. $|E_T| \leq \frac{K(b-a)^3}{12n^2} \leq 0.0001 \Leftrightarrow \frac{3e(2-1)^3}{12n^2} \leq 10^{-4} \Leftrightarrow$

$\frac{e}{4n^2} \leq \frac{1}{10^4} \Leftrightarrow n^2 \geq \frac{10^4 e}{4} \Leftrightarrow n \geq 83$. Take $n = 83$ for T_n . For E_M , again take $K = 3e$ in Theorem 3 to get

$|E_M| \leq 10^{-4} \Leftrightarrow n^2 \geq \frac{10^4 e}{8} \Leftrightarrow n \geq 59$. Take $n = 59$ for M_n .

21. $f(x) = \sin x, \Delta x = \frac{\pi - 0}{10} = \frac{\pi}{10}$

(a) $T_{10} = \frac{\pi}{10 \cdot 2} [f(0) + 2f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + \cdots + 2f\left(\frac{9\pi}{10}\right) + f(\pi)] \approx 1.983524$

$$M_{10} = \frac{\pi}{10} [f\left(\frac{\pi}{20}\right) + f\left(\frac{3\pi}{20}\right) + f\left(\frac{5\pi}{20}\right) + \cdots + f\left(\frac{19\pi}{20}\right)] \approx 2.008248$$

$$S_{10} = \frac{\pi}{10 \cdot 3} [f(0) + 4f\left(\frac{\pi}{10}\right) + 2f\left(\frac{2\pi}{10}\right) + 4f\left(\frac{3\pi}{10}\right) + \cdots + 4f\left(\frac{9\pi}{10}\right) + f(\pi)] \approx 2.000110$$

Since $I = \int_0^\pi \sin x \, dx = [-\cos x]_0^\pi = 1 - (-1) = 2$, $E_T = I - T_{10} \approx 0.016476$, $E_M = I - M_{10} \approx -0.008248$,

and $E_S = I - S_{10} \approx -0.000110$.

(b) $f(x) = \sin x \Rightarrow |f^{(n)}(x)| \leq 1$, so take $K = 1$ for all error estimates.

$$|E_T| \leq \frac{K(b-a)^3}{12n^2} = \frac{1(\pi-0)^3}{12(10)^2} = \frac{\pi^3}{1200} \approx 0.025839. \quad |E_M| \leq \frac{|E_T|}{2} = \frac{\pi^3}{2400} \approx 0.012919.$$

$$|E_S| \leq \frac{K(b-a)^5}{180n^4} = \frac{1(\pi-0)^5}{180(10)^4} = \frac{\pi^5}{1,800,000} \approx 0.000170.$$

The actual error is about 64% of the error estimate in all three cases.

(c) $|E_T| \leq 0.00001 \Leftrightarrow \frac{\pi^3}{12n^2} \leq \frac{1}{10^5} \Leftrightarrow n^2 \geq \frac{10^5 \pi^3}{12} \Rightarrow n \geq 508.3$. Take $n = 509$ for T_n .

$$|E_M| \leq 0.00001 \Leftrightarrow \frac{\pi^3}{24n^2} \leq \frac{1}{10^5} \Leftrightarrow n^2 \geq \frac{10^5 \pi^3}{24} \Rightarrow n \geq 359.4$$
. Take $n = 360$ for M_n .

$$|E_S| \leq 0.00001 \Leftrightarrow \frac{\pi^5}{180n^4} \leq \frac{1}{10^5} \Leftrightarrow n^4 \geq \frac{10^5 \pi^5}{180} \Rightarrow n \geq 20.3.$$

Take $n = 22$ for S_n (since n must be even).

29. $\Delta x = (b-a)/n = (6-0)/6 = 1$

(a) $T_6 = \frac{\Delta x}{2} [f(0) + 2f(1) + 2f(2) + 2f(3) + 2f(4) + 2f(5) + f(6)]$
 $\approx \frac{1}{2}[3 + 2(5) + 2(4) + 2(2) + 2(2.8) + 2(4) + 1]$
 $= \frac{1}{2}(39.6) = 19.8$

(b) $M_6 = \Delta x [f(0.5) + f(1.5) + f(2.5) + f(3.5) + f(4.5) + f(5.5)]$
 $\approx 1[4.5 + 4.7 + 2.6 + 2.2 + 3.4 + 3.2]$
 $= 20.6$

(c) $S_6 = \frac{\Delta x}{3} [f(0) + 4f(1) + 2f(2) + 4f(3) + 2f(4) + 4f(5) + f(6)]$
 $\approx \frac{1}{3}[3 + 4(5) + 2(4) + 4(2) + 2(2.8) + 4(4) + 1]$
 $= \frac{1}{3}(61.6) = 20.5\bar{3}$

37. Let $y = f(x)$ denote the curve. Using cylindrical shells, $V = \int_2^{10} 2\pi x f(x) dx = 2\pi \int_2^{10} x f(x) dx = 2\pi I_1$.

Now use Simpson's Rule to approximate I_1 :

$$\begin{aligned}I_1 &\approx S_8 = \frac{10-2}{3(8)} [2f(2) + 4 \cdot 3f(3) + 2 \cdot 4f(4) + 4 \cdot 5f(5) + 2 \cdot 6f(6) + 4 \cdot 7f(7) + 2 \cdot 8f(8) + 4 \cdot 9f(9) + 10f(10)] \\&\approx \frac{1}{3}[2(0) + 12(1.5) + 8(1.9) + 20(2.2) + 12(3.0) + 28(3.8) + 16(4.0) + 36(3.1) + 10(0)] \\&= \frac{1}{3}(395.2)\end{aligned}$$

Thus, $V \approx 2\pi \cdot \frac{1}{3}(395.2) \approx 827.7$ or 828 cubic units.