

Disparities in Air Pollution Exposure in the United States by Race/Ethnicity and Income, 1990–2010

Jiawen Liu,¹ Lara P. Clark,¹ Matthew J. Bechle,¹ Anjum Hajat,² Sun-Young Kim,³ Allen L. Robinson,⁴ Lianne Sheppard,^{5,6} Adam A. Szpiro,⁵ and Julian D. Marshall¹

¹Department of Civil & Environmental Engineering, University of Washington, Seattle, Washington, USA

²Department of Epidemiology, University of Washington, Seattle, Washington, USA

³Department of Cancer Control and Population Health, Graduate School of Cancer Science and Policy, National Cancer Center, Goyang-si, Gyeonggi-do, Korea

⁴Department of Mechanical Engineering & Department of Engineering and Public Policy, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

⁵Department of Biostatistics, University of Washington, Seattle, Washington, USA

⁶Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, USA

BACKGROUND: Few studies have investigated air pollution exposure disparities by race/ethnicity and income across criteria air pollutants, locations, or time.

OBJECTIVE: The objective of this study was to quantify exposure disparities by race/ethnicity and income throughout the contiguous United States for six criteria air pollutants, during the period 1990 to 2010.

METHODS: We quantified exposure disparities among racial/ethnic groups (non-Hispanic White, non-Hispanic Black, Hispanic (any race), non-Hispanic Asian) and by income for multiple spatial units (contiguous United States, states, urban vs. rural areas) and years (1990, 2000, 2010) for carbon monoxide (CO), nitrogen dioxide (NO₂), ozone (O₃), particulate matter with aerodynamic diameter ≤2.5 μm (PM_{2.5}; excluding year-1990), particulate matter with aerodynamic diameter ≤10 μm (PM₁₀), and sulfur dioxide (SO₂). We used census data for demographic information and a national empirical model for ambient air pollution levels.

RESULTS: For all years and pollutants, the racial/ethnic group with the highest national average exposure was a racial/ethnic minority group. In 2010, the disparity between the racial/ethnic group with the highest vs. lowest national-average exposure was largest for NO₂ [54% (4.6 ppb)], smallest for O₃ [3.6% (1.6 ppb)], and intermediate for the remaining pollutants (13%–19%). The disparities varied by U.S. state; for example, for PM_{2.5} in 2010, exposures were at least 5% higher than average in 63% of states for non-Hispanic Black populations; in 33% and 26% of states for Hispanic and for non-Hispanic Asian populations, respectively; and in no states for non-Hispanic White populations. Absolute exposure disparities were larger among racial/ethnic groups than among income categories (range among pollutants: between 1.1 and 21 times larger). Over the period studied, national absolute racial/ethnic exposure disparities declined by between 35% (0.66 μg/m³; PM_{2.5}) and 88% (0.35 ppm; CO); relative disparities declined to between 0.99 × (PM_{2.5}; i.e., nearly zero change) and 0.71 × (CO; i.e., a ~29% reduction).

DISCUSSION: As air pollution concentrations declined during the period 1990 to 2010, absolute (and to a lesser extent, relative) racial/ethnic exposure disparities also declined. However, in 2010, racial/ethnic exposure disparities remained across income levels, in urban and rural areas, and in all states, for multiple pollutants. <https://doi.org/10.1289/EHP8584>

Introduction

Air pollution is associated with ~100,000 annual premature deaths in the United States in 2017 (Stanaway et al. 2018) and has been linked to cardiovascular disease, respiratory disease, cancers, adverse birth outcomes, cognitive decline, and other health impacts (Cohen et al. 2017; Darrow et al. 2011; Lelieveld et al. 2015; Paul et al. 2019; Pope et al. 2009; Rivas et al. 2019; Stieb et al. 2012; Underwood 2017). Air pollution and its associated health impacts are not equitably distributed by race/ethnicity or income. Previous research has documented higher-than-average air pollution exposures for racial/ethnic minority populations and lower-income populations in the United States (Brulle and Pellow 2006; Evans and Kantrowitz 2002; Mohai et al. 2009), leading to disparities in attributable health impacts (Bowe et al. 2019; Fann et al. 2019; Gee and Payne-Sturges 2004). Most investigations of disparities in air pollution exposure involve a

single pollutant, location, and/or time point [see, e.g., literature reviews by Hajat et al. (2015) and Marshall et al. (2014); see Table S2]. Evidence from broader investigations suggests that exposure disparities by race/ethnicity and/or income can vary by pollutant (Rosofsky et al. 2018), location [e.g., by state (Bullock et al. 2018; Salazar et al. 2019), urbanicity (Mikati et al. 2018), metropolitan area (Zwickl et al. 2014; Downey et al. 2008)], and time point (Ard 2015; Clark et al. 2017; Kravitz-Wirtz et al. 2016; Colmer et al. 2020). However, to our knowledge, broad patterns in exposure disparities have not yet been investigated, using consistent methods, across pollutants, locations, and time points, for the contiguous U.S. population.

The objective of our research was to comprehensively and consistently investigate disparities in exposure to U.S. Environmental Protection Agency (U.S. EPA) criteria air pollutants for the two decades following the 1990 Clean Air Act Amendments in the United States. Specifically, we investigated the following questions regarding disparities in exposure to six criteria air pollutants: *a*) How do exposures vary by race/ethnicity and income? *b*) How do racial/ethnic exposure disparities vary by pollutant? *c*) How do racial/ethnic exposure disparities vary by location (state, urban vs. rural areas)? *d*) How have racial/ethnic exposure disparities changed over time? To address these questions, we combined demographic data from the U.S. Census (Manson et al. 2019) with predictions of outdoor average levels of six criteria air pollutants from a publicly available national empirical model derived from satellite, measurement, and other types of data (Kim et al. 2020) at the spatial scale of census block groups and census tracts. We then analyzed disparities in exposure to six criteria air pollutants [all criteria air pollutants except lead (Pb); i.e., carbon monoxide (CO), nitrogen dioxide (NO₂), ozone (O₃), fine and respirable suspended particulate matter with an aerodynamic diameter ≤2.5 μm

Address correspondence to Julian Marshall, Department of Civil & Environmental Engineering, University of Washington, 201 More Hall, Seattle, WA 98195 USA. Telephone: (206) 685-2591. Email: jdmars@uw.edu

Supplemental Material is available online (<https://doi.org/10.1289/EHP8584>).
The authors declare they have no actual or potential competing financial interest.

Received 29 October 2020; Revised 20 October 2021; Accepted 9 November 2021; Published 15 December 2021.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

(PM_{2.5}), particulate matter with aerodynamic diameter $\leq 10 \mu\text{m}$ (PM₁₀), and sulfur dioxide (SO₂) by race/ethnicity (four racial/ethnic groups: non-Hispanic White, non-Hispanic Black, Hispanic (any race), non-Hispanic Asian) and income (16 household income categories) across time points (decennial census years: 1990, 2000, and 2010) and spatial units (contiguous United States, state, urban vs. rural areas).

Methods

Demographic and Air Pollution Datasets

We obtained demographic data (i.e., population estimates by race/ethnicity, household income, and household income disaggregated by race/ethnicity) and map boundaries (e.g., states, census tracts, and census block groups) for the contiguous United States from the 1990, 2000, and 2010 decennial censuses from the IPUMS National Historic Geographic Information System (NHGIS) (Manson et al. 2019).

NHGIS provides, for each census block group, and for 1990, 2000, and 2010 (standardized to 2010 spatial boundaries), population estimates for six census self-reported racial groups: *a*) White alone, *b*) Black or African American alone, *c*) American Indian and Alaska Native alone, *d*) Asian and Pacific Islander alone, *e*) some other race alone, and *f*) two or more races. NHGIS reports population estimates for two census self-reported ethnic groups: *a*) Hispanic or Latino and *b*) not Hispanic or Latino. Thus, there are a total of 12 combined racial/ethnic groups in NHGIS (six racial groups, two ethnic groups). Our main analyses of racial/ethnic exposure disparities included the four largest racial/ethnic groups, which in total covered 307 million people (97.2% of the population) in the contiguous United States in 2010: *a*) not Hispanic or Latino, White alone (64% of the population; hereafter, “non-Hispanic White”), *b*) Hispanic or Latino of any race(s) (16%; hereafter, “Hispanic”), *c*) not Hispanic or Latino, Black or African American alone (12%; hereafter, “non-Hispanic Black”), and *d*) not Hispanic or Latino, Asian and Pacific Islander alone (4.6%; hereafter, “non-Hispanic Asian”).

For analyses by income in 2010, we used 2010 NHGIS household income estimates. For each block group, NHGIS reports the number of households in 16 annual household income categories (total covered in 2010: 114 million households) (in 2010 inflation-adjusted U.S. dollars): <10,000, 10,000–15,000, 15,000–20,000, 20,000–25,000, 25,000–30,000, 30,000–35,000, 35,000–40,000, 40,000–45,000, 45,000–50,000, 50,000–60,000, 60,000–75,000, 75,000–100,000, 100,000–125,000, 125,000–150,000, 150,000–200,000, and >200,000.

For analyses by income disaggregated by race/ethnicity in 2010, data from the 2010 NHGIS were available at the census tract level. For each census tract, NHGIS reports householder data for eight predefined race and/or ethnicity categories within each of the 16 census income categories, including one category based on both race and ethnicity (non-Hispanic White), one based on ethnicity regardless of race (Hispanic or Latino), and six based on race regardless of ethnicity (Black or African American alone, American Indian and Alaska Native alone, Asian alone, Native Hawaiian or Other Pacific Islander alone, some other race alone, and two or more race). To best match demographic variables used in race/ethnicity analysis at the census block group level, we reported results for four largest racial/ethnic groups (total covered in 2010: 113 million census householders, 98.5% of householders with data on income by race/ethnicity): not Hispanic or Latino, White alone (71% of householders; hereafter, “non-Hispanic White”), Hispanic or Latino (12%; hereafter, “Hispanic”), Black or African American alone (12%; hereafter, “Black”), and Asian

alone (3.8%; hereafter, “Asian”). Thus, for the data used for the household income by race/ethnicity analysis (but not for other analyses), Black and Asian categories included both Hispanic and non-Hispanic individuals; for these analyses (but not others), Hispanic Black populations ($\sim 0.40\%$ of the population) would be included in results for Hispanic and for Black populations, and Hispanic Asian populations ($\sim 0.08\%$) would be included in results for Hispanic and for Asian populations. Additionally, for the data used for the household income by race/ethnicity analysis (but not for other analyses), the Asian category does not also include Pacific Islander populations.

The U.S. Census Bureau defined census blocks as “urban” or “rural” based on population density and other characteristics (Ratcliffe et al. 2016). We used 2010 census urban/rural block definitions to define a 2010 census block group for all 3 y (1990, 2000, and 2010) as rural if all blocks inside it were rural, and we defined the remaining block groups as urban (i.e., each census block group and urban/rural designation was the same in 1990, 2000, and 2010).

Average estimates of ambient air pollution levels for U.S. EPA criteria pollutants were obtained from the Center for Air, Climate, and Energy Solutions (CACES) empirical models for the contiguous United States (www.caces.us/data). These models incorporate satellite-derived estimates of air pollution, satellite-derived land cover data, land use data, U.S. EPA monitoring station data, and universal Kriging (Kim et al. 2020); estimated pollution levels were available by census block at block centroids based on 2010 census boundaries for the years from 1990 to 2010 for all pollutants except PM_{2.5} (for which monitoring data and exposure models were only available starting in 1999). Estimated levels of O₃ from the CACES empirical model are 5-month summer averages (specifically, the average during May–September of the daily maximum 8-h moving average); for the remaining pollutants pollutants, estimated levels are annual averages.

CACES model performance during the years studied here (2000, 2010 for PM_{2.5}; 1990, 2000, 2010 for the other pollutants), as measured by cross-validated R^2 , was 0.84–0.89 for NO₂, 0.85 for PM_{2.5}, 0.62–0.82 for O₃, 0.56–0.62 for PM₁₀, 0.32–0.66 for SO₂, and 0.34–0.57 for CO (Kim et al. 2020). Mean error (ME) across the census years studied was between -0.02 and 0 ppm for CO, -0.04 to 0 ppb for O₃, -0.09 to -0.06 ppb for NO₂, -0.17 to -0.13 ppb for SO₂, -0.31 to $-0.26 \mu\text{g m}^{-3}$ for PM₁₀, and -0.05 to $-0.02 \mu\text{g m}^{-3}$ for PM_{2.5}. Mean bias (MB) was 13%–22% for SO₂, and <10% for the other pollutants (Table S1); further details about the models and model performance are in Kim et al. (2020) and Liu (2021).

Combining Demographic and Air Pollution Data

We matched the CACES empirical model results and the U.S. census demographic data using the 2010 census spatial boundary definitions (from finest to coarsest spatial resolution: block, block group, and tract boundaries) for the three census years (1990, 2000, 2010). We matched census block-level CACES model predictions for criteria air pollutants (blocks in 2010 in the contiguous United States: $n = \sim 7$ million; average: ~ 44 residents per block) to census block group-level demographic data (block groups: $n = \sim 220,000$; average: ~ 1400 residents per block group) by calculating population-weighted mean of the block-level predictions, for all blocks in that block-group. Similarly, to match census tract-level demographic data (tracts: $n = \sim 74,000$; ~ 4200 residents per tract), we calculated the population-weighted mean air pollution levels for all census block groups located within that tract.

Estimating Exposures to Pollutants

We estimated annual pollutant-specific exposures for 1990 (excluding PM_{2.5}), 2000, and 2010 based on population-weighted mean predicted ambient air pollution levels for each demographic group [race/ethnicity, income, and income by race/ethnicity; results for additional groups (income poverty ratio, age, language, mobility, travel time) are described in the Supplemental Material (SM)]. The data for the five additional groups (income poverty ratio, age, language, mobility, travel time) were extracted from NHGIS (i.e., we are directly employing values calculated by NHGIS; the values employed do not reflect our own data or calculations) (Manson et al. 2019). For all five additional groups, the rationale for including them is to explore whether exposures vary univariately for that demographic attribute. For all five additional groups, the categories used follow NHGIS categories and/or natural breaks in the data [e.g., for a ratio, separating values at, e.g., 0.5, 1.0, 1.5, 2.0; for age, separating young children as age 4 y or below, other children (who, typically, attend K12 education) as age 5–17 y, adults as age 18–64 y, and older adults as age 65+ y (reflecting an assumed retirement age)]. Income poverty ratio is defined by the U.S. Census as the ratio of income to poverty level in the past 12 months (Manson et al. 2019). The poverty level varies by number of people in the family and their ages; poverty level does not vary geographically (i.e., the same threshold is used throughout the United States) (U.S. Census Bureau 2021). In results shown in the SM for income to poverty ratio, we bin this ratio into five categories: <0.5, 0.5–1, 1–1.5, 1.5–2, and >2. The motivation for this analysis is to investigate income relative to the U.S. Census-defined poverty level. Age is binned into four categories: <5 y old, 5–17 y old, 18–64 y old, and 65+ y old. Language refers to language(s) spoken in the home. For households in which language(s) other than English are spoken, the U.S. Census subdivides household counts into *a*) households in which no one age 14 y and over speaks English only, and *b*) households in which one or more people age 14 y and over speaks English “very well.” We bin the NHGIS household language data into nine categories: English only, Spanish language and no English, English and a Spanish language, Asian language and no English, English and an Asian language, European language and no English, English and a European language, other language and no English, English and other language. Mobility refers to geographical mobility in the past year for current residence, based on metropolitan statistical areas (MSAs). We bin mobility into six categories: *a*) same house 1 y ago, *b*) different house: moved from same metropolitan, *c*) different house: moved from different metropolitan, *d*) different house: moved from micropolitan, *e*) different house: moved from not metropolitan nor micropolitan, and *f*) abroad 1 y ago. Travel time refers to travel time to work for workers age 16+ y who did not work at home. We divide the data into seven categories: <10 min, 10–20 min, 20–30 min, 30–40 min, 40–60 min, 60–90 min, and >90 min. This approach (average ambient air pollution level at residential census block group or tract) is broadly consistent with many examples in research and practice, including U.S. EPA monitors (Office of Air Quality Planning and Standards 2008), the National Ambient Air Quality Standards (e.g., Clean Air Scientific Advisory Committee 2010; Independent Particulate Matter Review Panel 2020; U.S. EPA 2019, 2020), many influential epidemiological studies (e.g., Di et al. 2017; Laden et al. 2006; Pope et al. 2009, 2020; Shi et al. 2016; Zanobetti and Schwartz 2009), and national empirical models for air pollution in the United States (e.g., Bechle et al. 2015; Di et al. 2020; Goldberg et al. 2019; Kim et al. 2020; Novotny et al. 2011; U.S. EPA 2016; Van Donkelaar et al. 2019; Young et al. 2016). We used the finest publicly available census spatial boundary data to estimate exposures for each analysis (income by race/ethnicity: tracts; all other analyses: block groups) based on availability of census demographic data.

The national annual (for O₃, 5-month average; for remaining pollutants, annual average) exposure (e_i) for demographic group i was calculated for a given pollutant and year as:

$$e_i = \frac{\sum_{j=1}^n c_j p_{ij}}{\sum_{j=1}^n p_{ij}}, \quad (1)$$

where c_j is the predicted average ambient pollution level for census block group or census tract j [here and after, we use c to represent ambient pollution level (observed or predicted) and e to represent population-weighted value for c], p_{ij} is the population of demographic group i in census block group or census tract j , and n is the number of census block groups or census tracts in the analyzed spatial level [the contiguous United States, each of the 49 “states” (including the District of Columbia plus the 48 contiguous states), and urban vs. rural areas].

National Exposure Disparities Analyses

Our primary exposure disparity metrics are based on absolute and relative differences in population-weighted mean air pollution exposures. We selected metrics based on mean pollution levels for consistency with our focus on broad national average patterns in exposure disparities among multiple pollutants. Absolute disparity metrics are often connect to pollutant-specific health impacts (Harper et al. 2013) (the present article focuses on pollution levels rather than health outcomes). Relative disparity metrics (e.g., ratios, relative percent differences) are relevant for quantifying disproportionality in exposure burdens, in a way that can be compared or summarized among different pollutants. An important limitation of these metrics (based on differences in mean exposures) is that they do not include information about disparities across the full exposure distributions (Harper et al. 2013). To address this limitation, we conducted supplemental analyses using inequality metrics accounting for full exposure distributions (Gini Coefficient and between-group Atkinson Index), as described in the SM, as well as sensitivity analyses comparing metrics based on other specific points of the exposure distribution (i.e., comparing specific exposure percentiles) as described below.

We calculated the absolute and relative exposure disparity metrics using two different approaches nationally: *a*) by race/ethnicity group and/or income category (i.e., the unit of analysis is a national subpopulation defined by race/ethnicity and/or income) and *b*) by local demographic characteristics (i.e., the unit of analysis is a set of census block groups defined based on proportion of racial/ethnic minority residents).

National exposure disparity metrics based on racial/ethnic group and/or income category. Our primary absolute disparity metric for quantifying national racial/ethnic exposure disparities is the pollutant-specific absolute difference in population-weighted average pollution level, as calculated using Equation 1 with block group-level data, between the racial/ethnic group with the highest national mean exposure (“most-exposed group”) and the racial/ethnic group with the lowest national mean exposure (“least-exposed group”) among the four racial/ethnic groups (non-Hispanic White, non-Hispanic Black, non-Hispanic Asian, Hispanic); here, the unit of analysis is a racial/ethnic group. In addition, we derived the percent difference relative to the model-predicted national mean exposure level for that pollutant $\{[(\text{population-weighted mean in most exposed} - \text{population-weighted mean in least exposed}) / \text{national mean exposure}] \times 100\%$. We also included relative exposure disparity metric as the pollutant-specific exposure ratio (i.e., population-weighted mean of most-exposed group / population-weighted mean of least-exposed group). Both the absolute and relative exposure disparity metrics are constructed

based on differences between most- and least-exposed racial/ethnic groups, to provide a measure of overall racial/ethnic disparities that avoids preselecting two specific groups for comparison and accounts for exposure disparities across multiple groups, in a consistent way for each pollutant (accounting for potential differences in the most- and least-exposed racial/ethnic groups by pollutant). We also report averages in relative disparities across pollutants as a representation of overall average inequalities in exposure to multiple pollutants, not as a representation of inequalities in health risks, which are pollutant-specific and depend on absolute levels of pollution exposure. Last, as a supplemental comparison among pollutants, we also calculated inequality metrics that account for the full exposure distributions: Gini coefficients by race/ethnicity and between-group Atkinson Indices.

To quantify national income-based exposure disparities, we calculated the pollutant-specific absolute difference in population-weighted average pollution level, using Equation 1 with block group-level data, between the lowest (<\$10,000) and the highest (>\$200,000) household income categories (of the 16 census categories). Additionally, as a relative disparity metric, we calculated the relative percent difference in mean exposures between the lowest and highest income categories. As a supplementary analysis, we calculated similar absolute and relative exposure disparity metrics between the income categories containing the 25th percentile (\$20,000–\$25,000) and the 75th percentile (\$75,000–\$100,000) of the income distribution.

To quantify national exposure disparities by race/ethnicity and income, we first calculated the absolute difference in population-weighted average pollution level between the most- and least-exposed racial/ethnic group (among the four racial/ethnic groups, not mutually exclusive with four racial/ethnic groups in racial/ethnic disparity, as described in “Demographic and Air Pollution Data Sets” in the “Methods” section) within each of the 16 census income categories, and then averaged that income category-specific racial/ethnic exposure disparity across all 16 income categories, for each pollutant. In the analyses for both race/ethnicity and income, we used census data for householders to calculate exposures for the four racial/ethnic groups using Equation 1 with tract-level data. Reflecting publicly available census data for racial/ethnic groups by income category, for this section only, the Black and Asian groups include Hispanic and non-Hispanic individuals, and the Asian group does not include Pacific Islander individuals. As a relative disparity metric, we divided the absolute exposure disparity metric by the national mean pollution level, for each of the pollutants.

National exposure disparity metrics based on local demographic characteristics (i.e., block group bins by proportion of racial/ethnic minority residents). We also investigated exposure disparities based on racial/ethnic minority resident percentages; here, the unit of analysis is bin of census block groups. Each block group bin was defined as single percentile (i.e., 1%) of all block groups stratified by the proportion of racial/ethnic minority residents. There were approximately 215,000 block groups in 2010, so each block group bin contained approximately 2,150 block groups. To investigate racial/ethnic disparities among block group bins, we rank ordered all census block group bins based on percent of racial/ethnic minority residents (i.e., people self-reporting any race/ethnicity other than non-Hispanic White alone). For example, the first block group bin was the first percentile and consisted of all block groups with between 0% and 0.67% racial/ethnic minority residents; the second block group bin was the second percentile, consisting of all block groups with 0.67%–0.97% racial/ethnic minority residents; the third block group bin consisted of all block groups with 0.97%–1.2% racial/ethnic minority residents, and so on through all 100 block group

bins. The last block group bin consisted of all block groups with 99.1%–99.6% racial/ethnic minority residents. The annual exposure (e_{ig}) for demographic group i for the g th percentile census block group bin (i.e., the average exposure across all block groups in the g th percentile for proportion of residents that belong to a racial/ethnic minority group) was calculated for a given pollutant and year as:

$$e_{ig} = \frac{\sum_{j=1}^{n_g} c_j p_{ij}}{\sum_{j=1}^{n_g} p_{ij}}, \quad (2)$$

where c_j is the predicted average ambient pollution level for census block group j , p_{ij} is the population of demographic group i in census block group j , and n_g is the number of census block groups in the g th percentile block group bin. The absolute disparity is calculated as the exposure difference between block groups with the highest vs. lowest deciles of proportion racial/ethnic minority residents, and, similarly, the relative disparity is calculated as the exposure ratio between block groups with the highest vs. lowest deciles of proportion racial/ethnic minority residents.

Sensitivity Analysis on Robustness of National Exposure Disparity Estimates

We conducted three sensitivity tests to investigate the robustness of conclusions based on estimated exposure disparities. First, as a sensitivity test for conclusions based on comparisons of mean values’ rank order for exposures between groups, we calculated disparities using different metrics of the exposure distribution (10th, 25th, 50th, 75th, 90th percentiles).

The remaining two sensitivity tests investigated whether conclusions here are robust to uncertainty in empirical model predictions. Specifically, in the second sensitivity test, we repeated the analysis of national mean exposures by racial/ethnic group, but for only the population living in a census block group with a U.S. EPA monitor in 2010. In this sensitivity test, for the pollution levels, we employ the monitor observations rather than the empirical model results. We then calculated Spearman rank order correlation of relative disparities by pollutant (between the most- and least-exposed group) between base case and sensitivity test.

In the third sensitivity test, we compared the magnitude of uncertainties in the estimated racial/ethnic exposure disparities with the magnitude of the estimated racial/ethnic exposure disparities. To assess the potential impact of model error on racial/ethnic disparities, we first calculated population-weighted mean error (ME_i) for each racial/ethnic group, i , using Equation 3:

$$ME_i = \frac{\sum_{j=1}^{n_o} (c_{jm} - c_{jo}) p_{ij}}{\sum_{j=1}^{n_o} p_{ij}}, \quad (3)$$

where c_{jm} is the predicted average ambient pollution level for census block group j , c_{jo} is the measured average ambient pollution level across all reporting U.S. EPA monitors within census block group j , p_{ij} is the population of demographic group i in block group j , and n_o is the total number of census block groups with EPA monitors. For each pollutant, the ME of disparity between two racial/ethnic groups i_1 and i_2 induced by the model was calculated as the difference between population-weighted ME for the most- and least-exposed racial/ethnic groups i_1 and i_2 . Calculated uncertainties are based on comparison with U.S. EPA measured pollution level in 2010. We then derived the ratio between the uncertainty due to exposure model error (i.e., the difference in population-weighted mean errors between racial/ethnic groups) and the estimated disparity in mean annual exposures between the most- and least-exposed racial/ethnic groups.

National Analysis of High-End Exposure Disparities in 2010

To quantify racial/ethnic disparities at the highest exposure levels, we analyzed the racial/ethnic composition of census block groups above the 90th percentiles of the pollution level among all census block groups. This analysis was done separately for each pollutant. First, for each of the four largest racial/ethnic groups, we estimated the proportion of that group's national population who lived in a high-exposure block group; here, our unit of analysis is a racial/ethnic group. This calculation reflects the proportion of a racial/ethnic group's total U.S. population who lived in heavily polluted (above the 90th percentile) block groups. We performed this calculation for each pollutant and each racial/ethnic group, using Equation 4.

$$a_i = \frac{\sum_{j=1}^{n_{90}} p_{ij}}{P_{total_national_i}} \times 100\%, \quad (4)$$

where a_i is the percent of racial/ethnic group i living in a block group with concentration above the 90th percentile for that pollutant, p_{ij} is the population of group i in census block group j , $P_{total_national_i}$ is the total population for demographic group i in the United States, and n_{90} is the number of census block groups with mean pollutant concentration >90th percentile.

In the second analysis, which was the converse of the first, we investigated the racial/ethnic composition of block groups above the 90th percentile for average pollution level. Here, our unit of analysis is all block groups above the 90th percentile. This calculation reflects the demographics of only people that lived in heavily polluted block groups. We completed this calculation for each pollutant and each racial/ethnic group using Equation 5.

$$b_i = \frac{\sum_{j=1}^{n_{90}} p_{ij}}{P_{total_block_group}} \times 100\%, \quad (5)$$

where b_i is (when considering only the people counted toward $P_{total_block_group}$) the percent of people who are in demographic group i , and $P_{total_block_group}$ is the total population of census block groups above the 90th percentile for that pollutant.

In addition, we explored differences in exposures to multiple pollutants by race/ethnicity by using data for 2010 and Equation 3 to estimate the proportion of each major race/ethnicity group's total U.S. population living in block groups with mean exposure levels above the 90th percentile for 0, 1, 2, 3, and ≥ 4 pollutants, respectively.

Counterfactual Analysis of Migration

We investigated whether changes in racial/ethnic exposure disparities over time were mainly attributable to changes in air pollution levels ("air pollution") or changes in where people lived (abbreviated as "migration", but also including immigration and other shifts in demographic patterns) as a sensitivity analysis. To do so, we employed two counterfactual scenarios (Clark et al. 2017) during two decades (1990 to 2000; 2000 to 2010). For each scenario and year, we calculated exposures for the four largest racial/ethnic groups for the contiguous U.S. population using Equation 1 based on census block group data. We then calculated the absolute racial/ethnic exposure disparity between the most- and least-exposed racial/ethnic groups (referred to in this section as "disparity") for all pollutants with available data (i.e., all except PM_{2.5} in 1990). To analyze the period 1990 to 2000, we calculated the change in disparity attributable to air pollution changing from 1990 to 2000 levels but with demographics remained constant at 1990 values (counterfactual scenario A—i.e., "counterfactual" because it includes consideration of year-

2000 pollution levels with year-1990 demographics) and, separately, used 1990 air pollution levels with demographic data changing from 1990 to 2000 values (counterfactual scenario B—includes consideration of year-1990 pollution levels with year-2000 demographics). To estimate the separate contribution of changes in air pollution during the period 1990 to 2000, we divided the disparity-changes from counterfactual scenario A by the "true" calculated disparity changes between 1990 and 2000 (i.e., using 1990 air pollution levels with 1990 demographic data, and using 2000 air pollution levels with 2000 demographic data). Similarly, to estimate the separate contribution of migration during 1990 to 2000, we divided the disparity changes from counterfactual scenario B by the "true" calculated disparity change between 1990 and 2000. Last, we used an analogous approach to analyze the next decade: 2000 to 2010.

Exposure Disparities Comparison Metrics for States

We investigated patterns among the 48 states of the contiguous United States plus the District of Columbia (DC) (hereafter, "states" refers to 48 states and DC, a total of 49 geographic units in state-level related calculations) using two metrics for absolute exposure disparity by race/ethnicity. First, for each state, pollutant, and race/ethnicity group, we calculated the normalized population-weighted disparity ($d1_i$) as the absolute difference in the annual exposure for racial/ethnic group i in the state (e_i) and the annual exposure for the state population as a whole (e_{state}) relative to the annual exposure across the contiguous United States ($e_{national}$):

$$d1_i = \frac{e_i - e_{state}}{e_{national}}. \quad (6)$$

Second, for each state, we used Equation 7 to calculate a normalized population-weighted disparity ($d2_m$) between the annual exposure for all non-Hispanic Black, non-Hispanic Asian, and Hispanic people combined (e_m), and for the non-Hispanic White population (e_{NHW}). This metric has the advantage of consistently comparing, for each state, exposures between racial/ethnic minority populations and the majority racial/ethnic group population (non-Hispanic White, 64% of the population).

$$d2_m = \frac{e_m - e_{NHW}}{e_{national}}. \quad (7)$$

Last, for each state, we averaged both metrics across the six pollutants.

Results

National Exposure Disparities by Race/Ethnicity and Income in 2010

By race/ethnicity. To investigate national disparities in exposure to criteria air pollution by race/ethnicity, we first compared national population-weighted mean exposures by U.S. Census self-reported race/ethnicity in 2010, the most recent decennial census year with available data. We first present results for differences among subpopulations (unit of analysis: racial/ethnic group), then we present differences among locations, depending on the proportion of each racial/ethnic group residents in that location (unit of analysis: census block groups binned by proportion of racial/ethnic minority residents).

Estimated national mean air pollution exposures for 2010 were higher for all three racial/ethnic minority groups than for the non-Hispanic White group for four of the six criteria pollutants (CO, NO₂, PM_{2.5}, and PM₁₀) (Table 1; Table S2–S3; Figure 1). For all six pollutants, the most-exposed group was a racial/ethnic

Table 1. Population distribution and population-weighted exposure distribution for six criteria pollutants for four main racial/ethnic groups and the national average in year 2010.

Demographic	Non-Hispanic White	Non-Hispanic Black	Hispanic	Non-Hispanic Asian	Entire population
Proportion of population	64%	12%	16%	4.6%	100%
PM _{2.5} (µg/m ³)					
10th percentile	6.1	7.9	6.5	6.7	6.3
25th percentile	7.7	9.2	7.7	8.2	7.9
50th percentile	9.3	10	9.6	9.7	9.5
Mean (SD)	9.1 (2.2)	10 (1.8)	9.4 (2.2)	9.4 (1.9)	9.3 (2.2)
75th percentile	11	11	11	11	11
90th percentile	12	13	12	12	12
NO ₂ (ppb)					
10th percentile	3.1	3.8	4.6	5.4	3.4
25th percentile	4.3	5.8	6.6	7.5	4.9
50th percentile	6.2	8.7	9.5	10	7.4
Mean (SD)	7.2 (4.1)	9.7 (5.3)	11 (6.1)	12 (5.9)	8.7 (5.1)
75th percentile	8.9	12	15	15	11
90th percentile	12.5	18	21	21	16
O ₃ (ppb)					
10th percentile	38	39	33	39	38
25th percentile	43	43	42	44	43
50th percentile	47	47	46	47	47
Mean (SD)	46 (6.0)	46 (6.1)	45 (7.2)	46 (5.9)	46 (6.2)
75th percentile	50	50	49	50	50
90th percentile	52	53	52	53	52
SO ₂ (ppb)					
10th percentile	0.91	1.0	0.83	0.79	0.95
25th percentile	1.1	1.2	1.0	1.0	1.2
50th percentile	1.5	1.6	1.3	1.2	1.5
Mean (SD)	1.6 (0.65)	1.7 (0.63)	1.4 (0.55)	1.4 (0.58)	1.6 (0.64)
75th percentile	1.9	2.1	1.7	1.7	2.0
90th percentile	2.4	2.5	2.2	2.3	2.5
PM ₁₀ (µg/m ³)					
10th percentile	12	14	15	14	13
25th percentile	14	16	17	16	15
50th percentile	17	19	20	19	18
Mean (SD)	18 (4.4)	19 (3.7)	21 (4.9)	20 (4.5)	18 (4.6)
75th percentile	21	21	23	22	22
90th percentile	23	23	28	25	24
CO (ppm)					
10th percentile	0.23	0.25	0.26	0.27	0.24
25th percentile	0.27	0.29	0.30	0.30	0.28
50th percentile	0.31	0.32	0.34	0.34	0.31
Mean (SD)	0.30 (0.057)	0.32 (0.067)	0.35 (0.079)	0.35 (0.071)	0.31 (0.066)
75th percentile	0.33	0.35	0.39	0.38	0.35
90th percentile	0.37	0.40	0.45	0.45	0.39

Note: CO, carbon monoxide; NO₂, nitrogen dioxide; O₃, ozone; PM_{2.5}, fine particulate matter with aerodynamic diameter less than or equal to 2.5 micrometers; PM₁₀ 10 micrometers, SD, standard deviation; SO₂, sulfur dioxide.

minority group: for PM_{2.5} and SO₂, national mean exposures were highest for the non-Hispanic Black population; for CO, NO₂, and O₃, the non-Hispanic Asian population; and for PM₁₀, the Hispanic population. For CO, NO₂, PM_{2.5}, and PM₁₀, national mean exposures were lowest for non-Hispanic White population; for O₃, Hispanic population; and for SO₂, non-Hispanic Asian population. Disparities between the most- and least-exposed racial/ethnic groups were largest (based on the relative disparity ratio) for NO₂ [absolute disparity: 4.6 ppb (54%), relative disparity (ratio): 1.6]; intermediate for SO₂ [0.29 ppb (19%), 1.2], PM₁₀ [3.0 µg/m³ (17%), 1.2], CO [0.044 ppm (16%), 1.1], and PM_{2.5} [1.2 µg/m³ (13%), 1.1]; and lowest for O₃ [1.6 ppb (3.6%), 1.0] (Table S4). Across the five pollutants, normalized disparities were also largest for NO₂ and smallest for O₃ for all the additional demographic groups considered (income poverty ratio, age, language, mobility, and travel time) (Table S5). Among those additional demographic groups, disparities that stand out as comparatively larger are income poverty ratio (NO₂), mobility (NO₂, CO), and travel time (NO₂) (see Figure S1; Table S5).

Sensitivity tests on robustness of conclusions based on mean values showed that, for all pollutants, the rank order (i.e., most-

to least-exposed racial/ethnic group, among the four racial/ethnic groups) was consistent throughout the exposure distributions (Figure 1). Results for the supplemental inequality metrics (Gini coefficient; between-group Atkinson Index) indicate that exposure inequality was largest for NO₂ and smallest for O₃ (Tables S6 and S7). This finding is consistent with the findings based on our primary metrics. The remaining two sensitivity tests investigated whether conclusions here are robust to uncertainty in exposure model predictions. Results reveal that the conclusions are robust to exposure model uncertainty. Results for analyzing only the population living in a census block group with a U.S. EPA monitor in 2010 were essentially the same as results using exposure model predictions: the non-Hispanic White group was the least-exposed group on average for most pollutants (CO, NO₂, PM_{2.5}, PM₁₀, and O₃), and the relative disparities by pollutant (between the most- and least-exposed group on average) were highly correlated (Spearman rank order correlation between base case and sensitivity test: 0.89) (Tables S8 and S9). The ratio between the uncertainties in estimated racial/ethnic exposure disparities and the estimated racial/ethnic disparities between the most- and least-exposed racial/ethnic groups were small: on




Figure 1. Distribution of exposure to pollutants in years 1990, 2000, and 2010, stratified by racial/ethnic group, for (A) PM_{2.5}, (B) NO₂, (C) O₃, (D) SO₂, (E) PM₁₀, and (F) CO. For all panels, the highest/lowest bound represents the 90th/10th percentile value, the box shows the 25th and 75th percentiles, and the horizontal line in the box represents the median. Color circles indicate the national population-weighted mean. PM_{2.5} has no estimates in 1990 because of a lack of monitoring data prior to 1999. Note: CO, carbon monoxide; Hispanic, Hispanic people of any race(s); NH, non-Hispanic; NO₂, nitrogen dioxide; O₃, ozone; PM_{2.5}, fine particulate matter with aerodynamic diameter less than or equal to 2.5 micrometers; PM₁₀ 10 micrometers; SO₂, sulfur dioxide.

average across the six pollutants, 0.0073 (if using absolute values of the ratio, 0.083). The largest absolute ratio was -0.17 (O₃). That result indicated that the uncertainty in the exposure model predictions was always small in comparison with the predicted racial/ethnic exposure disparities (Tables S10 and S11).

We also performed an analysis to determine whether average air pollution levels varied based on the racial/ethnic composition of a given census block group. For CO, NO₂, PM_{2.5}, and PM₁₀, average pollution levels were higher in census block groups with higher proportions of racial/ethnic minority residents (Figure 2). For O₃, estimated average levels were approximately equal across census block group bins, regardless of census block group racial/ethnic characteristics (Figure 2). For SO₂, estimated average levels were generally higher in census block group bins with the highest and lowest proportions of racial/ethnic minority residents (i.e., higher in more racially segregated census block groups) (Figure 2). This approach also reveals that the disparities were much larger for NO₂ than for other pollutants. The disparity in average air pollution levels between block groups with the highest vs. lowest deciles of proportion racial/ethnic minority residents (block groups with >88% vs. <4% racial/ethnic minority residents) was larger for NO₂ [absolute disparity: 9.4 ppb, relative disparity (ratio): 3.1] than for other pollutants [relative disparity (ratio) range: 0.8–1.4, median: 1.1] (Table S12).

Last, we investigated racial/ethnic disparities in exposure to the highest air pollution levels. First, for each racial/ethnic group

we calculated the proportion of people nationally who lived in a block group with air pollution levels above the 90th percentile for each pollutant. Averaged across all pollutants, the proportion of people nationally who lived in those highest-exposure block groups was: 9.6% for the overall population, 17% for the Hispanic population, 15% for the non-Hispanic Asian population, 12% for the non-Hispanic Black population, and 7.2% for the non-Hispanic White population. Racial/ethnic minority populations were more likely than non-Hispanic White populations to live in a census block group with air pollution levels above the 90th percentile, for all pollutants (range: $1.0 \times$ to $4.1 \times$, median: $2.1 \times$) except SO₂ ($0.88 \times$) (Figure S2; Table S13). Next, we calculated the racial-ethnic composition of the block groups with air pollution levels above the 90th percentile for each pollutants; the proportion of the population in those block groups that is non-Hispanic White is less than the national average, for all pollutants except SO₂ (Figure S3; Table S14). Racial/ethnic minority populations were also disproportionately likely to live in a census block group having *multiple* pollutants with levels above the 90th percentile. For example, the proportion of population living in a census block group with levels above the 90th percentile for four or more criteria pollutants was 5.2% for the Hispanic population (3.6 times the national population average proportion), 2.2% for the non-Hispanic Asian population (1.5 times the average), 1.9% for the non-Hispanic Black population (1.3 times the average), and 0.36% for the non-Hispanic White population (0.25 times the




Figure 2. Relationship between the proportion of racial/ethnic minority residents in census block groups and average criteria air pollution concentrations in the years 1990, 2000, and 2010 for (A) $PM_{2.5}$, (B) NO_2 , (C) O_3 , (D) SO_2 , (E) PM_{10} , and (F) CO. For each panel, the thicker portion of the line indicates the 25th to 75th percentile of census block groups, the thin line indicates the 10th to 90th percentiles, the dashed line indicates the 1st to 99th percentiles, and the diamond icon indicates the median. Note: CO, carbon monoxide; Hispanic, Hispanic people of any race(s); NH, non-Hispanic; NO_2 , nitrogen dioxide; O_3 , ozone; $PM_{2.5}$, fine particulate matter with aerodynamic diameter less than or equal to 2.5 micrometers; PM_{10} , 10 micrometers; SO_2 , sulfur dioxide.

average) (for comparison: 1.4% for the overall U.S. population) (Table S14). The ratio of the non-Hispanic White population relative to the national population average in each block group category declined monotonically as the number of pollutants above the 90th percentile increased from 0 to ≥ 4 (ratios from 1.1 to 0.25), whereas corresponding ratios increased monotonically for non-Hispanic Black (from 0.88 to 1.3) and for Hispanic populations (from 0.84 to 3.6) and increased nonmonotonically for non-Hispanic Asian populations (from 0.88 for 0 pollutants to 2.3 and 1.5 for 3 and ≥ 4 pollutants >90 th percentile, respectively) (Figure S4; Table S15).

By income. To investigate national exposure disparities by income, we first compared national mean exposures to criteria air pollution by census income category in 2010. For all pollutants except O_3 , national mean exposures were higher for lowest-income ($< \$10,000$; 7.2% of the households with income data) than for highest-income ($> \$200,000$; 4.2%) households, with all pollutants except NO_2 (and, to a lesser extent, CO and O_3) exhibiting a monotonic trend (Figure S5). (Consistent with those findings, we also found that for the remaining three pollutants (SO_2 , $PM_{2.5}$, PM_{10}), but not for O_3 , NO_2 , and CO, the most-exposed income category is the lowest-income category and the least-exposed income category is the highest-income category; see Table S16). Relative to the overall population-weighted mean exposure for all households in 2010, the absolute difference between mean exposures

among those in the lowest- vs. highest-income category households were 16% (relative to national mean exposure) higher for SO_2 , 6.6% higher for $PM_{2.5}$, and 5.2% higher for PM_{10} . For NO_2 , CO, and O_3 , exposures for lowest- and highest-income households were similar ($\sim \pm 2\%$) (Table S17). (For comparison, for NO_2 , CO, and O_3 , exposure differences between the most- and least-exposed income categories were 2.5% to 9.4%; see Table S16.)

Based on differences in average exposures between the approximate 25th and 75th percentiles for income [$\$20,000$ – $\$25,000$ (midpoint: $\$22,500$) and $\$75,000$ – $\$100,000$ (midpoint: $\$87,500$)], a $\$10,000$ increase in income was associated with an average reduction in concentration (expressed as a percent of the national mean concentration) of 0.90% for SO_2 , 0.41% for $PM_{2.5}$, 0.36% for NO_2 , and 0.22% for PM_{10} and CO, and an increase of 0.16% for O_3 . For NO_2 , the change in average exposure per $\$10,000$ increase in income was 0.59% between the 25th and 50th [$\$40,000$ – $\$45,000$ (midpoint: $\$42,500$)] percentiles, and 0.26% between the 50th and 75th percentile (Table S18).

By both race/ethnicity and income. In this section, we present exposure disparities accounting for both race/ethnicity and income together for census householders (hereafter, “households”). For all six pollutants in 2010, the absolute exposure disparity between the most- and least-exposed racial/ethnic groups was larger [on average, ~ 6 times larger; 1.1 times (i.e., 10% larger) for SO_2 , 21 times for NO_2 , and 1.4 (i.e., 40% larger) to 6.8 times for the remaining




Figure 3. Population-weighted criteria air pollution concentration in 2010 for 16 household income groups, stratified by race/ethnicity, for (A) $PM_{2.5}$, (B) NO_2 , (C) O_3 , (D) SO_2 , (E) PM_{10} , and (F) CO. For all panels, each data point represents pollution exposure for one income category and racial/ethnic group. Values plotted for household income are, for values below \$200,000 (i.e., for the first 15 income categories), the midpoint value; for the highest income category (“>\$200,000”), the value plotted is the low end of the range (\$200,000). Note: Asian, Hispanic and non-Hispanic Asian people; Black, Hispanic and non-Hispanic Black people; CO, carbon monoxide; Hispanic, Hispanic people of any race(s); NH White, non-Hispanic White people; NO_2 , nitrogen dioxide; O_3 , ozone; $PM_{2.5}$, fine particulate matter with aerodynamic diameter less than or equal to 2.5 micrometers; PM_{10} 10 micrometers; SO_2 , sulfur dioxide.

pollutants] than the absolute exposure disparity between the lowest- and highest-income categories [relative disparity: on average, ~ 1.2 times (i.e., 20% larger)]. The absolute exposure disparity between the most- and least-exposed racial/ethnic groups is 5.8 times for NO_2 , 1.1 times (i.e., 10% larger) for SO_2 , and 1.4 to 4.4 times for remaining pollutants than the absolute exposure disparity between the most- and least-exposed income categories (Table S19). For all income levels and pollutants, the most-exposed racial/ethnic group was a racial/ethnic minority group (Figure 3; Table S20). For five of the six pollutants (not SO_2 ; Figure 3), average exposures were higher on average for Black households at the approximate 75th percentile for income (income category midpoint: \$87,500) than for non-Hispanic White households at the approximate 25th percentile for income (midpoint: \$22,500). Racial/ethnic exposure disparities tended to be comparatively smaller at higher incomes than at lower incomes (except for O_3), but the size of that effect was modest. For example, the absolute exposure disparity between the most- and least-exposed racial/ethnic groups (Figure 3) was, on average, 9.5% lower for households at the approximate 75th percentile than at the approximate 25th percentile of income.

Income distributions varied by racial/ethnic group. For example, non-Hispanic White households represented 61% of the lowest income category (<\$10,000) and 85% of the highest income category (>\$200,000), vs. 23% and 3.5%, respectively, for Black households, 13% and 4.3% for Hispanic households, and 3.5% and 6.9% for Asian households (Table S21). To quantify racial/ethnic exposure disparities after accounting for racial/ethnic income

distribution variation, we calculated the absolute exposure disparity between the most- and least-exposed racial/ethnic groups within each income category in 2010 and then averaged across all 16 income categories. The resulting national absolute exposure disparity between most- and least-exposed racial/ethnic groups averaged across income categories and normalized to national mean exposure (i.e., expressed as a percent of the national mean concentration) was 58% for NO_2 , 4.5% for O_3 , 12%–17% for the remaining pollutants. Conversely, to quantify income exposure disparities after accounting for race/ethnicity, we calculated the absolute income disparity within each racial/ethnic group and averaged across the four racial/ethnic groups. The resulting national absolute exposure disparity between lowest and highest income categories normalized to national mean exposure was 15% for SO_2 , -2.9% for O_3 , and 2.7%–6.3% for the remaining pollutants (Table S22). In conclusion, the results given here, consistent with Liu (2021), indicate that racial/ethnic exposure disparities were distinct from, and larger than, exposure disparities by income.

Racial/ethnic Exposure Disparities by State and by Urbanicity in 2010

By state. We explored how exposures varied by state, pollutant, and racial/ethnic group in 2010 (Figure 4). The analysis separately considers the District of Columbia (DC) plus the 48 states of the contiguous United States (hereafter, “states” refers to 48 states and DC, a total of 49 geographic units in state-level related calculations). There are 294 pollutant-state combinations (6 pollutants \times 49 units) and




Figure 4. State racial/ethnic disparities in pollution exposure in 2010, showing the difference between (1) NH White vs. state average, (2) NH Black vs. state average, (3) Hispanic vs. state average, (4) NH Asian vs. state average, and (5) Minority vs. NH White for the six pollutants. (A) PM_{2.5}, (B) NO₂, (C) O₃, (D) SO₂, (E) PM₁₀, and (F) CO, and (G) average across the six pollutants. Columns 1–4: exposure disparity relative to state average; calculated as mean exposure for a racial/ethnic group in that state minus the overall mean for that state, then divided by the national overall mean. Column 5: exposure disparity for racial/ethnic minorities relative to the racial/ethnic majority group; calculated as mean exposure for racial/ethnic minorities minus mean exposure for non-Hispanic White people, then divided by the national overall mean. Mean values are population-weighted. States displayed in white indicate that the disparity is within $\pm 5\%$ of the national overall mean. Purple shading indicates that mean exposures are higher than average by more than 5% of the national overall mean (columns 1–4) or that mean exposures are higher for racial/ethnic minorities than for non-Hispanic White people, by more than 5% of the national overall mean (column 5). Orange shading indicates the reverse: mean exposures are lower than average for that group (columns 1–4) or mean exposures are lower for racial/ethnic minorities than for non-Hispanic White people (column 5), and the disparity is greater than 5% of the national overall mean. See Excel Table S1 for corresponding numeric data. Note: CO, carbon monoxide; Hispanic, Hispanic people of any race(s); NH, non-Hispanic; NO₂, nitrogen dioxide; O₃, ozone; PM_{2.5}, fine particulate matter with aerodynamic diameter less than or equal to 2.5 micrometers; PM₁₀, 10 micrometers; SO₂, sulfur dioxide.

1,176 pollutant-state-group combinations (294 pollutant-states \times 4 racial/ethnic groups). For this section, we define $\pm 5\%$ (all percentages used in this section were expressed as a percent of the national mean exposure in 2010) as “similar to” and therefore report examples where exposures differ from the average by $>5\%$ (or, in a sensitivity test, $>20\%$). For example, “ $>5\%$ lower-than-average” means the exposure is lower than state average by an amount greater than 5% of the pollutant’s national mean.

Overall, several spatial patterns emerge across states. First, racial/ethnic exposure disparities were ubiquitous among U.S. states. In all 48 states and DC in 2010, one or more racial/ethnic groups experienced exposures disparities $>5\%$ of the pollutant’s national mean. Second, racial/ethnic minority populations within states were much more likely to have been more exposed vs. less exposed than the state average; in contrast, none of the non-Hispanic White populations within states experienced exposures $>5\%$ above the state average. Third, having exposures $>5\%$ lower than average within a state was much more likely to happen for non-Hispanic White populations than for racial/ethnic minority (non-Hispanic Black, non-Hispanic Asian, and Hispanic populations combined) populations (Figure 4, right column). Fourth, racial/ethnic exposure disparities were most pronounced (in magnitude and with regard to the number of states affected)

for NO₂, whereas mean O₃ exposures were similar among all racial/ethnic groups in all states.

Those findings reflect underlying trends across states, pollutants, and racial/ethnic groups. For example, for the non-Hispanic White group, 87% of the 294 pollutant-states had exposures that were similar ($\pm 5\%$) to the average, 13% had exposures $>5\%$ less than average, and none were $>5\%$ greater than average. In contrast, for exposures for the three racial/ethnic minority groups, 42% (of 882 pollutant-state-group combinations) were $>5\%$ greater than average, 55% were $\pm 5\%$ of the average, and only 4% were $>5\%$ lower than average. Thus, within individual states, the non-Hispanic White group was exposed to pollution levels that were similar to or cleaner than average, whereas the three racial/ethnic minority groups were more likely to be exposed to dirtier rather than cleaner pollution levels. For example, averaged across pollutants, the proportion of the states for which exposures were $>5\%$ greater than average is 73% for non-Hispanic Black populations, 57% for Hispanic populations, 35% for non-Hispanic Asian populations, and zero for non-Hispanic White populations.

The three racial/ethnic minority groups were disproportionately likely to be the *most*-exposed group, and disproportionately *unlikely* to be the *least*-exposed group of the four racial/ethnic groups across

states. For example, the most-exposed group (for all cases, not just cases >5% greater than average) was the non-Hispanic Black group for 45% of the 294 pollutant-states, the Hispanic group for 29%, the non-Hispanic Asian group for 18%, and non-Hispanic White group for 7.5%. In contrast, the least-exposed group was rarely a racial/ethnic minority group (of the 294 pollutant-states, ~8%, each for the non-Hispanic Black and the Hispanic groups, 15% for the non-Hispanic Asian group) and was usually (70% of 294 pollutant-states) the non-Hispanic White group.

In a sensitivity test, we changed the analysis threshold to exposures >20% (rather than >5%) greater than average and similarly found that the air pollution disproportionately impacted racial/ethnic minority groups. For example, exposure disparities >20% of national mean exposure for one or more pollutant-groups occurred for 67% of states (Figure 4, left four columns for six pollutants, darkest two purple shades), further emphasizing that disparities were widespread across states in 2010.

Figure 4 reveals differences among states. For example, the four most populous states (California, Florida, New York, and Texas), all have large, racially/ethnically diverse urban areas. However, average disparities between racial/ethnic minority populations and non-Hispanic White populations (Figure 4, bottom right) were notably larger (on average, 6 times larger) for California and New York than for Florida and Texas (Excel Table S1). Some small, relatively rural states also had substantial exposure disparities; examples include NO₂ in Nebraska (19%) and PM_{2.5} in Nebraska (8.1%).

By urbanicity. We investigated racial/ethnic and income-based exposure disparities in 2010 separately for block groups that were defined as urban (89% of the population) vs. rural (11% of the population). Overall, urban populations experienced larger exposure than that of rural populations for all pollutants (Table S23).

The most- and least-exposed of the four racial/ethnic groups differed between urban and rural areas for SO₂ and O₃. For SO₂, the most-exposed racial/ethnic group was the non-Hispanic Black group in urban areas and the non-Hispanic White group in rural areas. For O₃, the most-exposed racial/ethnic group was the non-Hispanic Asian group in urban areas and the non-Hispanic White group in rural areas. For the remaining four pollutants, the most-exposed group was a racial/ethnic minority group in both urban and rural areas (Table S24).

The racial/ethnic exposure disparities were generally larger for urban than for rural block groups. Specifically, the average exposure disparity between the most- and least-exposed racial/ethnic group was 5.5 times larger for absolute disparity [1.2 times for relative disparity (ratio between relative disparity in urban areas and relative disparity in rural areas)] for urban block groups than for rural block groups for NO₂, 3.1 times (1.0 times) larger for O₃, 2.4 times (1.1 times) larger for CO, 1.3 times (1.0 times) larger for SO₂, and 1.2 times (1.0 times) larger for PM₁₀. [Here, 1.2 times larger would indicate 20% larger, and 1.0 times larger would indicate 0% larger (i.e., not larger).] In contrast, for PM_{2.5}, the average racial/ethnic exposure disparity was 1.2 times (1.0 times) larger for rural block groups than for urban block groups (Table S24).

Exposure disparities by income category were also larger in urban than in rural areas. Absolute exposure disparities between lowest and highest income category were 1.1 times (PM_{2.5}) to 25 times (O₃) (median: 3.5 times) greater [for relative disparity (ratio), range: 0.98–1.1 times; median: 1.0 times] in urban than in rural areas (Table S25). Of the 12 pollutant-urbanicity categories (6 pollutants × 2 urbanicities), exposures were higher for the lowest-income category than for the highest-income category in all cases except for O₃ in urban areas and NO₂ in rural areas (Table S25).

Changes in National Exposures and Exposure Disparities from 1990 to 2010

Criteria air pollution levels have declined in the United States in the decades following the 1990 Clean Air Act amendments (U.S. EPA 2020) (Table S26). To investigate whether these reductions have led to reductions in racial/ethnic exposure disparities, we compared average exposures by racial/ethnic group from 1990 to 2010, for five of the pollutants. Exposure model results for PM_{2.5} were available only from 2000 to 2010, so those results are presented separately.

National mean pollution levels for all six pollutants fell over the study period. For example, from 1990 to 2010, the national mean exposures decreased for all five pollutants by an average of 40% relative to national mean exposures in 1990 [range: –6% (O₃) to –71% (SO₂); –34% to –55% for remaining three pollutants]. PM_{2.5} exposures decreased 29% from 2000 to 2010 (Table S27).

Average racial/ethnic exposure disparities also declined from 1990 to 2010. The amount of change depends in part on whether one considers *absolute* or *relative* disparities. In terms of *absolute* disparities, the disparities between the most- and least-exposed racial/ethnic groups decreased on average by 69% relative to absolute disparity in 1990 across the five pollutants. The largest change was an 88% decrease for CO disparities [0.40 ppm in 1990, 0.044 ppm in 2010, a 0.35 ppm (i.e., 88%) change], and the smallest change was a 54% decrease for NO₂ [9.8 ppb (1990), 4.6 ppb (2010), a 5.3 ppb (54%) change]. From 2000 to 2010, PM_{2.5} disparities decreased by 35% [1.9 µg/m³ (2000), 1.2 µg/m³ (2010), a 0.66 µg/m³ change] (Table S28).

In terms of *relative* disparities, the greatest change during the period 1990–2010 was a decrease for CO [disparities: 1.63 (1990), 1.15 (2010), 0.71 times (i.e., 29% reduction)], and the smallest was a decrease for O₃ [1.10 (1990), 1.04 (2010), 0.95 times (i.e., 5% reduction)]; remaining three pollutants (NO₂, PM₁₀, SO₂) were between 0.94 times and 0.95 times (i.e., 5%–6% reduction in relative disparity). PM_{2.5} relative disparity remained nearly constant (0.99 times) during the period 2000–2010 (Table S28).

Absolute disparities between census block group bins with the highest vs. lowest deciles of proportions of racial/ethnic minority residents (90th–100th vs. 1st–10th percentiles in Figure 2) decreased for CO, NO₂, PM₁₀, and SO₂ [by 10% (SO₂) to 164% (CO)] and decreased by 17% from 2000 to 2010 for PM_{2.5} (Table S29). For O₃, absolute disparities increased slightly, from –1.7 ppb in 1990 to –1.3 ppb (which is 0.74% of the national mean exposure) in 2010.

In addition to national changes, we investigated changes in absolute racial/ethnic exposure disparities from 1990 to 2010 by state and by urban vs. rural areas. Most states (>75%) experienced a reduction in racial/ethnic exposure disparities for pollutants, except for PM₁₀ (and, except for PM_{2.5} during the period 2000–2010) (Figure S6; Table S30). Urban areas experienced larger reductions in racial/ethnic exposure disparities than did rural areas for NO₂ and PM₁₀ (13 times larger reductions in urban areas, for both pollutants), CO (2.4 times), and SO₂ (1.2 times). Conversely, PM_{2.5} (during the period 2000–2010) and O₃ (during the period 1990–2010) had larger reductions in absolute racial/ethnic disparities for rural than for urban (2.4 times and 3.4 times larger in rural areas, respectively) (Figure S7; Table S31).

Finally, we investigated whether the changes in absolute racial/ethnic exposure disparities from 1990 to 2010 were more attributable to changes in air pollution levels or to changes in demographic patterns (migration, immigration, and other factors). Based on a counterfactual analysis, reductions in racial/ethnic exposure disparities between the most- and least-exposed racial/ethnic groups were mainly attributable to changes in air pollution levels rather than to changes in demographic patterns. On average across all pollutants, 87% of the reduction in the absolute racial/ethnic disparity

metric was attributable to changes in air pollution levels from 1990 to 2000 (excluding PM_{2.5} based on lack of available data), and 97% from 2000 to 2010 (Tables S32 and S33).

Discussion

Our research provides the first national investigation of air pollution exposure disparities by income and race/ethnicity for all criteria pollutants (except lead). Our results reveal trends by pollutant and across time and space.

In 2010, on average nationally, racial/ethnic minority populations were exposed to higher average levels of transportation-related air pollution (CO, NO₂) and particulate matter (PM_{2.5}, PM₁₀) than were non-Hispanic White populations. This finding, which holds even after accounting for uncertainties in the predictions from exposure models, is consistent with prior national studies of NO₂, PM_{2.5}, and PM₁₀ (Clark et al. 2017; Kravitz-Wirtz et al. 2016; Mikati et al. 2018; Tessum et al. 2019; Colmer et al. 2020). Disparities for the remaining pollutants (CO, O₃, and SO₂) had not been previously studied in detail for the national population, and few studies have considered how disparities for any pollutant have changed across 20 y (Kravitz-Wirtz et al. 2016; Bullard et al. 2008).

Our findings on “which group was most exposed over time?” (on average, nationally) varied by pollutant, but in all six cases the most exposed group was a racial/ethnic minority group. That result is consistent with prior national studies, which have reported, for example, highest average NO₂ exposures for Hispanic Black and non-Hispanic Asian populations (Clark et al. 2017) and highest average proximities to industrial PM_{2.5} emissions (Mikati et al. 2018) and highest average exposures to industrial air toxins (Ard 2015) for non-Hispanic Black populations.

We found that racial/ethnic minority populations were more than two times as likely than non-Hispanic white populations to live in a census block group with highest air pollution levels (above 90th percentile) on average. Those results are consistent with existing literature on disproportionate environmental risks for racial/ethnic minority populations (Collins 2016) and on groups or locations with higher risks for one environmental factor having higher risks for other factors, too (Morello-Frosch and Lopez 2006; Su et al. 2012).

We found that air pollution exposures were generally higher for lower-income than for higher-income households (for all pollutants except O₃). This finding is consistent with previous national research [e.g., for industrial PM_{2.5} emissions (Mikati et al. 2018), industrial air toxins (Ard 2015), and PM_{2.5} and NO₂ (Clark et al. 2014; Kravitz-Wirtz et al. 2016)]. Additionally, we found that, in 2010, absolute racial/ethnic exposure disparities were distinct from and were larger than (on average, ~6 times larger than) absolute exposure disparities by income. The findings here are *inconsistent* with the idea that racial/ethnic exposure disparities can be explained by, or are “merely” a reflection of, income disparities among racial/ethnic groups (Liu 2021).

The findings from this study can be used to compare relative exposure disparities for different criteria air pollutants in a consistent way, providing additional context for previous studies of single pollutant. We found that in 2010, relative racial/ethnic exposure disparities (i.e., ratios of average exposures between the most- and least-exposed groups) were largest for NO₂ and smallest for O₃. Relative income-based exposure disparities (i.e., ratios of average exposures between the lowest and highest income groups), although smaller than racial/ethnic exposure disparities for each pollutant, were largest for SO₂ and smallest (and similar) for NO₂, CO, and O₃. (These results provide information on the rank order of relative disparities in air pollution levels by pollutant; information on the rank order of relative disparities in associated health impacts by pollutant would require further analysis, as discussed next.)

Exposure disparities often connect with health disparities. Based on the magnitude of exposure disparities (e.g., 2010 national average PM_{2.5} exposures for non-Hispanic Black people were 1.0 µg/m³ higher than average), the resulting health disparities may be substantial (Liu 2021). Future research could usefully extend our exposure disparity results to provide rigorous, comprehensive investigation of the associated health impacts.

State-level results may be especially useful given the important role that states play in air pollution and environmental policy making (Abel et al. 2015). Exposures >5% greater than the national mean exposure within states were common for racial/ethnic minority populations, but not for non-Hispanic White populations. Indeed, we found no case (no state and no pollutant) for which the non-Hispanic White group experiences exposures >5% greater than the state average. This finding reflects disparity in exposure as well as non-Hispanic White populations representing a large percentage of states' populations. Exposure disparities varied substantially among states, even among states with similar characteristics (e.g., urbanicity, population, region). Our results emphasize differences among states in the level and makeup of exposure disparities, yet also demonstrate that exposure disparities were ubiquitous, including both large and small states, and states in all regions of the United States, in 2010.

Our analyses by urbanicity were in part motivated by and reflect urban–rural differences in demographics and air pollution levels (Clark et al. 2017; Mikati et al. 2018; Rosofsky et al. 2018). Racial/ethnic disparities were larger for urban block groups for all pollutants except PM_{2.5}. Of the six pollutants, the largest ratio between urban and rural racial/ethnic absolute disparities (5.5 times larger) was for NO₂ (Table S24). The NO₂ results are consistent with prior research (Clark et al. 2014, 2017). Over our study period, reductions in absolute racial/ethnic exposure disparities for PM_{2.5} and O₃ were larger for rural than for urban areas. Analyzing urban and rural block groups separately, exposures were mostly higher for the lowest income category than the highest. Absolute income-based exposure disparities were also 7.5 times larger on average in urban than in rural areas.

The results by state and by urbanicity reflect that exposure disparities differ by spatial units (e.g., urban/rural, and by state); future research could explore these aspects further, for example, through a spatial decomposition of national exposure disparities.

Regulations such as the 1990 Clean Air Act Amendments have achieved substantial reductions in the concentrations of many pollutants. Our analysis reveals that, as a concomitant benefit, falling pollution levels have reduced absolute exposure disparities among racial/ethnic groups. These findings are consistent with previous national research for NO₂, PM_{2.5}, and industrial air toxins (Ard 2015; Clark et al. 2017; Kravitz-Wirtz et al. 2016; Colmer et al. 2020). We found that a larger share of the racial/ethnic exposure disparity reduction was attributable to air pollution level reduction rather than changes in demographic and residential patterns.

Our study described patterns in exposure disparities but did not investigate aspects such as underlying causes or ethical or legal aspects. Systemic racism and racial segregation are two major causes discussed in multiple previous studies (Jones et al. 2014; Morello-Frosch and Lopez 2006; Schell et al. 2020). Future longitudinal research could further investigate the underlying causes of exposure disparities. One important dimension not considered here is responsibility for generating pollution. Recent analysis suggests that Hispanic and Black populations have disproportionately lower consumption of goods and services whose emissions lead to PM_{2.5} air pollution (Tessum et al. 2019).

Our study has several limitations. The finest spatial scale of publicly available census demographic data for race/ethnicity and

income, at consistent spatial geographies across time (Manson et al. 2019), is at the census block group level; race/ethnicity across income data is at census tract level with slightly different categories (see “Methods” section); we were unable to assess disparities at finer spatial scales than publicly available census data; we only included the four main racial/ethnic groups. Our analysis of exposures by income is based on national-level income distribution data and does not account for spatial variations in income distributions (e.g., among states). Our disparity estimates do not account for *a*) daily mobility for work, shopping, recreation, and other activities; *b*) direct indoor exposure to indoor sources such as cigarette smoke, cooking, or incense; *c*) indoor–outdoor relationships in pollution levels, such as particle losses during airflow in ducts or ozone losses to indoor surfaces; or *d*) occupational exposures. Our exposure disparity estimates were limited by uncertainties in the CACES exposure model predictions and in census demographic data. Our uncertainty analysis (but not our main analysis) was limited to U.S. EPA monitoring locations; we were not able to test potential exposure errors at locations without monitors on the national scale. However, sensitivity analyses (Results section) indicate that the general results are robust to model uncertainty.

To our knowledge, our study provides the first national analysis of air pollution exposure disparities among income and racial/ethnic groups, for all criteria pollutants (except Pb), including trends across time (by decade, 1990–2010) and spatial location (by state and for urban vs. rural areas). On average, exposures were generally higher for racial/ethnic minority populations than for non-Hispanic White populations. Among pollutants, national racial/ethnic exposure disparities were largest for NO₂ and smallest for O₃. Exposures were also, on average, higher for the lowest-income households than for the highest-income households. However, exposure disparities by race/ethnicity were not explained by disparities in income. Racial/ethnic exposure disparities declined from 1990 to 2010 (on an absolute basis, and to a lesser extent, on a relative basis), but still existed in all states in 2010.

Acknowledgments

J.L. conducted the research and most of the analysis; developed the methods; visualized the data; and wrote, reviewed, and edited the text. M.B. and L.P.C. designed the research; performed the research; analyzed the data; and wrote, reviewed, and edited the text. A.H., S.Y.K., A.L.R., L.S., and A.A.S. conceptualized and designed the research, oversaw the methods and analysis, and reviewed and edited the writing. J.D.M. conceptualized and supervised the project; designed the research, methods, and analysis; and wrote, reviewed, and edited the text. A.L.R. and J.D.M. acquired the funding.

The authors thank K. Harper, Harper Health & Science Communications, LLC, for providing editorial support in accordance with Good Publication Practice (GPP3) guidelines. We also thank the CACES Science Advisory Committee (SAC) for feedback on earlier versions of this research.

This publication was developed as part of the Center for Air, Climate, and Energy Solutions (CACES), which was supported under Assistance Agreement No. R835873 awarded by the U.S. EPA. It has not been formally reviewed by the U.S. EPA. The views expressed in this document are solely those of authors and do not necessarily reflect those of the U.S. EPA. The U.S. EPA does not endorse any products or commercial services mentioned in this publication.

All data used are publicly available. Demographic data are available via IPUMS NHGIS (www.nhgis.org); air pollution estimates are available via the U.S. EPA CACES project (www.caces.us).

References

- Abel TD, Salazar DJ, Robert P. 2015. States of environmental justice: redistributive politics across the United States, 1993–2004. *Rev Policy Res* 32(2):200–225, <https://doi.org/10.1111/ropr.12119>.
- Ard K. 2015. Trends in exposure to industrial air toxins for different racial and socioeconomic groups: a spatial and temporal examination of environmental inequality in the U.S. from 1995 to 2004. *Soc Sci Res* 53:375–390, PMID: 26188461, <https://doi.org/10.1016/j.ssresearch.2015.06.019>.
- Bechle MJ, Millet DB, Marshall JD. 2015. National spatiotemporal exposure surface for NO₂: monthly scaling of a satellite-derived land-use regression, 2000–2010. *Environ Sci Technol* 49(20):12297–12305, PMID: 26397123, <https://doi.org/10.1021/acs.est.5b02882>.
- Bowe B, Xie Y, Yan Y, Al-Aly Z. 2019. Burden of cause-specific mortality associated with PM_{2.5} air pollution in the United States. *JAMA Netw Open* 2(11):e1915834, PMID: 31747037, <https://doi.org/10.1001/jamanetworkopen.2019.15834>.
- Brulle RJ, Pellow DN. 2006. Environmental justice: human health and environmental inequalities. *Annu Rev Public Health* 27(1):103–124, PMID: 16533111, <https://doi.org/10.1146/annurev.publhealth.27.021405.102124>.
- Bullard RD, Mohai P, Saha R, Wright B. 2008. Toxic wastes and race at twenty: why race still matters after all of these years. *Environ Law* 38(2):371–411. <http://www.jstor.org/stable/43267204>.
- Bullock C, Ard K, Saalman G. 2018. Measuring the relationship between state environmental justice action and air pollution inequality, 1990–2009. *Rev Policy Res* 35(3):466–490, <https://doi.org/10.1111/ropr.12292>.
- Clark LP, Millet DB, Marshall JD. 2014. National patterns in environmental injustice and inequality: outdoor NO₂ air pollution in the United States. *PLoS One* 9(4):e94431, PMID: 24736569, <https://doi.org/10.1371/journal.pone.0094431>.
- Clark LP, Millet DB, Marshall JD. 2017. Changes in transportation-related air pollution exposures by race-ethnicity and socioeconomic status: outdoor nitrogen dioxide in the United States in 2000 and 2010. *Environ Health Perspect* 125(9):097012, PMID: 28930515, <https://doi.org/10.1289/EHP959>.
- Clean Air Scientific Advisory Committee. 2010. Letter to Hon. Lisa P. Jackson, administrator, Environmental Protection Agency, Re: CASAC Review of Policy Assessment for the Review of the PM NAAQS – Second external review draft. [https://yosemite.epa.gov/sab/sabproduct.nsf/CCF9F4C0500C500F8525779D0073C593/\\$File/EPA-CASAC-10-015-unsigned.pdf](https://yosemite.epa.gov/sab/sabproduct.nsf/CCF9F4C0500C500F8525779D0073C593/$File/EPA-CASAC-10-015-unsigned.pdf) [accessed 18 October 2021].
- Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, et al. 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 389(10082):1907–1918, PMID: 28408086, [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6).
- Collins MB, Munoz I, Jaja J. 2016. Linking “toxic outliers” to environmental justice communities. *Environ Res Lett* 11(1):015004, <https://doi.org/10.1088/1748-9326/11/1/015004>.
- Colmer J, Hardman I, Shimshack J, Voorheis J. 2020. Disparities in PM_{2.5} air pollution in the United States. *Science* 369(6503):575–578, PMID: 32732425, <https://doi.org/10.1126/science.aaz9353>.
- Darrow LA, Klein M, Strickland MJ, Mulholland JA, Tolbert PE. 2011. Ambient air pollution and birth weight in full-term infants in Atlanta, 1994–2004. *Environ Health Perspect* 119(5):731–737, PMID: 21156397, <https://doi.org/10.1289/ehp.1002785>.
- Di Q, Amini H, Shi L, Kloog I, Silvern R, Kelly J, et al. 2020. Assessing NO₂ concentration and model uncertainty with high spatiotemporal resolution across the contiguous United States using ensemble model averaging. *Environ Sci Technol* 54(3):1372–1384, PMID: 31851499, <https://doi.org/10.1021/acs.est.9b03358>.
- Di Q, Wang Y, Zanobetti A, Wang Y, Koutrakis P, Choirat C, et al. 2017. Air pollution and mortality in the Medicare population. *N Engl J Med* 376(26):2513–2522, PMID: 28657878, <https://doi.org/10.1056/NEJMoa1702747>.
- Downey L, Dubois S, Hawkins B, Walker M. 2008. Environmental inequality in metropolitan America. *Organ Environ* 21(3):270–294, PMID: 19960094, <https://doi.org/10.1177/1086026608321327>.
- Evans GW, Kantrowitz E. 2002. Socioeconomic status and health: the potential role of environmental risk exposure. *Annu Rev Public Health* 23(1):303–331, PMID: 11910065, <https://doi.org/10.1146/annurev.publhealth.23.112001.112349>.
- Fann N, Coffman E, Hajat A, Kim SY. 2019. Change in fine particle-related premature deaths among US population subgroups between 1980 and 2010. *Air Qual Atmos Health* 12(6):673–682, <https://doi.org/10.1007/s11869-019-00686-9>.
- Gee GC, Payne-Sturges DC. 2004. Environmental health disparities: a framework integrating psychosocial and environmental concepts. *Environ Health Perspect* 112(17):1645–1653, PMID: 15579407, <https://doi.org/10.1289/ehp.7074>.
- Goldberg DL, Gupta P, Wang K, Jena C, Zhang Y, Lu Z, et al. 2019. Using gap-filled MAIAC AOD and WRF-Chem to estimate daily PM_{2.5} concentrations at 1 km resolution in the Eastern United States. *Atmos Environ* 199:443–452, <https://doi.org/10.1016/j.atmosenv.2018.11.049>.
- Hajat A, Hsia C, O’Neill MS. 2015. Socioeconomic disparities and air pollution exposure: a global review. *Curr Environ Health Rep* 2(4):440–450, PMID: 26381684, <https://doi.org/10.1007/s40572-015-0069-5>.

- Harper S, Ruder E, Roman HA, Geggel A, Nweke O, Payne-Sturges D, et al. 2013. Using inequality measures to incorporate environmental justice into regulatory analyses. *Int J Environ Res Public Health* 10(9):4039–4059, PMID: 23999551, <https://doi.org/10.3390/ijerph10094039>.
- Independent Particulate Matter Review Panel. 2020. The need for a tighter particulate-matter air-quality standard. *N Engl J Med* 383:680–683, PMID: 32521130, <https://doi.org/10.1056/NEJMs2011009>.
- Jones MR, Diez-Roux AV, Hajat A, Kershaw KN, O'Neill MS, Guallar E, et al. 2014. Race/ethnicity, residential segregation, and exposure to ambient air pollution: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Public Health* 104(11):2130–2137, PMID: 25211756, <https://doi.org/10.2105/AJPH.2014.302135>.
- Kim SY, Bechle M, Hankey S, Sheppard L, Szpiro AA, Marshall JD. 2020. Concentrations of criteria pollutants in the contiguous U.S., 1979–2015: role of prediction model parsimony in integrated empirical geographic regression. *PLoS One* 15(2):e0228535, PMID: 32069301, <https://doi.org/10.1371/journal.pone.0228535>.
- Kravitz-Wirtz N, Crowder K, Hajat A, Sass V. 2016. The long-term dynamics of racial/ethnic inequality in neighborhood air pollution exposure, 1990–2009. *Du Bois Rev* 13(2):237–259, PMID: 28989341, <https://doi.org/10.1017/S1742058X16000205>.
- Laden F, Schwartz J, Speizer FE, Dockery DW. 2006. Reduction in fine particulate air pollution and mortality: extended follow-up of the Harvard Six Cities study. *Am J Respir Crit Care Med* 173(6):667–672, PMID: 16424447, <https://doi.org/10.1164/rccm.200503-4430C>.
- Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. 2015. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525(7569):367–371, PMID: 26381985, <https://doi.org/10.1038/nature15371>.
- Liu J. 2021. Auxiliary Analysis of Air Pollution Disparities in the U.S. <https://bit.ly/3B4PhKa>. [accessed 20 October 2021].
- Manson S, Schroeder J, Van Riper D, Ruggles S. 2019. IPUMS National Historical Geographic Information System: Version 14.0 [Database]. <http://doi.org/10.18128/D050.V14.0> [accessed 24 May 2020].
- Marshall JD, Swor KR, Nguyen NP. 2014. Prioritizing environmental justice and equality: diesel emissions in Southern California. *Environ Sci Technol* 48(7):4063–4068, PMID: 24559220, <https://doi.org/10.1021/es405167f>.
- Mikati I, Benson AF, Luben TJ, Sacks JD, Richmond-Bryant J. 2018. Disparities in distribution of particulate matter emission sources by race and poverty status. *Am J Public Health* 108(4):480–485, PMID: 29470121, <https://doi.org/10.2105/AJPH.2017.304297>.
- Mohai P, Pellow D, Roberts JT. 2009. Environmental justice. *Annu Rev Environ Resour* 34(1):405–430, <https://doi.org/10.1146/annurev-environ-082508-094348>.
- Morello-Frosch R, Lopez R. 2006. The riskscape and the color line: examining the role of segregation in environmental health disparities. *Environ Res* 102(2):181–196, PMID: 16828737, <https://doi.org/10.1016/j.envres.2006.05.007>.
- Novotny EV, Bechle MJ, Millet DB, Marshall JD. 2011. National satellite-based land-use regression: NO₂ in the United States. *Environ Sci Technol* 45(10):4407–4414, PMID: 21520942, <https://doi.org/10.1021/es103578x>.
- Office of Air Quality Planning and Standards. 2008. Ambient Air Monitoring Strategy for State, Local, and Tribal Air Agencies. <https://www.epa.gov/amtic/ambient-air-monitoring-strategy-state-local-and-tribal-air-agencies>. [accessed 01 December 2021].
- Paul KC, Haan M, Mayeda ER, Ritz BR. 2019. Ambient air pollution, noise, and late-life cognitive decline and dementia risk. *Annu Rev Public Health* 40:203–220, PMID: 30935305, <https://doi.org/10.1146/annurev-publhealth-040218-044058>.
- Pope CA, Coleman N, Pond ZA, Burnett RT. 2020. Fine particulate air pollution and human mortality: 25+ years of cohort studies. *Environ Res* 183:108924, PMID: 31831155, <https://doi.org/10.1016/j.envres.2019.108924>.
- Pope CA 3rd, Ezzati M, Dockery DW. 2009. Fine-particulate air pollution and life expectancy in the United States. *N Engl J Med* 360(4):376–386, PMID: 19164188, <https://doi.org/10.1056/NEJMs0805646>.
- Ratcliffe M, Burd C, Holder K, Fields A. 2016. Defining Rural at the U.S. Census Bureau (U.S. Census Bureau). https://www2.census.gov/geo/pdfs/reference/ua/Defining_Rural.pdf [accessed 7 July 2020].
- Rivas I, Basagaña X, Cirach M, López-Vicente M, Suades-González E, Garcia-Esteban R, et al. 2019. Association between early life exposure to air pollution and working memory and attention. *Environ Health Perspect* 127(5):57002, PMID: 31070940, <https://doi.org/10.1289/EHP3169>.
- Rosofsky A, Levy JI, Zanobetti A, Janulewicz P, Fabian MP. 2018. Temporal trends in air pollution exposure inequality in Massachusetts. *Environ Res* 161(2018):76–86, PMID: 29101831, <https://doi.org/10.1016/j.envres.2017.10.028>.
- Salazar DJ, Clauson S, Abel TD, Clauson A. 2019. Race, income, and environmental inequality in the U.S. states, 1990–2014. *Soc Sci Q* 100(3):592–603, <https://doi.org/10.1111/ssqu.12608>.
- Schell CJ, Dyson K, Fuentes TL, Des Roches S, Harris NC, Miller DS, et al. 2020. The ecological and evolutionary consequences of systemic racism in urban environments. *Science* 369(6510):, PMID: 32792461, <https://doi.org/10.1126/science.aay4497>.
- Shi L, Zanobetti A, Kloog I, Coull BA, Koutrakis P, Melly SJ, et al. 2016. Low-concentration PM_{2.5} and mortality: estimating acute and chronic effects in a population-based study. *Environ Health Perspect* 124(1):46–52, PMID: 26038801, <https://doi.org/10.1289/ehp.1409111>.
- Stanaway JD, Afshin A, Gakidou E, Lim SS, Abate D, Abate KH, et al. 2018. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 392(10159):1923–1994, PMID: 30496105, [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6).
- Stieb DM, Chen L, Eshoul M, Judek S. 2012. Ambient air pollution, birth weight and preterm birth: a systematic review and meta-analysis. *Environ Res* 117(2012):100–111, PMID: 22726801, <https://doi.org/10.1016/j.envres.2012.05.007>.
- Su JG, Jerrett M, Morello-Frosch R, Jesdale BM, Kyle AD. 2012. Inequalities in cumulative environmental burdens among three urbanized counties in California. *Environ Int* 40:79–87, PMID: 22280931, <https://doi.org/10.1016/j.envint.2011.11.003>.
- Tessum CW, Apte JS, Goodkind AL, Muller NZ, Mullins KA, Paoletta DA, et al. 2019. Inequity in consumption of goods and services adds to racial-ethnic disparities in air pollution exposure. *Proc Natl Acad Sci U S A* 116(13):6001–6006, PMID: 30858319, <https://doi.org/10.1073/pnas.1818859116>.
- Underwood E. 2017. The polluted brain. *Science* 355(6323):342–345, PMID: 28126768, <https://doi.org/10.1126/science.355.6323.342>.
- U.S. Census Bureau. 2021. Poverty Thresholds. <https://www.census.gov/data/tables/time-series/demo/income-poverty/historical-poverty-thresholds.html> [accessed 5 November 2021].
- U.S. EPA (U.S. Environmental Protection Agency). 2016. EPA Downscaler Model for Predicting Daily Air Pollution. https://19january2017snapshot.epa.gov/air-research/downscaler-model-predicting-daily-air-pollution_.html. [accessed 20 October 2021].
- U.S. EPA. 2019. Integrated Science Assessment (ISA) for Particulate Matter (Final Report, December 2019). <https://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=347534>. [accessed 13 October 2021].
- U.S. EPA. 2020. Our Nation's Air: Status and Trends through 2019. <https://gispub.epa.gov/air/trendsreport/2020/> [accessed 7 July 2020].
- U.S. EPA. 2020. Review of the National Ambient Air Quality Standards for Particulate Matter. <https://www.federalregister.gov/documents/2020/04/30/2020-08143/review-of-the-national-ambient-air-quality-standards-for-particulate-matter>. [accessed 18 October 2021].
- Van Donkelaar A, Martin RV, Li C, Burnett RT. 2019. Regional estimates of chemical composition of fine particulate matter using a combined geoscience-statistical method with information from satellites, models, and monitors. *Environ Sci Technol* 53(5):2595–2611, PMID: 30698001, <https://doi.org/10.1021/acs.est.8b06392>.
- Young MT, Bechle MJ, Sampson PD, Szpiro AA, Marshall JD, Sheppard L, et al. 2016. Satellite-based NO₂ and model validation in a national prediction model based on universal kriging and land-use regression. *Environ Sci Technol* 50(7):3686–3694, PMID: 26927327, <https://doi.org/10.1021/acs.est.5b05099>.
- Zanobetti A, Schwartz J. 2009. The effect of fine and coarse particulate air pollution on mortality: a national analysis. *Environ Health Perspect* 117(6):898–903, PMID: 19590680, <https://doi.org/10.1289/ehp.0800108>.
- Zwickl K, Ash M, Boyce JK. 2014. Regional variation in environmental inequality: industrial air toxics exposure in U.S. cities. *Ecol Econ* 107:494–509, <https://doi.org/10.1016/j.ecolecon.2014.09.013>.