


Evaluation and Prognosis of Patients with Cirrhosis



Marion G. Peters, MD
John V. Carbone, MD, Endowed Chair
Professor of Medicine
Chief of Hepatology Research
University of California San Francisco

Survival Time From First Liver Decompensation to Death in HCV



- Death during study
 - 366/1037 HCV
 - 100/180 HIV/HCV
- Risk factors for death:
 - HIV
 - Baseline CTP
 - MELD >13
 - Age

Ascites



- **Diuretic-responsive ascites**
 - Sodium restriction
 - Spironolactone (75 mg-100 mg) and furosemide (20 mg-40 mg)
- **Refractory ascites**
 - Large volume paracentesis with 25% albumin (50 cc/L)
 - TIPS: higher orthotopic liver transplantation (OLT)-free survival, higher portosystemic encephalopathy (PSE)
- **Hyponatremia**
 - Fluid restriction, vasopressin 2 R antagonists
- **Spontaneous bacterial peritonitis complication**


Spontaneous Bacterial Peritonitis



- Most common type of bacterial infection in hospitalized patients with cirrhosis
- Clinical suspicion:
 - $< 50\%$: fever, abdominal pain or tenderness, and leukocytosis
 - unexplained encephalopathy, jaundice
 - worsening renal failure
- Diagnose: tap ascites: WCC > 500 , PMN > 250 cells/ μ L
 - Place ascites in blood culture bottles
- Start treatment immediately before culture results

Variceal Surveillance

All patients with cirrhosis require esophagogastroduodenoscopy



Hepatorenal Syndrome



- Results from vasodilatation and marked reduction in effective arterial blood volume leading to renal vasoconstriction
- Occurs in patients with refractory ascites and/or hyponatremia
- **Type 1 HRS:** rapidly progressive renal failure in 2 weeks
- **Type 2 HRS:** slowly progressive - median survival: ~ 6 months
 - Albumin, midodrine, and octreotide-induced vasoconstriction
 - OLT

Hepatic Encephalopathy



- Treatment aims to reduce production of ammonia from the colon through
 - nonabsorbable disaccharides
 - ✦ lactulose, lactitol, and lactose
 - nonabsorbable antibiotics
 - ✦ neomycin, rifaximin
- Protein restriction promotes protein degradation and, if maintained for long periods, worsens nutritional status and decreases muscle mass
 - No longer recommended

Summary: ESLD and HCV



- All HCV patients should be assessed for fibrosis stage prior to starting therapy
- Progression to decompensated cirrhosis can occur with interferon alfa-based therapies
- All patients with cirrhosis require monitoring for
 - Varices with EGD
 - HCC with imaging positive and negative alpha-fetoprotein
 - Ascites and infection (SBP prophylaxis)
- Consider OLT when first decompensation occurs

End



This presentation is brought to you by the International Antiviral Society-USA (IAS-USA)
in collaboration with Hepatitis Web Study & the Hepatitis C Online Course

Funded by a grant from the Centers for Disease Control and Prevention

Disclosure Information



Dr Peters has reported the following financial relationships with commercial firms:

- Consultant: Merck & Co, Inc, Theravance, and Roche
- Data safety monitoring board: Biotron
- Scientific advisor: Clinical Care Options
- Her spouse is employed by Genentech (Roche)

Outline




- Cirrhosis staging
- Clinical evidence of portal hypertension
 - Low platelets
 - Low white cell count
 - Splenomegaly
 - Spider nevi
- Management of cirrhosis
- How to recognize decompensated liver disease
- Management of decompensated liver disease

Distinguishing Compensated and Decompensated Cirrhosis



Natural History of End-Stage Liver Disease (ESLD)



- Alcohol
- Hepatitis C or B virus
- NASH
- Cholestatic
- Autoimmune

- Variceal hemorrhage
- Ascites
- Encephalopathy
- Jaundice

Decompensation

- Ascites (HRS, SBP)
- Encephalopathy
- Bleeding varices
- Coagulopathy

Natural History of ESLD



- Transition to decompensated cirrhosis: 5% to 7% of patients per year
- Best predictor of decompensation: hepatic venous pressure gradient (HVPG) > 10 mmHg
- HCC
 - can trigger decompensation
 - predictor of death in decompensated cirrhosis
- Tools for predicting disease severity and death in decompensated cirrhosis
 - Child-Turcotte-Pugh (CTP) score
 - Model for End-Stage Liver Disease (MELD) score

Child-Pugh-Turcotte Score

Points	1 (normal)	2	3
Hepatic encephalopathy	None	1-2	3-4
Ascites	None	slight	mod
Bilirubin (mg/dL)	<2	2-3	>3
Albumin (g/dL)	>3.5	2.8-3.5	<2.8
Prothrombin time	<4 secs ↑	4-6 secs	>6 secs
or International Normalized Ratio	<1.7	1.7-2.3	>2.3

A: 5-6

B: 7-9

C: > 9

Classification and Prognostic Systems for Patients with Cirrhosis



MELD: Model for ESLD

- | | | | | |
|----------------------|-----------|-----------|-----------|-----------|
| • Bilirubin (mg/dL) | 2 | 5 | 5 | 5 |
| • INR | 1.1 | 2.0 | 2.0 | 3.0 |
| • Creatinine (mg/dL) | 1.0 | 1.0 | 2.0 | 2.0 |
| • MELD | 10 | 20 | 27 | 31 |
- **Predicts 3-month mortality post TIPS/surgery**
 - **Used for allocation of liver donors**