

COGNITIVE ASSESSMENT TOOLKIT

A guide to detect cognitive impairment quickly and efficiently during the Medicare Annual Wellness Visit



alzheimer's  association®

800.272.3900 | alz.org®

TABLE OF CONTENTS

Overview	3
Medicare Annual Wellness Visit Algorithm for Assessment of Cognition	4
General Practitioner Assessment of Cognition (GPCOG)	5
Memory Impairment Screen (MIS)	7
Mini-Cog™	9
Short Form of the Informant Questionnaire on Cognitive Decline in the Elderly (Short IQCODE)	11
Eight-item Informant Interview to Differentiate Aging and Dementia (AD8)	14
Alzheimer’s Association Recommendations for Operationalizing the Detection of Cognitive Impairment During the Medical Annual Wellness Visit in a Primary Care Setting	17

OVERVIEW

The Alzheimer's Association – dedicated to fueling the advancement of early detection and diagnosis of dementia – has developed an easy-to-implement process to assess cognition during the Medicare Annual Wellness Visit. Developed by a group of clinical dementia experts, the recommended process outlined on page 4 allows you to efficiently identify patients with probable cognitive impairment while giving you the flexibility to choose a cognitive assessment tool that works best for you and your patients.


This Cognitive Assessment Toolkit contains:

- The Medicare Annual Wellness Visit Algorithm for Assessment of Cognition, incorporating patient history, clinician observations, and concerns expressed by the patient, family or caregiver
- Three validated patient assessment tools: the General Practitioner Assessment of Cognition (GPCOG), the Memory Impairment Screen (MIS) and the Mini-Cog™. All tools:
 - › Can be administered in 5 minutes or less
 - › Are equal to or superior to the Mini-Mental State Exam (MMSE) for detecting dementia
 - › Are easily administered by medical staff members who are not physicians
 - › Are relatively free from educational, language and/or cultural bias
- Three validated informant assessment of patient tools: the Short Form of the Informant Questionnaire on Cognitive Decline in the Elderly (Short IQCODE), the Eight-item Informant Interview to Differentiate Aging and Dementia (AD8) and the GPCOG
- The “Alzheimer's Association Recommendations for Operationalizing the Detection of Cognitive Impairment During the Medical Annual Wellness Visit in a Primary Care Setting,” as published in the journal *Alzheimer's and Dementia*.

For more information on the detection, diagnosis and treatment of Alzheimer's disease, as well as direct access to patient and caregiver resources, please visit our Health Care Professionals and Alzheimer's center at alz.org/hcps.

ALZHEIMER'S ASSOCIATION®

Medicare Annual Wellness Visit Algorithm for Assessment of Cognition



* No one tool is recognized as the best brief assessment to determine if a full dementia evaluation is needed. Some providers repeat patient assessment with an alternate tool (e.g., SLUMS, or MoCA) to confirm initial findings before referral or initiation of full dementia evaluation.

AD8 = Eight-item Informant Interview to Differentiate Aging and Dementia; **AWV** = Annual Wellness Visit; **GPCOG** = General Practitioner Assessment of Cognition; **HRA** = Health Risk Assessment; **MIS** = Memory Impairment Screen; **MMSE** = Mini Mental Status Exam; **MoCA** = Montreal Cognitive Assessment; **SLUMS** = St. Louis University Mental Status Exam; **Short IQCODE** = Short Informant Questionnaire on Cognitive Decline in the Elderly

Cordell CB, Borson S, Boustani M, Chodosh J, Reuben D, Verghese J, et al. Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual Wellness Visit in a primary care setting. *Alzheimers Dement*. 2013;9(2):141-150. Available at <http://download.journals.elsevierhealth.com/pdfs/journals/1552-5260/PIIS1552526012025010.pdf>.

Patient name: _____

Date: _____

GPCOG Screening Test

Step 1: Patient Examination

Unless specified, each question should only be asked once

Name and Address for subsequent recall test

1. *"I am going to give you a name and address. After I have said it, I want you to repeat it. Remember this name and address because I am going to ask you to tell it to me again in a few minutes: John Brown, 42 West Street, Kensington." (Allow a maximum of 4 attempts).*

Time Orientation

Correct Incorrect

2. *What is the date? (exact only)*

☐
☐

Clock Drawing – use blank page

3. *Please mark in all the numbers to indicate the hours of a clock (correct spacing required)*
4. *Please mark in hands to show 10 minutes past eleven o'clock (11.10)*

☐
☐
☐
☐

Information

5. *Can you tell me something that happened in the news recently? (Recently = in the last week. If a general answer is given, eg "war", "lot of rain", ask for details. Only specific answer scores).*

☐
☐

Recall

6. *What was the name and address I asked you to remember*

John

Brown

42

West (St)

Kensington

(To get a total score, add the number of items answered correctly)
Total correct (score out of 9)

/9

If patient scores 9, no significant cognitive impairment and further testing not necessary.

If patient scores 5-8, more information required. Proceed with Step 2, informant section.

If patient scores 0-4, cognitive impairment is indicated. Conduct standard investigations.

Informant Interview

Date: _____

Informant's name: _____

Informant's relationship to patient, i.e. informant is the patient's: _____

These six questions ask how the patient is compared to when s/he was well, say 5 – 10 years ago

Compared to a few years ago:

- | | Yes | No | Don't Know | N/A |
|--|--------------------------|--------------------------|--------------------------|--------------------------|
| ▪ Does the patient have more trouble remembering things that have happened recently than s/he used to? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| ▪ Does he or she have more trouble recalling conversations a few days later? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| ▪ When speaking, does the patient have more difficulty in finding the right word or tend to use the wrong words more often? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | |
| ▪ Is the patient less able to manage money and financial affairs (e.g. paying bills, budgeting)? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| ▪ Is the patient less able to manage his or her medication independently? | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| ▪ Does the patient need more assistance with transport (either private or public)?
(If the patient has difficulties due only to physical problems, e.g. bad leg, tick 'no') | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

(To get a total score, add the number of items answered 'no', 'don't know' or 'N/A')

Total score (out of 6)

If patient scores 0-3, cognitive impairment is indicated. Conduct standard investigations.

MEMORY IMPAIRMENT SCREEN (MIS)

Instructions for Administration

1. Show patient a sheet of paper with the 4 items to be recalled in 24-point or greater uppercase letters (on other side), and ask patient to read the items aloud.
2. Tell patient that each item belongs to a different category. Give a category cue and ask patient to indicate which of the words belongs in the stated category (eg, "Which one is the game?"). Allow up to 5 attempts. Failure to complete this task indicates possible cognitive impairment.
3. When patient identifies all 4 words, remove the sheet of paper. Tell patient that he or she will be asked to remember the words in a few minutes.
4. Engage patient in distractor activity for 2 to 3 minutes, such as counting to 20 and back, counting back from 100 by 7, spelling WORLD backwards.
5. FREE RECALL — 2 points per word: Ask patient to state as many of the 4 words he or she can recall. Allow at least 5 seconds per item for free recall. Continue to step 6 if no more words have been recalled for 10 seconds.
6. CUED RECALL — 1 point per word: Read the appropriate category cue for each word not recalled during free recall (eg, "What was the game?").

Word	Cue	Free recall (2 pts.)	Cued Recall (1 pts)
Checkers	Game		
Saucer	Dish		
Telegram	Message		
Red Cross	Organization		

Scoring

The maximum score for the MIS is 8.

- 5-8 No cognitive impairment
- ≤ 4 Possible cognitive impairment

Copyright © 1999 Albert Einstein College of Medicine. All rights reserved.

WORD LIST

CHECKERS

SAUCER

TELEGRAM

RED CROSS

Copyright © 1999 Albert Einstein College of Medicine. All rights reserved.

ID: _____ Date: _____

Step 1: Three Word Registration

Look directly at person and say, “Please listen carefully. I am going to say three words that I want you to repeat back to me now and try to remember. The words are [select a list of words from the versions below]. Please say them for me now.” If the person is unable to repeat the words after three attempts, move on to Step 2 (clock drawing).

The following and other word lists have been used in one or more clinical studies.¹⁻³ For repeated administrations, use of an alternative word list is recommended.

Version 1

 Banana
Sunrise
Chair

Version 2

 Leader
Season
Table

Version 3

 Village
Kitchen
Baby

Version 4

 River
Nation
Finger

Version 5

 Captain
Garden
Picture

Version 6

 Daughter
Heaven
Mountain

Step 2: Clock Drawing

Say: “Next, I want you to draw a clock for me. First, put in all of the numbers where they go.” When that is completed, say: “Now, set the hands to 10 past 11.”

Use preprinted circle (see next page) for this exercise. Repeat instructions as needed as this is not a memory test. Move to Step 3 if the clock is not complete within three minutes.

Step 3: Three Word Recall

Ask the person to recall the three words you stated in Step 1. Say: “What were the three words I asked you to remember?” Record the word list version number and the person’s answers below.


Word List Version: _____ Person’s Answers: _____

Scoring

Word Recall: _____ (0-3 points)	1 point for each word spontaneously recalled without cueing.
Clock Draw: _____ (0 or 2 points)	Normal clock = 2 points. A normal clock has all numbers placed in the correct sequence and approximately correct position (e.g., 12, 3, 6 and 9 are in anchor positions) with no missing or duplicate numbers. Hands are pointing to the 11 and 2 (11:10). Hand length is not scored. Inability or refusal to draw a clock (abnormal) = 0 points.
Total Score: _____ (0-5 points)	Total score = Word Recall score + Clock Draw score. A cut point of <3 on the Mini-Cog™ has been validated for dementia screening, but many individuals with clinically meaningful cognitive impairment will score higher. When greater sensitivity is desired, a cut point of <4 is recommended as it may indicate a need for further evaluation of cognitive status.

Clock Drawing

ID: _____ Date: _____



References

1. Borson S, Scanlan JM, Chen PJ et al. The Mini-Cog as a screen for dementia: Validation in a population-based sample. *J Am Geriatr Soc* 2003;51:1451–1454.
2. Borson S, Scanlan JM, Watanabe J et al. Improving identification of cognitive impairment in primary care. *Int J Geriatr Psychiatry* 2006;21: 349–355.
3. Lessig M, Scanlan J et al. Time that tells: Critical clock-drawing errors for dementia screening. *Int Psychogeriatr*. 2008 June; 20(3): 459–470.
4. Tsoi K, Chan J et al. Cognitive tests to detect dementia: A systematic review and meta-analysis. *JAMA Intern Med*. 2015; E1–E9.
5. McCarten J, Anderson P et al. Screening for cognitive impairment in an elderly veteran population: Acceptability and results using different versions of the Mini-Cog. *J Am Geriatr Soc* 2011; 59: 309–213.
6. McCarten J, Anderson P et al. Finding dementia in primary care: The results of a clinical demonstration project. *J Am Geriatr Soc* 2012; 60: 210–217.
7. Scanlan J & Borson S. The Mini-Cog: Receiver operating characteristics with the expert and naive raters. *Int J Geriatr Psychiatry* 2001; 16: 216–222.

**Short Form of the Informant Questionnaire on Cognitive
Decline in the Elderly (Short IQCODE)¹**

by A. F. Jorm

**Centre for Mental Health Research
The Australian National University
Canberra, Australia**

There is no copyright on the Short IQCODE. However, the author appreciates being kept informed of research projects which make use of it.

Note: As used in published studies, the IQCODE was preceded by questions to the informant on the subject's sociodemographic characteristics and physical health.

Now we want you to remember what your friend or relative was like 10 years ago and to compare it with what he/she is like now. 10 years ago was in 20__.* Below are situations where this person has to use his/her memory or intelligence and we want you to indicate whether this has improved, stayed the same or got worse in that situation over the past 10 years. Note the importance of comparing his/her present performance with 10 years ago. So if 10 years ago this person always forgot where he/she had left things, and he/she still does, then this would be considered "Hasn't changed much". Please indicate the changes you have observed by circling the appropriate answer.

Compared with 10 years ago how is this person at:

	1	2	3	4	5
1. Remembering things about family and friends e.g. occupations, birthdays, addresses	Much improved	A bit improved	Not much change	A bit worse	Much worse
2. Remembering things that have happened recently	Much improved	A bit improved	Not much change	A bit worse	Much worse
3. Recalling conversations a few days later	Much improved	A bit improved	Not much change	A bit worse	Much worse
4. Remembering his/her address and telephone number	Much improved	A bit improved	Not much change	A bit worse	Much worse
5. Remembering what day and month it is	Much improved	A bit improved	Not much change	A bit worse	Much worse
6. Remembering where things are usually kept	Much improved	A bit improved	Not much change	A bit worse	Much worse
7. Remembering where to find things which have been put in a different place from usual	Much improved	A bit improved	Not much change	A bit worse	Much worse
8. Knowing how to work familiar machines around the house	Much improved	A bit improved	Not much change	A bit worse	Much worse

9. Learning to use a new gadget or machine around the house	Much improved	A bit improved	Not much change	A bit worse	Much worse
10. Learning new things in general	Much improved	A bit improved	Not much change	A bit worse	Much worse
11. Following a story in a book or on TV	Much improved	A bit improved	Not much change	A bit worse	Much worse
12. Making decisions on everyday matters	Much improved	A bit improved	Not much change	A bit worse	Much worse
13. Handling money for shopping	Much improved	A bit improved	Not much change	A bit worse	Much worse
14. Handling financial matters e.g. the pension, dealing with the bank	Much improved	A bit improved	Not much change	A bit worse	Much worse
15. Handling other everyday arithmetic problems e.g. knowing how much food to buy, knowing how long between visits from family or friends	Much improved	A bit improved	Not much change	A bit worse	Much worse
16. Using his/her intelligence to understand what's going on and to reason things through	Much improved	A bit improved	Not much change	A bit worse	Much worse

*The original tool was published in 1994.

The Alzheimer's Association updated the year 19__ as published in the original tool to 20__ .

Tool Reference: Jorm AF. A short form of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE): development and cross-validation. Psychol Med 1994; 24: 145–153.

AD8 Dementia Screening Interview

Patient ID#: _____

CS ID#: _____

Date: _____

Remember, "Yes, a change" indicates that there has been a change in the last several years caused by cognitive (thinking and memory) problems.	YES, A change	NO, No change	N/A, Don't know
1. Problems with judgment (e.g., problems making decisions, bad financial decisions, problems with thinking)			
2. Less interest in hobbies/activities			
3. Repeats the same things over and over (questions, stories, or statements)			
4. Trouble learning how to use a tool, appliance, or gadget (e.g., VCR, computer, microwave, remote control)			
5. Forgets correct month or year			
6. Trouble handling complicated financial affairs (e.g., balancing checkbook, income taxes, paying bills)			
7. Trouble remembering appointments			
8. Daily problems with thinking and/or memory			
TOTAL AD8 SCORE			

Adapted from Galvin JE et al, The AD8, a brief informant interview to detect dementia, Neurology 2005;65:559-564

Copyright 2005. The AD8 is a copyrighted instrument of the Alzheimer's Disease Research Center, Washington University, St. Louis, Missouri.

All Rights Reserved.

The AD8 Administration and Scoring Guidelines

A spontaneous self-correction is allowed for all responses without counting as an error.

The questions are given to the respondent on a clipboard for self-administration or can be read aloud to the respondent either in person or over the phone. It is preferable to administer the AD8 to an informant, if available. If an informant is not available, the AD8 may be administered to the patient.

When administered to an informant, specifically ask the respondent to rate change in the patient.

When administered to the patient, specifically ask the patient to rate changes in his/her ability for each of the items, **without** attributing causality.

If read aloud to the respondent, it is important for the clinician to carefully read the phrase as worded and give emphasis to note changes due to cognitive problems (not physical problems). There should be a one second delay between individual items.

No timeframe for change is required.

The final score is a sum of the number items marked “Yes, A change”.

Interpretation of the AD8 (Adapted from Galvin JE et al, The AD8, a brief informant interview to detect dementia, *Neurology* 2005;65:559-564)

A screening test in itself is insufficient to diagnose a dementing disorder. The AD8 is, however, quite sensitive to detecting early cognitive changes associated many common dementing illness including Alzheimer disease, vascular dementia, Lewy body dementia and frontotemporal dementia.

Scores in the impaired range (see below) indicate a need for further assessment. Scores in the “normal” range suggest that a dementing disorder is unlikely, but a very early disease process cannot be ruled out. More advanced assessment may be warranted in cases where other objective evidence of impairment exists.


Based on clinical research findings from 995 individuals included in the development and validation samples, the following cut points are provided:

- 0 – 1: Normal cognition
- 2 or greater: Cognitive impairment is likely to be present

Administered to either the informant (preferable) or the patient, the AD8 has the following properties:

- Sensitivity > 84%
- Specificity > 80%
- Positive Predictive Value > 85%
- Negative Predictive Value > 70%
- Area under the Curve: 0.908; 95%CI: 0.888-0.925

Receiver Operator Characteristics (ROC) curve for AD8



Copyright 2005. The *Eight-item Informant Interview to Differentiate Aging and Dementia* is a copyrighted instrument of Washington University, St. Louis, Missouri. All Rights Reserved.

Permission Statement

Washington University grants permission to use and reproduce the *Eight-item Informant Interview to Differentiate Aging and Dementia* exactly as it appears in the PDF available here without modification or editing of any kind solely for end user use in investigating dementia in clinical care or research in clinical care or research (the "Purpose"). For the avoidance of doubt, the Purpose does not include the (i) sale, distribution or transfer of the *Eight-item Informant Interview to Differentiate Aging and Dementia* or copies thereof for any consideration or commercial value; (ii) the creation of any derivative works, including translations; and/or (iii) use of the *Eight-item Informant Interview to Differentiate Aging and Dementia* as a marketing tool for the sale of any drug. All copies of the AD8 shall include the following notice: "Reprinted with permission. Copyright 2005. The *Eight-item Informant Interview to Differentiate Aging and Dementia* is a copyrighted instrument of Washington University, St. Louis, Missouri. All Rights Reserved." Please contact morrisj@abraxas.wustl.edu for use of the *Eight-item Informant Interview to Differentiate Aging and Dementia* for any other intended purpose.

Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual Wellness Visit in a primary care setting

Cyndy B. Cordell^{a,*}, Soo Borson^{b,c}, Malaz Boustani^{d,e,f}, Joshua Chodosh^{g,h}, David Reuben^h, Joe Vergheseⁱ, William Thies^a, Leslie B. Fried^{j,k}; for the Medicare Detection of Cognitive Impairment Workgroup

^aAlzheimer's Association, Chicago, IL, USA

^bDepartment of Psychiatry and Behavioral Sciences, University of Washington School of Medicine, Seattle, WA, USA

^cMemory Disorders Clinic and Dementia Health Services, University of Washington School of Medicine, Seattle, WA, USA

^dIndiana University Center for Aging Research, Indianapolis, IN, USA

^eRegenstrief Institute, Inc., Indianapolis, IN, USA

^fDepartment of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA

^gVeterans Administration Greater Los Angeles Healthcare System, Los Angeles, CA, USA

^hDivision of Geriatrics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA

ⁱDepartment of Neurology, Albert Einstein College of Medicine, Bronx, NY, USA

^jAmerican Bar Association, Washington, DC, USA

^kAlzheimer's Association Medicare Advocacy Project, Washington, DC, USA

Abstract

The Patient Protection and Affordable Care Act added a new Medicare benefit, the Annual Wellness Visit (AWV), effective January 1, 2011. The AWV requires an assessment to detect cognitive impairment. The Centers for Medicare and Medicaid Services (CMS) elected not to recommend a specific assessment tool because there is no single, universally accepted screen that satisfies all needs in the detection of cognitive impairment. To provide primary care physicians with guidance on cognitive assessment during the AWV, and when referral or further testing is needed, the Alzheimer's Association convened a group of experts to develop recommendations. The resulting Alzheimer's Association Medicare Annual Wellness Visit Algorithm for Assessment of Cognition includes review of patient Health Risk Assessment (HRA) information, patient observation, unstructured queries during the AWV, and use of structured cognitive assessment tools for both patients and informants. Widespread implementation of this algorithm could be the first step in reducing the prevalence of missed or delayed dementia diagnosis, thus allowing for better healthcare management and more favorable outcomes for affected patients and their families and caregivers.

© 2013 The Alzheimer's Association. All rights reserved.

Keywords:

Annual Wellness Visit; AWV; Cognitive impairment; Assessment; Screen; Dementia; Alzheimer's disease; Medicare; Algorithm; Patient Protection and Affordable Care Act

1. Introduction

The Patient Protection and Affordable Care Act of 2010 added a new Medicare benefit, the Annual Wellness Visit

(AWV), effective January 1, 2011. The AWV includes routine measurements such as height, weight, and blood pressure; a review of medical and family history; an assessment to detect cognitive impairment; and establishment of a list of current medical providers, medications, and schedule for future preventive services. In addition, during the first AWV only, beneficiaries are to be screened for depression (if

*Corresponding author. Tel.: 312-335-5867. Fax: 866-699-1246.

E-mail address: cyndy.cordell@alz.org

not completed under a separate Medicare benefit) and for functional difficulties using nationally recognized appropriate screening questions or standardized questionnaires. Although the U.S. Preventive Services Task Force (USPSTF) in 2003 concluded that there was insufficient published evidence of better clinical outcomes as a result of routine screening for cognitive impairment in older adults, the Task Force recognized that the use of cognitive assessment tools can increase the detection of cognitive impairment [1]. As per the Centers for Medicare and Medicaid Services (CMS) regulation, the AWPV requires detection of cognitive impairment by "... assessment of an individual's cognitive function by direct observation, with due consideration of information obtained by way of patient report, concerns raised by family members, friends, caretakers, or others" [2]. During the public comment period, several organizations, including the Alzheimer's Association, noted that the use of a standardized tool for assessment of cognitive function should be part of the AWPV.

These comments are supported by a number of studies showing that cognitive impairment is unrecognized in 27%–81% of affected patients in primary care [3–7]. The use of a brief, structured cognitive assessment tool correctly classifies patients with dementia or mild cognitive impairment (MCI) more often than spontaneous detection by the patients' own primary care physicians (83% vs 59%, respectively) [8].

In response to concerns submitted during public comment, CMS elected not to recommend a specific tool for the final AWPV benefit because "There is no nationally recognized screening tool for the detection of cognitive impairments at the present time..." [9]. However, CMS recognizes that without clarification, the full intended benefits of the AWPV cognitive assessment may not be realized [10]. CMS is working with other governmental agencies (e.g., National Institutes on Aging) on recommendations for use of specific tools.

Understanding that, under the present regulation, each healthcare provider who conducts an AWPV would have to determine how best to "detect cognitive impairment," the Alzheimer's Association convened the Medicare Detection of Cognitive Impairment Workgroup to develop recommendations for operationalizing the cognitive assessment component in primary care settings. This workgroup was comprised of geographically dispersed USA experts with published works in the field of detecting cognitive impairment during primary care visits. The focus on primary care was deliberate, as most Medicare beneficiaries will receive their AWPV in this setting.

2. Guiding principles for recommendations

2.1. Consensus on general principles

Based on their expertise, the workgroup agreed on the following general principles to guide the development of recommendations for cognitive assessment:

- Detection of cognitive impairment is a stepwise, iterative process.
- Informal observation alone by a physician is not sufficient (i.e., observation without a specific cognitive evaluation).
- Detection of cognitive impairment can be enhanced by specifically asking about changes in memory, language, and the ability to complete routine tasks.
- Although no single tool is recognized as the "gold standard" for detection of cognitive impairment, an initial structured assessment should provide either a baseline for cognitive surveillance or a trigger for further evaluation.
- Clinical staff can offer valuable observations of cognitive and functional changes in patients who are seen over time.
- Counseling before and after cognitive assessment is an essential component of any cognitive evaluation.
- Informants (family member, caregiver, etc.) can provide valuable information about the presence of a change in cognition.

2.2. Principles specific to the AWPV

- The AWPV requires the completion of a Health Risk Assessment (HRA) by the patient either before or during the visit. The HRA should be reviewed for any reported signs and symptoms indicative of possible dementia.
- The AWPV will likely occur in a primary care setting. Tools for initial cognitive assessments should be brief (<5 min), appropriately validated, easily administered by non-physician clinical staff, and available free of charge for use in a clinical setting.
- If further evaluation is indicated based on the results of the AWPV, a more detailed evaluation of cognition should be scheduled for a follow-up visit in primary care or through referral to a specialist.

3. Review of available brief tools for use during the AWPV

3.1. Workgroup review process

Although there is no single cognition assessment tool that is considered to be the gold standard, there is a plethora of tools in the literature. A MEDLINE (PubMed) search conducted in October 2011, using the key words "screening or detection of dementia or cognitive impairment," yielded over 500 publications. To narrow the search to tools more applicable to the AWPV, the workgroup sought to determine whether the literature offered a consensus regarding brief cognitive assessment during time-limited primary care visits.

The workgroup focused on systematic evidence review (SER) studies published since 2000 resulting in four studies by Lorentz et al, Brodaty et al, Holsinger et al, and Milne et al [11–14]. Although each SER had a similar objective—to determine which tools were best for administration during

Table 1
Review articles of brief cognitive assessment tools—select inclusion and comparison criteria

	Lorentz et al, 2002 [11]	Brodaty et al, 2006 [12]	Holsinger et al, 2007 [13]	Milne et al, 2008 [14]	Ismail et al, 2010 [15]	Kansagara and Freeman, 2010 [16]
Inclusion criteria	<ul style="list-style-type: none"> • Admin ≤ 10 min • Performance characteristics evaluated in ≥ 1 community or clinical setting 	<ul style="list-style-type: none"> • Admin ≤ 5 min and simple • Validated in community or PC • Misclassification rate \leq MMSE • NPV \geq MMSE 	<ul style="list-style-type: none"> • Studied in patients ≥ 60 years • Criterion to diagnose dementia acceptable 	<ul style="list-style-type: none"> • Admin time suitable for PC in UK • Geriatric PC screens for cognitive change 	<ul style="list-style-type: none"> • Tools most frequently used in PC • Tools recommended or newly used in PC 	<ul style="list-style-type: none"> • Tools identified by the VA as alternatives to the MMSE
Comparison criteria	<ul style="list-style-type: none"> • Face validity, sensitivity, and specificity • Sociodemographic biases • Comparison with MMSE • Acceptability • Ease of use by nonspecialists 	<ul style="list-style-type: none"> • Study validity • Applicability to PC • Psychometric properties • Administration characteristics 	<ul style="list-style-type: none"> • Admin time • Study quality • Likelihood ratios • Domains tested • Utility in special situations 	<ul style="list-style-type: none"> • Practicality • Feasibility • Applicability • Psychometric properties 	<ul style="list-style-type: none"> • Summary of other studies and strength/weaknesses of tools • Newer tools that address weaknesses 	<ul style="list-style-type: none"> • Relevance of study to the VA setting • Admin time • Sensitivity • Specificity • Cost

Abbreviations: MMSE, Mini-Mental State Examination; NPV, negative predictive value; PC, primary care; UK, United Kingdom; VA, US Department of Veteran Affairs.

primary care visits—different comparison criteria to select the tools were applied (Table 1). Two other studies were also considered relevant to the development of the workgroup recommendations: Ismail et al [15] conducted a literature review designed to identify widely used and most promising newer brief cognitive tools being used in primary care and geriatrics, and an SER by Kansagara and Freeman [16] of six brief cognitive assessment tools that could serve as possible alternatives to the Mini-Mental State Examination (MMSE) for use by the U.S. Department of Veterans Affairs (VA). Neither study was designed to determine which brief tool is the “best,” but both provided evidence related to primary care use and performance characteristics of brief assessments of cognition (Table 1).

3.2. Workgroup review results

Of the five publications that focused specifically on identifying brief cognitive assessments most suitable or most used in primary care settings [11–15], all selected the Memory Impairment Screen (MIS), and four of these publications [11,12,14,15] also selected the General Practitioner Assessment of Cognition (GPCOG) and the Mini-Cog (Table 2).

The following attributes of the GPCOG, Mini-Cog, and the MIS contributed to their selection as most suited for routine use in primary care:

- Requires 5 minutes or less to administer.
- Is validated in a primary care or community setting.
- Is easily administered by medical staff members who are not physicians.
- Has good to excellent psychometric properties.
- Is relatively free from educational, language, and/or culture bias.
- Can be used by clinicians in a clinical setting without payment for copyrights.

Charging a fee for clinical use of brief cognitive assessment tool has become an issue because of increased enforcement of the MMSE copyright. First published in 1975 [17], the MMSE copyright is now held by Psychological Assessment Resources, Inc., which charges a fee for each use (for exact fees see www.parinc.com). The comparative SER within the VA [16] evaluated alternatives to the proprietary MMSE, including the GPCOG and the Mini-Cog, along with four other brief tools (Table 2). The Mini-Cog and MIS are copyrighted, but the owners, Soo Borson, MD, and Albert Einstein College of Medicine, respectively, allow free use by clinicians as clinical tools with distribution restrictions for other entities (e.g., commercial companies). The GPCOG has similar use rules.

3.3. Patient structured cognitive assessment tools recommended for AWW

In alignment with the workgroup's guiding principles and supported by data in the six selected SERs/reviews,

Table 2
Brief cognitive assessment tools evaluated in multiple review articles

Assessment Tool	Lorentz et al, 2002 [11]	Brodaty et al, 2006 [12]	Holsinger et al, 2007 [13]	Milne et al, 2008 [14]	Ismail et al, 2010 [15]	Kansagara and Freeman, 2010* [16]
7-Minute Screener	X	X	X	X		
AMT		X	X	X	X	
CAMCOG		X	Suited [†]			
CDT	X	X	Suited [‡]	X	X	
GPCOG	Most suited	Most suited	X	Most suited	Most suited	X
Mini-Cog	Most suited	Most suited	X	Most suited	Most suited	X
MIS	Most suited	Most suited	Suited [‡]	Most suited	Most suited	
MMSE	X	X	Suited [§]	X	X	
MoCA			Suited [†]		X	X
RUDAS		X			X	
SAS-SI	X	X	X			
SBT (BOMC, 6-CIT)	X	X	X	X		X
SPMSQ	X			X		
STMS	X	X	X			X
T&C	X	X				

Abbreviations: 6-CIT, 6-Item Cognitive Impairment Test; AMT, Abbreviated Mental Test; BOMC, 6-item Blessed Orientation-Memory-Concentration Test; CAMCOG, Cambridge Cognitive Examination; CDT, Clock Drawing Test; GPCOG, General Practitioner Assessment of Cognition; MIS, Memory Impairment Screen; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment; RUDAS, Rowland Universal Dementia Assessment; SAS-SI, Short and Sweet Screening Instrument; SBT, Short Blessed Test; SLUMS, St Louis Mental Status; SPMSQ, Short Portable Mental Status Questionnaire; STMS, Short Test of Mental Status; T&C, Time and Change Test.

X = assessment reviewed, but not identified as most suited for general use in primary care.

Suited = tool appropriate for the following clinical issue: [†] available time is not limited; [‡] available time is limited; and [§] cognitive impairment is at least moderate. Most suited = tool identified as most suited for routine use in primary care.

*Kansagara and Freeman evaluated six tools, including the SLUMS, which was not evaluated in any other review.

the GPCOG, Mini-Cog, and MIS are brief structured tools that are suitable for assessment of cognitive function during the AWV. Each tool has unique benefits. The GPCOG has patient and informant components that can be used alone or together to increase specificity and sensitivity [18]. The Mini-Cog has been validated in population-based studies and in community-dwelling older adults heterogeneous with respect to language, culture, and education [19–22]. The MIS is a verbally administered word-recall task that tests encoding as well as retrieval [23], and is an option for patients who have motor impairments that prevent use of paper and pencil.

3.4. Structured cognitive assessment tools for use with informants

Cognitive assessment combined with informant-reported data improves the accuracy of assessment [24–27]. If an informant is present during the AWV, use of a structured informant tool is recommended. Similar to cognitive assessment tools for use with patients, there is no single “gold standard” informant tool; however, relatively few brief informant tools have been validated in community and/or primary care settings. Brief tools appropriately validated include the Short IQCODE [25], the AD8 [28], which can be administered in-person or by telephone, and the aforementioned GPCOG [18], which has both patient and informant components.

4. Recommended algorithm for detection of cognitive impairment during the AWV


4.1. Incorporating assessment of cognition during the AWV

The Alzheimer's Association Medicare Annual Wellness Visit Algorithm for Assessment of Cognition for consistency (Figure 1) illustrates a stepwise process. The process is intended to detect patients with a high likelihood of having dementia. The AWV algorithm includes both structured assessments discussed previously and other less structured patient- and informant-based evaluations. By assessing and documenting cognitive status on an annual basis during the AWV, clinicians can more easily determine gradual cognitive decline over time in an individual patient—a key criterion for diagnosing dementia due to Alzheimer's disease and other progressive conditions affecting cognition.

For patients with a previous diagnosis of MCI or dementia, this should be documented and included in their AWV list of health risk factors. Annual unstructured and structured cognitive assessments could be used to monitor significant changes in cognition and potentially lead to a new diagnosis of dementia for those with MCI or new care recommendations for those with dementia.

4.2. Detection of cognitive impairment during the AWV—initial HRA review, conversations, and observations

The first step in detection of cognitive impairment during the AWV (Fig. 1, Step A), involves a conversation between



AWV = Annual Wellness Visit; GPCOG = General Practitioner Assessment of Cognition; HRA = Health Risk Assessment; MIS = Memory Impairment Screen; MMSE = Mini Mental Status Exam; MoCA = Montreal Cognitive Assessment; SLUMS = St. Louis University Mental Status Exam; Short IQCODE = short Informant Questionnaire on Cognitive Decline in the Elderly

Fig. 1. Alzheimer's Association Medicare Annual Wellness Visit Algorithm for Assessment of Cognition.

a clinician and the patient and, if present, any family member or other person who can provide collateral information. This introduces the purpose and content of the AWV, which includes: a review of the HRA; observations by clinicians (medical and associated staff); acknowledgment of any self-reported or informant-reported concerns; and conversational queries about cognition directed toward the patient and others present. If any concerns are noted, or if an informant is not present to provide confirmatory information, further evaluation of cognition with a structured tool should be performed.

Patient completion of an HRA is a required element of the AWV and can be accomplished with the help of a family member or other knowledgeable informants, including a professional caregiver. Published CMS guidance offers healthcare professionals flexibility as to the specific format, questions, and delivery methods that can be used for an AWV HRA [29]. The following questions may be suitable for the AWV HRA and have been tested and evaluated in the general popu-

lation through the Behavioral Risk Factor Surveillance System or presented as HRA example questions:

1. During the past 12 months, have you experienced confusion or memory loss that is happening more often or is getting worse [30]?
2. During the past 7 days, did you need help with others to perform everyday activities such as eating, getting dressed, grooming, bathing, walking, or using the toilet [29]?
3. During the past 7 days, did you need help from others to take care of things such as laundry and housekeeping, banking, shopping, using the telephone, food preparation, transportation, or taking your own medications [29]?

A noted deficit in activities of daily living (ADLs) (e.g., eating and dressing) or instrumental activities of daily living (IADLs) (e.g., shopping and cooking) that cannot be

attributed to physical limitations should prompt concern, as there is a strong correlation between decline in function and decline in cognitive status across the full spectrum of dementia [31]. In addition to clinically observed concerns, any patient- or informant-reported concerns should trigger further evaluation [13]. Positive responses to conversational queries, such as “Have you noticed any change in your memory or ability to complete routine tasks, such as paying bills or preparing a meal?” should be followed up with a structured assessment of cognition.

Upon realizing the time constraints of a typical primary care visit, if no cognitive concerns surface during the initial evaluation and this information is corroborated by an informant, the clinician may elect not to perform a structured cognitive assessment and assume that the patient is not currently demented. This approach is supported by studies in populations with low rates of dementia that suggest the absence of memory difficulties reported by informants and patients reduces the likelihood that dementia is present [32,33].

4.3. Structured cognitive assessment tools for use with patients and informants during the AWV

The second step in detection of cognitive impairment during the AWV (Figure 1, Step B) requires cognitive assessment using a structured tool. Based on synthesis of data from the six review articles previously discussed, patient tools suitable for the initial structured assessment are the GPCOG, Mini-Cog, and MIS.

Recognizing that there is no single optimal tool to detect cognitive impairment for all patient populations and settings, clinicians may select other brief tools to use in their clinical practice, such as those listed in Table 3. The 15 brief tools listed were evaluated in multiple review articles (passed through at least two review search criteria for tools possibly suited for primary care) or are used in the VA. Tools listed in Table 3 are subject to the inclusion/exclusion criteria of each review and do not represent the entire listing of the >100 brief cognitive assessment tools that may be suitable for primary care practices.

If an informant is present, defined as someone who can attest to a patient's change in memory, language, or function over time, it is suitable to use the AD8, the informant component of the GPCOG, or the Short IQCODE, during the AWV.

4.4. Primary care workflow considerations

According to the algorithm, any patient who does not have an informant present should be assessed with a structured tool. For such patients (and for practices that implement structured assessments during all AWVs), completion of this structured assessment can be administered by trained medical staff as the first step for cognitive impairment detection. This could improve office efficiency. To increase acceptance of a structured assessment, the reason provided to

the patient can be normalized with a statement such as, “This is something I do for all of my older patients as part of their annual visit.” When the initial assessment prompts further evaluation, explanation of results should be deferred until a more comprehensive evaluation has been completed. “There are many reasons for not getting every answer correct. More evaluation will help us determine that,” is an example statement that may encourage patients to pursue further testing.

5. Full dementia evaluation

Patients with assessments that indicate cognitive impairment during the AWV should be further evaluated to determine appropriate diagnosis (e.g., MCI, Alzheimer's disease) or to identify other causes. As reflected in the algorithm (Figure 1, Step C), initiation of a full dementia evaluation is outside the scope of the AWV, but can occur in a separate visit either on the same day, during a newly scheduled visit, or through referral to a specialist. Specialists who have expertise in diagnosing dementia include geriatricians, geriatric psychiatrists, neurologists, and neuropsychologists. The two-visit approach has been cited as a time-effective process to evaluate suspected dementia in primary care [34] and is consistent with the two-step approach widely used in epidemiologic research on dementia. Regardless of the timing and setting, clinicians are encouraged to counsel patients to include an informant in the diagnostic process.

Components of a full dementia evaluation can vary depending on the presentation and include tests to rule in or out the various causes of cognitive impairment and establish its severity. Diagnostic evaluations include a complete medical history; assessment of multiple cognitive domains, including episodic memory, executive function, attention, language, and visuospatial skills; neurologic exam (gait, motor function, reflexes); ADL and IADL functioning; assessment for depression; and review for medications that may adversely affect cognition. Standard laboratory tests include thyroid-stimulating hormone (TSH), complete blood count (CBC), serum B₁₂, folate, complete metabolic panel, and, if the patient is at risk, testing for sexually transmitted diseases (human immunodeficiency virus, syphilis). Structural brain imaging, including magnetic resonance imaging (MRI) or computed tomography (CT), is a supplemental aid in the differential diagnosis of dementia, especially if neurologic physical exam findings are noted. An MRI or CT can be especially informative in the following cases: dementia that is of recent onset and is rapidly progressing; younger onset dementia (<65 years of age); history of head trauma; or neurologic symptoms suggesting focal disease.

6. Discussion

Unfortunately, up to 81% of patients who meet the criteria for dementia have never received a documented diagnosis

Table 3

Key advantages and limitations of brief cognitive assessment tools evaluated in multiple reviews and/or for use in the VA

Assessment*	Time (~min)	Advantages	Limitations
7-Minute Screener [48]	7–12	<ul style="list-style-type: none"> • Little or no education bias • Validated in primary care 	<ul style="list-style-type: none"> • Difficult to administer • Complex logarithmic scoring
AMT [49]	5–7	<ul style="list-style-type: none"> • Easy to administer • Verbal memory test (no writing/drawing) 	<ul style="list-style-type: none"> • Education/language/culture bias • Limited use in US (mostly used in Europe) • Does not test executive function or visuospatial skills
CAMCOG [50]	20	<ul style="list-style-type: none"> • Tests many separate domains (7) 	<ul style="list-style-type: none"> • Difficult to administer • Long administration time
CDT [51]	≤1	<ul style="list-style-type: none"> • Very brief administration time • Minimal education bias 	<ul style="list-style-type: none"> • Lacks standards for administration and scoring
GPCOG [†] [18]			
Patient	2–5	<ul style="list-style-type: none"> • Developed for and validated in primary care 	<ul style="list-style-type: none"> • Patient component scoring has an indeterminate range that requires an informant score to assess as pass or fail
Informant	1–3	<ul style="list-style-type: none"> • Informant component useful when initial complaint is informant-based • Little or no education bias • Multiple languages accessible at www.gpcog.com.au 	<ul style="list-style-type: none"> • Informant component alone has low specificity • Lacks data on any language/culture biases
Mini-Cog [†] [8, 19]	2–4	<ul style="list-style-type: none"> • Developed for and validated in primary care and multiple languages/cultures • Little or no education/language/race bias • Short administration time 	<ul style="list-style-type: none"> • Use of different word lists may affect failure rates • Some study results based on longer tests with the Mini-Cog elements reviewed independently
MIS [23,52]	4	<ul style="list-style-type: none"> • Verbal memory test (no writing/drawing) • Little or no education bias 	<ul style="list-style-type: none"> • Does not test executive function or visuospatial skills
MMSE [17]	7–10	<ul style="list-style-type: none"> • Most widely used and studied worldwide • Often used as reference for comparative evaluations of other assessments • Required for some drug insurance reimbursements 	<ul style="list-style-type: none"> • Education/age/language/culture bias • Ceiling effect (highly educated impaired subjects pass) • Proprietary—unless used from memory, test needs to be purchased at www.parinc.com • Best performance for at least moderate cognitive impairment
MoCA [†] [53]	10–15	<ul style="list-style-type: none"> • Designed to test for mild cognitive impairment • Multiple languages accessible at www.mocatest.org • Tests many separate domains (7) 	<ul style="list-style-type: none"> • Lacks studies in general practice settings • Education bias (≤12 years) • Limited use and evidence due to published data relatively new (2005) • Admin time ≥10 min
RUDAS [54]	10	<ul style="list-style-type: none"> • Designed for multicultural populations • Little or no education/language bias 	<ul style="list-style-type: none"> • Validated in Australian community • Limited use and evidence due to published data relatively new (2004)
SAS-SI [55]	10	<ul style="list-style-type: none"> • Detected dementia better than neuropsychologic testing in a community population 	<ul style="list-style-type: none"> • Does not test memory • Lacks data on any education/language/culture biases
SBT (BOMC [†] and 6-CIT) [56,57]	4–6	<ul style="list-style-type: none"> • Verbal test (no writing/drawing) 	<ul style="list-style-type: none"> • Education/language/cultural/race bias • Scoring can be cumbersome • Does not test executive function
SLUMS [†] [58]	7	<ul style="list-style-type: none"> • No education bias • Tests many separate domains (7) • Available at: http://aging.slu.edu/pdfsurveys/mentalstatus.pdf 	<ul style="list-style-type: none"> • Limited use and evidence due to published data relatively new (2006) • Studied in VA geriatric clinic (predominantly white males)
SPMSQ [59]	3–4	<ul style="list-style-type: none"> • Verbal test (no writing/drawing) 	<ul style="list-style-type: none"> • Scoring can be cumbersome • Does not test short-term memory
STMS [†] [60]	5	<ul style="list-style-type: none"> • Validated in primary care • Tests many separate domains (7) 	<ul style="list-style-type: none"> • Education/language/race bias • Studied in relatively educated subjects, may not be applicable to general population
T&C [61]	≤1	<ul style="list-style-type: none"> • Very brief administration time • Little or no education bias 	<ul style="list-style-type: none"> • Strong language/cultural bias

Abbreviations: 6-CIT, 6-Item Cognitive Impairment Test; AMT, Abbreviated Mental Test; BOMC, 6-item Blessed Orientation-Memory-Concentration Test; CAMCOG, Cambridge Cognitive Examination; CDT, Clock Drawing Test; GPCOG, General Practitioner Assessment of Cognition; MIS, Memory Impairment Screen; MMSE, Mini-Mental State Examination; MoCA, Montreal Cognitive Assessment; RUDAS, Rowland Universal Dementia Assessment; SAS-SI, Short and Sweet Screening Instrument; SBT, Short Blessed Test; SLUMS, St Louis University Mental Status; SPMSQ, Short Portable Mental Status Questionnaire; STMS, Short Test of Mental Status; T&C, Time and Change Test.

*References provide descriptions of assessments.

[†]Brief tools used in the VA healthcare system reviewed by Kansagara and Freeman.

[35]. Delayed or missed diagnosis deprives affected individuals of available treatments, care plans, and services that can improve their symptoms and help maintain independence. Studies show that interventions tailored to patients with dementia can improve quality of care, reduce unfavorable dementia-related behaviors, increase access to community services for both the patient and their caregivers, and result in less caregiver stress and depression [36–42]. Early diagnosis of dementia also provides families and patients an opportunity to plan for the future while the affected individual is still able to participate in the decision-making processes.

Early detection and medical record documentation may improve medical care. The medical record could inform all clinicians, including those who may be managing comorbidities on a sporadic basis, that treatment and care should be adjusted to accommodate cognitive impairment. According to a 2004 Medicare beneficiary survey, among patients with dementia, 26% had coronary heart disease, 23% had diabetes, and 13% had cancer [43].

It is important to note that the unstructured and structured cognitive assessments being recommended for the AWV are only the first steps in diagnosing dementia, and cognitive assessment is best as an iterative process. For example, clinicians concerned with HRA information about decline in function may proceed directly to a structured assessment or continue to query the patient for additional information; a self-reported memory concern coupled with a failed structured cognitive assessment should always result in a full dementia evaluation.

Not all who are referred for further assessment will ultimately receive a dementia diagnosis. In a USA primary care population aged ≥ 65 years ($N = 3340$), 13% failed a brief screen for cognitive impairment and approximately half ($n = 227$) agreed to be further evaluated for dementia [7]. Among the 107 patients ultimately diagnosed with dementia, 81% were newly diagnosed based on the absence of any medical record of dementia, thus facilitating appropriate medical and psychosocial interventions [7].

Despite the many advantages of early dementia diagnosis, several barriers to diagnosis still exist. These include physician concerns of the time burden resulting from testing and counseling [35] and stigma concerns among physicians, patients, and caregivers [35,44,45]. Despite these barriers, successful widespread implementation of a brief cognitive assessment has been reported. McCarten et al [22] evaluated the Mini-Cog for routine cognitive assessment of veterans presenting for primary care. Of the 8342 veterans approached, >96% agreed to be assessed and those that failed the brief assessment exhibited no serious reactions upon disclosure of test results.

The AWV provides an unprecedented opportunity to overcome current barriers and initiate discussions about cognitive function among the growing population most at risk

for Alzheimer's disease. Detection of cognitive impairment during the AWV is further supported by previously published quality indicators that state all vulnerable elders (defined as persons ≥ 65 years who are at risk for death or functional decline) should be evaluated annually for cognitive and functional status [46].

There are limitations to these recommendations. They are based on assessment of recommendations from review articles and on expert opinion, not on a new, comprehensive review of original research to define the optimal approach to detection of cognitive impairment or review of emerging technologies that could assist in testing (e.g., use of online or electronic tablet applications). Further complicating SERs of brief cognitive assessment tools is that sensitivity and specificity will vary depending on the dementia prevalence of the study population, the tool(s) used, and the cut score selected for each tool. Brodaty et al [12] recognized that published research concerning cognitive impairment screening tools is uneven in quantity and quality. The literature also is lacking in comparative validity of brief cognitive assessment tools in low-education or illiterate populations.

The Alzheimer's Association Medicare Annual Wellness Visit Algorithm for Assessment of Cognition is based on current validated tools and commonly used rule-out assessments. The use of biomarkers (e.g., CSF tau and beta amyloid proteins, amyloid tracer positron emission tomography scans) was not considered as these measures are not currently approved or widely available for clinical use.

In 2011, greater than two million Medicare beneficiaries received their AWV preventive service [47]. There are no data available as to what methods were used to detect cognitive impairment or how many beneficiaries were assessed as having cognitive impairment. For future AWVs, the Alzheimer's Association Medicare Annual Wellness Visit Algorithm for Assessment of Cognition provides guidance to primary care practices on a process to operationalize this required AWV element. With widespread implementation of the algorithm, the AWV could be the first step in reducing the prevalence of missed or delayed dementia diagnoses, thus allowing for better healthcare management and more favorable outcomes for affected patients and their families and caregivers.

7. Author Disclosures

Soo Borson is the developer of the Mini-Cog and is the owner of its copyrights.

Over the past 5 years, Malaz Boustani has received research support for investigator-initiated projects from Forest Pharmaceutical and Novartis; honoraria from Novartis and Pfizer, Inc.; and research support for investigator-initiated projects from the NIH and AHRQ. Dr Boustani was a member of the US Preventive Services Task Force that published

the systematic evidence review, *Dementia Screening*, for the AHRQ in 2003.

RESEARCH IN CONTEXT

1. Systematic review: Our research included comparing five systematic evidence reviews (SER) of brief dementia screening tools published since 2000 and a 2010 literature review of newer brief assessments of cognition. Our research focused on determining if there was a consensus among the published SERs as to which tool is most suited for primary care and if there were any common results across the publications.
2. Interpretation: Our research concluded there is a consensus in the literature concerning suitable tools for screening for dementia in primary care. We also reaffirmed that many validated tools are available, and that screening for dementia should not be solely based on a tool, but should be a stepwise process to include other assessments.
3. Future directions: Further validation of existing and emerging screening tools (e.g., iPad applications, gait monitoring) may result in newer tools being recognized more suitable and practical for primary care settings.

References

- [1] Boustani M, Peterson B, Harris R, Lux L, Krasnov C, Sutton S, et al. Screening for dementia. Rockville, MD: Agency for Healthcare Research and Quality. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK42773/>; 2003. Accessed September 3, 2011.
- [2] Anonymous. Patient Protection and Affordable Care Act of 2010, 42 CFR. §410.15(a). 2010. Available at: <http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?c=ecfr&sid=6b50669da0f96db4eea346533db23747&rgn=div8&view=text&node=42:2.0.1.2.10.2.35.4&idno=42>.
- [3] Chodosh J, Petitti DB, Elliott M, Hays RD, Crooks VC, Reuben DB, et al. Physician recognition of cognitive impairment: evaluating the need for improvement. *J Am Geriatr Soc* 2004; 52:1051–9.
- [4] Camicioli R, Willert P, Lear J, Grossmann S, Kaye J, Butterfield P. Dementia in rural primary care practices in Lake County, Oregon. *J Geriatr Psychiatry Neurol* 2000;13:87–92.
- [5] Callahan CM, Hendrie HC, Tierney WM. Documentation and evaluation of cognitive impairment in elderly primary care patients. *Ann Intern Med* 1995;122:422–9.
- [6] Valcour VG, Masaki KH, Curb JD, Blanchette PL. The detection of dementia in the primary care setting. *Arch Intern Med* 2000; 160:2964–8.
- [7] Boustani M, Callahan CM, Unverzagt FW, Austrom MG, Perkins AJ, Fultz BA, et al. Implementing a screening and diagnosis program for dementia in primary care. *J Gen Intern Med* 2005; 20:572–7.
- [8] Borson S, Scanlan JM, Watanabe J, Tu S-P, Lessig M. Improving identification of cognitive impairment in primary care. *Int J Geriatr Psychiatry* 2006;21:349–55.
- [9] Anonymous. Medicare coverage of Annual Wellness Visit providing a personalized prevention plan. *Fed Regist* 2010;75:73401.
- [10] US Department of Health and Human Services. Advisory Council on Alzheimer's research, care, and services: opportunities and gaps. 2011. Available at: <http://aspe.hhs.gov/daltcp/napa/092711/Mtg1-Slides3.pdf>. Accessed October 14, 2011.
- [11] Lorentz WJ, Scanlan JM, Borson S. Brief screening tests for dementia. *Can J Psychiatry* 2002;47:723–33.
- [12] Brodaty H, Low L-F, Gibson L, Burns K. What is the best dementia screening instrument for general practitioners to use? *Am J Geriatr Psychiatry* 2006;14:391–400.
- [13] Holsinger T, Deveau J, Boustani M, Williams JW Jr. Does this patient have dementia? *JAMA* 2007;297:2391–404.
- [14] Milne A, Culverwell A, Guss R, Tuppen J, Whelton R. Screening for dementia in primary care: a review of the use, efficacy and quality of measures. *Int Psychogeriatr* 2008;20:911–26.
- [15] Ismail Z, Rajji TK, Shulman KI. Brief cognitive screening instruments: an update. *Int J Geriatr Psychiatry* 2010;25:111–20.
- [16] Kansagara D, Freeman M. A systematic evidence review of the signs and symptoms of dementia and brief cognitive tests. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21155200>. Accessed June 7, 2011.
- [17] Folstein MF, Folstein SE, McHugh PR. "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–98.
- [18] Brodaty H, Pond D, Kemp NM, Luscombe G, Harding L, Berman K, et al. The GPCOG: a new screening test for dementia designed for general practice. *J Am Geriatr Soc* 2002;50:530–4.
- [19] Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The Mini-Cog: a cognitive "vital signs" measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry* 2000;15:1021–7.
- [20] Borson S, Scanlan JM, Watanabe J, Tu S-P, Lessig M. Simplifying detection of cognitive impairment: comparison of the Mini-Cog and Mini-Mental State Examination in a multiethnic sample. *J Am Geriatr Soc* 2005;53:871–4.
- [21] Borson S, Scanlan JM, Chen P, Ganguli M. The Mini-Cog as a screen for dementia: validation in a population-based sample. *J Am Geriatr Soc* 2003;51:1451–4.
- [22] McCarten JR, Anderson P, Kuskowski MA, McPherson SE, Borson S. Screening for cognitive impairment in an elderly veteran population: acceptability and results using different versions of the Mini-Cog. *J Am Geriatr Soc* 2011;59:309–13.
- [23] Buschke H, Kuslansky G, Katz M, Stewart WF, Sliwinski MJ, Eckholdt HM, et al. Screening for dementia with the memory impairment screen. *Neurology* 1999;52:231–8.
- [24] Mackinnon A, Mulligan R. Combining cognitive testing and informant report to increase accuracy in screening for dementia. *Am J Psychiatry* 1998;155:1529–35.
- [25] Jorm AF. A short form of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE): development and cross-validation. *Psychol Med* 1994;24:145–53.
- [26] Ayalon L. The IQCODE versus a single-item informant measure to discriminate between cognitively intact individuals and individuals with dementia or cognitive impairment. *J Geriatr Psychiatry Neurol* 2011;24:168–73.
- [27] Galvin JE, Roe CM, Morris JC. Evaluation of cognitive impairment in older adults: combining brief informant and performance measures. *Arch Neurol* 2007;64:718–24.
- [28] Galvin JE, Roe CM, Xiong C, Morris JC. Validity and reliability of the AD8 informant interview in dementia. *Neurology* 2006; 67:1942–8.
- [29] Goetzel R, Staley P, Ogden L, Strange P, Fox J, Spangler J, et al. A framework for patient-centered health risk assessments—providing health promotion and disease prevention services to Medicare beneficiaries. Atlanta, GA: US Department of Health and Human Services,

- Centers for Disease Control and Prevention. Available at: <http://www.cdc.gov/policy/opth/hra/>; 2011.
- [30] Anonymous. Behavioral Risk Factor Surveillance System Survey Questionnaire. Bethesda, MD: Centers for Disease Control and Prevention. Available at: <http://www.cdc.gov/brfss/questionnaires/pdf-ques/2011brfss.pdf>; 2011. Accessed January 10, 2012.
 - [31] Njegovan V, Hing MM, Mitchell SL, Molnar FJ. The hierarchy of functional loss associated with cognitive decline in older persons. *J Gerontol A Biol Sci Med Sci* 2001;56:M638–43.
 - [32] Carr DB, Gray S, Baty J, Morris JC. The value of informant versus individual's complaints of memory impairment in early dementia. *Neurology* 2000;55:1724–6.
 - [33] Tobiansky R, Blizard R, Livingston G, Mann A. The Gospel Oak Study stage IV: the clinical relevance of subjective memory impairment in older people. *Psychol Med* 1995;25:779–86.
 - [34] Simmons BB, Hartmann B, DeJoseph D. Evaluation of suspected dementia. *Am Fam Physician* 2011;84:895–902.
 - [35] Bradford A, Kunik ME, Schulz P, Williams SP, Singh H. Missed and delayed diagnosis of dementia in primary care: prevalence and contributing factors. *Alzheimer Dis Assoc Disord* 2009;23:306–14.
 - [36] Bass DM, Clark PA, Looman WJ, McCarthy CA, Eckert S. The Cleveland Alzheimer's managed care demonstration: outcomes after 12 months of implementation. *Gerontologist* 2003;43:73–85.
 - [37] Callahan CM, Boustani MA, Unverzagt FW, Austrom MG, Damush TM, Perkins AJ, et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA* 2006;295:2148–57.
 - [38] Fortinsky RH, Unson CG, Garcia RI. Helping family caregivers by linking primary care physicians with community-based dementia care services: the Alzheimer's Service Coordination Program. *Dementia* 2002;1:227–40.
 - [39] Reuben DB, Roth CP, Frank JC, Hirsch SH, Katz D, McCreath H, et al. Assessing care of vulnerable elders—Alzheimer's disease: a pilot study of a practice redesign intervention to improve the quality of dementia care. *J Am Geriatr Soc* 2010;58:324–9.
 - [40] Vickrey BG, Mittman BS, Connor KI, Pearson ML, Della Penna RD, Ganiats TG, et al. The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial. *Ann Intern Med* 2006;145:713–26.
 - [41] Olazarán J, Reisberg B, Clare L, Cruz I, Peña-Casanova J, Del Ser T, et al. Nonpharmacological therapies in Alzheimer's disease: a systematic review of efficacy. *Dement Geriatr Cogn Disord* 2010;30:161–78.
 - [42] Auclair U, Epstein C, Mittelman M. Couples counseling in Alzheimer's disease: additional clinical findings from a novel intervention study. *Clin Gerontol* 2009;32:130–46.
 - [43] Thies W, Bleiler L. 2011 Alzheimer's disease facts and figures. *Alzheimers Dement* 2011;7:208–44.
 - [44] Justiss MD, Boustani M, Fox C, Katona C, Perkins AJ, Healey PJ, et al. Patients' attitudes of dementia screening across the Atlantic. *Int J Geriatr Psychiatry* 2009;24:632–7.
 - [45] Boustani MA, Justiss MD, Frame A, Austrom MG, Perkins AJ, Cai X, et al. Caregiver and noncaregiver attitudes toward dementia screening. *J Am Geriatr Soc* 2011;59:681–6.
 - [46] Feil DG, MacLean C, Sultzer D. Quality indicators for the care of dementia in vulnerable elders. *J Am Geriatr Soc* 2007;55(Suppl 2):S293–301.
 - [47] Anonymous. Preventive New Media. Centers for Medicare & Medicaid Services. Available at: http://www.cms.gov/NewMedia/02_preventive.asp; 2011. Accessed January 10, 2012.
 - [48] Solomon PR, Pendlebury WW. Recognition of Alzheimer's disease: the 7 Minute Screen. *Fam Med* 1998;30:265–71.
 - [49] Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. *Age Ageing* 1972;1:233–8.
 - [50] Roth M. CAMDEX: the Cambridge examination for mental disorders of the elderly. Cambridge: Cambridge University Press; 1988.
 - [51] Shulman KI. Clock-drawing: is it the ideal cognitive screening test? *Int J Geriatr Psychiatry* 2000;15:548–61.
 - [52] Kuslansky G, Buschke H, Katz M, Sliwinski M, Lipton RB. Screening for Alzheimer's disease: the memory impairment screen versus the conventional three-word memory test. *J Am Geriatr Soc* 2002;50:1086–91.
 - [53] Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005;53:695–9.
 - [54] Storey JE, Rowland JTI, Basic D, Conforti DA, Dickson HG. The Rowland Universal Dementia Assessment Scale (RUDAS): a multicultural cognitive assessment scale. *Int Psychogeriatr* 2004;16:13–31.
 - [55] Belle SH, Mendelsohn AB, Seaberg EC, Ratcliff G. A brief cognitive screening battery for dementia in the community. *Neuroepidemiology* 2000;19:43–50.
 - [56] Katzman R, Brown T, Fuld P, Peck A, Schechter R, Schimmel H. Validation of a short Orientation-Memory-Concentration Test of cognitive impairment. *Am J Psychiatry* 1983;140:734–9.
 - [57] Brooke P, Bullock R. Validation of a 6 item cognitive impairment test with a view to primary care usage. *Int J Geriatr Psychiatry* 1999;14:936–40.
 - [58] Tariq SH, Tumosa N, Chibnall JT, Perry MH 3rd, Morley JE. Comparison of the Saint Louis University mental status examination and the mini-mental state examination for detecting dementia and mild neurocognitive disorder—a pilot study. *Am J Geriatr Psychiatry* 2006;14:900–10.
 - [59] Pfeiffer E. A short portable mental status questionnaire for the assessment of organic brain deficit in elderly patients. *J Am Geriatr Soc* 1975;23:433–41.
 - [60] Kokmen E, Naessens JM, Offord KP. A short test of mental status: description and preliminary results. *Mayo Clin Proc* 1987;62:281–8.
 - [61] Inouye SK, Robison JT, Froehlich TE, Richardson ED. The time and change test: a simple screening test for dementia. *J Gerontol A Biol Sci Med Sci* 1998;53:M281–6.