

# ABDOMINAL COMPLETE ULTRASOUND PROTOCOL

UW Medicine

**BILLING CODE: UABDC**

**PATIENT PREP:** NPO at least 4 hours. Exceptions can be made for urgent and ER exams. Include in the report that the patient was not NPO for exam when relevant.

**\*\*A limited/focused exam may be done for the area of concern only in the following circumstances:**

*(If this occurs, billing code to be used is UABDL)*

- Inpatient: If other imaging has been performed within last 72 hours, a limited area of concern ultrasound may be done as long as there are no other findings are mentioned in report that should be evaluated. Examples: Eval renal mass, eval for cholecystitis, eval for Portal HTN with recent abdominal ultrasound.
- ER Patient: Perform full protocol unless order specifically states "Limited Abd to Eval xxx"
- Outpatient: Perform full protocol regardless of recent prior imaging.

**\*\*When screening for cirrhosis or HCC is requested, the following should be added to the exam:**

- Linear images of the capsule of LHL to show degree of nodularity
- Spleen images with measurement of size
- LIRADS should be completed in Viewpoint report. See additional info for LIRADS below.

**\*\*When rule out cholecystitis is requested, or patient is being seen for RUQ pain and has findings suspicious for cholecystitis, the following should be added to the exam:**

- Cystic artery (add billing code UORGDL)
- GB length measurement

**\*\*Organ donor and recipient screenings:**

- Pre-liver or kidney transplant donation screening: Complete abdomen, no extra images required
- Pre-renal transplant recipient screening: Complete abdomen, no extra images required
- Pre-liver transplant recipient screening: Perform spectral dopplers of liver vessels if requested. See abdomen doppler protocol.

**\*\*If a patient is coming in for pain, document within the indication section what the current status of pain is. For instance, how long the patient has had pain, if it's getting worse or better, and where exactly the pain is. Always take an image where the patient is hurting the most, labeled as area of pain. When describing the pain, write "per patient, ..." in the indication. Example: Per the patient, his pain is now in the RLQ and getting worse.**

**\*\*"Not well seen" to be stated if structure is not able to be completely evaluated. Include the reason why in relevant section of the report.**

**\*\*Any masses, cysts, stones or abnormalities should be measured in three dimensions and have a 2D picture and a color image documented. MFI should be routinely used to evaluate perfusion and low flow structures as needed. Measure the 2 largest or most worrisome masses/cysts in any given structure and comment on the presence of additional if relevant.**

**\*\*Cine clips to be added as needed for any abnormality seen.**

# IMAGES TO OBTAIN

## ABDOMINAL AORTA

- Sagittal image of proximal, mid, and distal aorta with measurements (outer to outer).
- Transverse image of proximal, mid, and distal aorta.
- Document any AAA in three dimensions, 2D and color Doppler.
- Document the iliac arteries sagittal and transverse if AAA is seen. Measure in sagittal.

## IVC

- Sagittal image of superior IVC.

## PANCREAS

- Transverse image of head, body (showing splenic vein) and tail.
- Sagittal image of head, body and tail.
- Transverse image of head showing porto-splenic confluence.
- Document and measure pancreatic duct if visible.
- Take image of "Pancreas Area" if not well seen.

## LIVER

### Left lobe – subcostal/epigastric approach

- Sagittal images:
  - Left lobe with left portal vein and ligamentum teres.
  - Left lobe with hepatic vein
  - Cine clip sweeping through LHL in sagittal from medial to lateral
- Transverse images:
  - Left lobe visualizing dome of liver
  - Left hepatic vein confluence into IVC
  - Left lobe with left portal vein
  - Cine clip sweeping through LHL in transverse from superior to inferior

### Caudate lobe – subcostal/epigastric approach

- Sagittal image of the caudate lobe.
- Transverse image of the caudate lobe.

### Right lobe - subcostal or intercostal approach

- Sagittal images:
  - Right hemidiaphragm to assess for pleural effusions and ringdown.
  - Right lobe and right portal vein.
  - Main interlobar fissure with gallbladder and CHD and MPV
  - Right lobe showing echotexture between liver and right kidney.
  - Right lobe and right kidney documenting approximate liver size. Measurement of length of the liver is not needed unless requested.
  - Cine clip sweeping through RHL in sagittal from medial to lateral
- Transverse images:
  - Right lobe to visualize dome of liver
  - Right liver with middle hepatic vein draining into IVC
  - Right lobe at right portal vein.
  - Right lobe and right kidney.
  - Cine clip sweeping through RHL in transverse from superior to inferior
  - 2D image through MPV
  - COLOR image of MPV showing patency and direction of flow

*NOTE: If lateral edge or dome of liver is not seen in its entirety with patient in supine position, turn patient in LLD or raise head of bed and repeat RHL cine clips to visualize better.*

## LIVER CAPSULE -FOR CIRRHOSIS AND HCC SCREENING ONLY

- Image of contour of liver capsule taken with linear transducer to assess for nodularity, preferably of the left lobe.

## BILE DUCTS:

- Sagittal image of CBD and CHD with measurements at the level of the porta hepatis. Color doppler should be used to distinguish ducts from vessels.
- If dilated, follow CBD as distal as possible to look for stones/mass and measure as distal as possible as well.
- Document and measure any intrahepatic bile duct dilatation with 2D and color imaging.

## GALLBLADDER

- Sagittal image of gallbladder.
- Transverse image of gallbladder.
- Measurement of gallbladder wall in sagittal section only. Do not include liver surface. If liver edge is edematous, try to measure free GB wall. Normal wall thickness is <3mm.
- Take image and cine through GB if there is any abnormality (stones, polyp, adenomyomatosis.) Take color image of any mass seen.
- LLD POSITION: Transverse and sagittal image of gallbladder with patient in LLD position to check for stones/polyps and mobility.
- If evaluating for cholecystitis, or if suspicious GB findings such as thickened wall or distension:
  - Evaluate for Murphy's sign and include in the report.
  - Gallbladder length measurement. Normal <8cm
  - Color doppler of gallbladder wall to assess for hypervascularity
  - Spectral doppler velocity of the cystic artery which can be seen within wall of anterior gallbladder.
    - Normal velocity <40cm/s
    - If artery cannot be visualized, include color doppler of area and comment in report that it could not be seen.
    - Add code UORGDL if cystic artery is sampled.



## RIGHT KIDNEY

- Sagittal image of right kidney in medial, middle, and lateral views.
- Sagittal measurement of right kidney.
- Transverse images of superior, mid, inferior right kidney.
- Demonstrate calculi, hydronephrosis or pelviectasis with a color image of the renal pelvis if present.
- **IF RENAL STONES ARE PRESENT:**
  - Measure renal stones in one largest dimension.
  - Demonstrate acoustic shadowing if possible.
  - Image with color to look for twinkle (can still be a stone if no twinkle shows).
- **IF HYDRONEPHROSIS, PELVIECTASIS, OR DILATED URETER PRESENT:**
  - Patient should void bladder and images should be taken to reassess degree of dilation with 2D and color images.

## SPLEEN

- Sagittal and transverse images through spleen
- Sagittal length measurement of spleen
- Sagittal image through left hemidiaphragm and spleen to assess for pleural effusion and ringdown.
- Compare echotexture between spleen and left kidney.
- Color Doppler image of any abnormality.

## LEFT KIDNEY

- Sagittal image of left kidney in medial, middle, and lateral views.
- Sagittal measurement of left kidney.
- Transverse images of superior, mid, inferior left kidney.
- Demonstrate hydronephrosis or pelviectasis with a color image of the renal pelvis if present.
- **IF RENAL STONES ARE PRESENT:**
  - Measure renal stones in one largest dimension.
  - Demonstrate acoustic shadowing if possible.
  - Image with color to look for twinkle (can still be a stone if no twinkle shows).
- **IF HYDRONEPHROSIS, PELVIECTASIS, OR DILATED URETER PRESENT:**
  - Patient should void bladder and images should be taken to reassess degree of dilation with 2D and color images.

## BLADDER

- Urinary bladder in sagittal and transverse planes.
- If bladder stones or debris are seen, demonstrate mobility by rolling patient.
- Demonstrate ureteral jets within bladder if hydronephrosis or dilated ureter is seen. (Observe for up to 5 mins).

## LOWER QUADRANTS:

- Document RLQ and LLQ (to check for ascites).

## ADDITIONAL IMAGES TO OBTAIN AS NEEDED

### AREA OF PAIN:

- Additional images in the area of pain using the curvilinear or linear transducer as need. Label images "Area of pain."

### CIRRHOSIS and HCC SCREENING:

- Image of contour of liver capsule taken with linear transducer to assess for nodularity, preferably of the left lobe.
- Spleen length and 2D images of spleen.
- **LI-RADS** : Viewpoint report should include the following LI-RADS statements:  
*NOTE: If lateral edge or dome of liver is not seen in its entirety with patient in supine position, turn patient in LLD or raise head of bed and RHL repeat cine clips to visualize better.*

1. LI-RADS visualization score:

VIS-A: no or minimal limitations

- A. limitations if any present are unlikely to affect sensitivity
- B. example: liver completely or near completely visualized, with little beam attenuation/shadowing

VIS-B: moderate limitations

- C. limitations that may obscure small (<10 mm) observations
- D. example: moderate beam attenuation with some portions of liver or diaphragm not visualized

VIS-C: severe limitations

- E. limitations that significantly limits detection of observations
- F. example: severe beam attenuation with most (>50%) of either liver lobe not visualized

2. LI-RADS observation description -Any mass seen should be referred to as an "observation" in the report with a description of the appearance. Do not label with the term mass. If it is clearly a cyst, it is okay to use the word "cyst."

### REPORTING EXAMPLE FOR LIVER SECTION WITH LIRADS:

LIVER:

Normal size, normal shape and contour, increased echogenicity.

LI-RADS visualization score: VIS-A: no or minimal limitations

LI-RADS observations:

- 1- Hypoechoic observation in LHL measuring 0.7 x 0.4 x 0.9 cm
- 2- Cystic/anechoic observation in LHL measuring 0.7 x 0.5 x 0.3 cm

### PRE-LIVER OR KIDNEY TRANSPLANT DONATION SCREENING:

- Complete Abdomen, no extra images required

### PRE-RENAL TRANSPLANT RECIPIENT SCREENING:

- Complete Abdomen, no extra images required

### PRE-LIVER TRANSPLANT RECIPIENT SCREENING:

- Perform spectral dopplers of liver vessels if requested. See ABDOMEN DOPPLER PROTOCOL.

## COMPLETE ABDOMEN ULTRASOUND PROTOCOL

IMAGE	MODE
IVC Sag	2D
Aorta Prox Sag w/ measurement	2D +
Aorta Prox Trv / IVC Trans	2D
Aorta Mid Sag w/ measurement	2D+
Aorta Mid Trans	2D
Aorta Dist Sag w/ measurement	2D+
Aorta Dist Trv	2D
Panc Trans H/B/T	2D
Panc Sag H/B/T	2D
<i>Panc Duct if dilated</i>	<i>2D +</i>
Left Liver Sag (medial portion)	2D
Left Liver Sag (lateral portion)	2D
Left Liver Sag M-L	Cine
Caudate Liver Sag	2D
Caudate Liver Trans	2D
Left Liver Trans (at hepatic vein)	2D
Left Liver Trans (at portal vein)	2D
Left Liver Trans S-I	Cine
Right Liver Trans (at hepatic veins)	2D
Right Liver Trans (at portal vein)	2D
Right Liver Trans at RK	2D
Right Liver Trans S-I	Cine
Right Liver Sag/Rt Chest	2D
Right Liver Sag	2D
Right Liver Sag / RK	2D
Right Liver Sag M-L	Cine
<i>Extra RHL sweeps in LLD or with head of bed elevated if needed</i>	<i>Cine</i>
MPV	2D
MPV w/ color	Color
GB Sag	
GB Wall w/measurement	2D+
GB Sag w/ color	Color
GB Trans	2D
GB Sag LLD	2D
GB Trans LLD	2D
<i>GB length if r/o chole</i>	<i>2D+</i>
<i>Cystic duct if r/o chole</i>	<i>Spectral</i>

IMAGE	MODE
CHD w/measurement and color	Color+
CBD w/measurement and color	Color+
Right Kidney Sag Mid	2D
Right Kidney Sag Mid w/ measurement	2D +
Right Kidney Sag Med	2D
Right Kidney Sag Lat	2D
Right Kidney Trans Sup	2D
Right Kidney Trans Mid	2D
Right Kidney Trans Inf	2D
Spleen Sag x2	2D x2
Spleen Sag w/ measurement	2D +
Spleen Sag/ Lt Chest	2D
Spleen Trans	2D
Spleen/LK	2D
Left Kidney Sag Mid	2D
Left Kidney Sag Mid w/ measurement	2D +
Left Kidney Sag Med	2D
Left Kidney Sag Lat	2D
Left Kidney Trans Sup	2D
Left Kidney Trans Mid	2D
Left Kidney Trans Inf	2D
Bladder Sag	2D
Bladder Trans	2D
<i>Bladder jets if hydro</i>	<i>Color</i>
RLQ	2D
LLQ	2D
<i>Liver capsule - Cirrhosis/HCC screening</i>	<i>Linear</i>
<i>MFI for lesions</i>	<i>MFI</i>

### ABDOMINAL COMPLETE PROTOCOL HISTORY

	Date	Changes made	By whom
Updated	07/09/21		Becky Marion
Updated	10/6/22	<p><b>ADDED:</b></p> <ul style="list-style-type: none"> <li>-Limited exam section when prior imaging within 72 hours</li> <li>- Cine sweeps of liver</li> <li>-LHL image w LHV</li> <li>-LI-RADS section</li> <li>-Cystic artery and gallbladder length for R/O chole cases and RUQ pain w suspicious GB findings</li> </ul> <p><b>REMOVED:</b></p> <ul style="list-style-type: none"> <li>-TRV IVC image</li> </ul>	<p>Protocol Meeting</p> <p><b>Attendees:</b></p> <ul style="list-style-type: none"> <li>Manjiri Dighe</li> <li>Shaun Bornemeier</li> <li>Dalene Edden</li> <li>Katie Toth</li> <li>Becky Marion</li> <li>Renee Betit Fitz</li> </ul>
Updated	6/1/2023	<p><b>-CHANGED:</b> Linear and LIRADS to be done for rule out cirrhosis cases as well as HCC</p> <p><b>-ADDED:</b> LIRADS - If it is clearly a cyst, it is okay to use the word "cyst."</p> <p><b>-ADDED:</b> MFI should be used to evaluate perfusion and low flow structures as needed on a routine basis.</p> <p><b>-CHANGED:</b> Measure the 2 largest or most worrisome masses/cysts in any given structure and comment on the presence of additional if relevant. (No longer 3)</p> <p><b>-CHANGED:</b> Sagittal image of CBD and CHD with measurements at the level of the porta hepatis. Color doppler should be used to distinguish ducts from vessels. (Multiple images with and without color and calipers no longer needed.)</p>	<p>Protocol Meeting</p> <p><b>Attendees:</b></p> <ul style="list-style-type: none"> <li>Manjiri Dighe</li> <li>Shaun Bornemeier</li> <li>Katie Toth</li> <li>Becky Marion</li> <li>Renee Betit Fitz</li> </ul>
Added	4/15/2024	Added Image list	Renee Betit Fitzgerald
Reviewed	5/23/2024	<p>Added: NOTE: If lateral edge or dome of liver is not seen in its entirety with patient in supine position, turn patient in LLD or raise head of bed and repeat RHL cine clips to visualize better.</p> <p>Updated LIRADS visualization scores</p>	<p>Protocol Meeting</p> <p><b>Attendees:</b></p> <ul style="list-style-type: none"> <li>Manjiri Dighe</li> <li>Dalene Edden</li> <li>Becky Marion</li> <li>Renee Betit Fitz</li> </ul>