

Table 1. Guidelines for Redating Based on Ultrasonography ↩

Gestational Age Range*	Method of Measurement	Discrepancy Between Ultrasound Dating and LMP Dating That Supports Redating
≤13 6/7 wk	CRL	
• ≤ 8 6/7 wk		More than 5 d
• 9 0/7 wk to 13 6/7 wk		More than 7 d
14 0/7 wk to 15 6/7 wk	BPD, HC, AC, FL	More than 7 d
16 0/7 wk to 21 6/7 wk	BPD, HC, AC, FL	More than 10 d
22 0/7 wk to 27 6/7 wk	BPD, HC, AC, FL	More than 14 d
28 0/7 wk and beyond†	BPD, HC, AC, FL	More than 21 d

Abbreviations: AC, abdominal circumference; BPD, biparietal diameter; CRL, crown–rump length; FL, femur length; HC, head circumference; LMP, last menstrual period.

*Based on LMP.

†Because of the risk of redating a small fetus that may be growth restricted, management decisions based on third-trimester ultrasonography alone are especially problematic and need to be guided by careful consideration of the entire clinical picture and close surveillance.

REFERENCE: [Methods for Estimating the Due Date | ACOG](#) . ACOG Committee Opinion Number 700 (Replaces Committee Opinion Number 611, October 2014. Reaffirmed 2022)